


# NICAM - Nonhydrostatic Icosahedral Atmospheric Model

## Masaki Satoh

(AORI, the Univ. of Tokyo/RIGC, JAMSTEC)



**1<sup>th</sup> Workshop on Dynamical Cores for Climate Models**

**December 14th-16th, 2011, Carlo V Castle - Lecce**

Group web page <http://nicam.jp>




# Contents

- **NICAM**
- **3.5km mesh NICAM simulation**
  - Tropical cyclone and synoptic scale waves
  - Cloud properties and evaluation
- **Athena Project**
- **NICAM-JMA collaboration**
- *Japan models application package*




# NICAM: Nonhydrostatic Icosahedral Atmospheric Model

- Development since 2000  
Tomita and Satoh(2005, *Fluid Dyn. Res.*), Satoh et al.(2008, *J. Comp. Phys.*)
- First global  $dx=3.5\text{km}$  run in 2004 using the Earth Simulator (JAMSTEC)  
Tomita et al.(2005, *Geophys. Res. Lett.*), Miura et al.(2007, *Science*)
- Toward higher resolution global simulation  
 $dx=1.7\text{km}, 880\text{m}, 440\text{m}$  using K-computer (10PF; Kobe,Riken,2012)
- International collaborations
  - Athena project (2009-10): COLA, NICS, ECMWF, JAMSTEC, Univ. of Tokyo
  - G8 ICOMEX (2011-): Germany, UK, France, US, Japan



# Nonhydrostatic Icosahedrall Atmospheric Model

- Icosahedral grid with spring dynamics smoothing
  - Tomita et al.(2001, *J. Comp. Phys.*)
  - Tomita et al.(2002, *J. Comp. Phys.*)
- Flux-form conservative nonhydrostatic scheme
  - Split-explicit time integration
  - mass and total energy & momentum conserving
    - Satoh (2002, *Mon. Wea. Rev.*)
    - Satoh (2003, *Mon. Wea. Rev.*)



Area distribution of cell: before and after

# Nonhydrostatic scheme

$$\frac{\partial}{\partial t} R + \nabla_h \cdot \mathbf{V}_h + \frac{\partial}{\partial \xi} \left( \frac{W}{G^{1/2}} + \mathbf{G}^3 \cdot \mathbf{V}_h \right) = 0$$

$$\frac{\partial}{\partial t} \mathbf{V}_h + \nabla_h P + \frac{\partial}{\partial \xi} (\mathbf{G}^3 P) = \mathbf{ADV}_h + \mathbf{F}_{Coriolis}$$

$$\frac{\partial}{\partial t} W + \frac{\partial}{\partial \xi} \left( \frac{P}{G^{1/2}} \right) + Rg = ADV_z + F_{z,Coriolis}$$

$$\frac{\partial}{\partial t} E_{total} + \nabla_h \cdot [(h + k + \Phi) \mathbf{V}_h] + \frac{\partial}{\partial \xi} \left[ (h + k + \Phi) \left( \frac{W}{G^{1/2}} + \mathbf{G}^3 \cdot \mathbf{V}_h \right) \right] = 0$$

## Prognostic variables

- density
- horizontal momentum  $R = G^{1/2} \rho$
- vertical momentum  $\mathbf{V}_h = G^{1/2} \rho \mathbf{v}_h$
- total energy  $W = G^{1/2} \rho w$

$$E_{total} = \rho G^{1/2} (e_{in} + k + \Phi)$$

## Metrics

$$G^{1/2} = \left( \frac{\partial z}{\partial \xi} \right)_{x,y}$$

$$\mathbf{G}^3 = (\nabla_h \xi)_z$$

$$\xi = \frac{H(z - z_s)}{H - z_s}$$

Satoh (2002, 2003)

# Model description

## ■ Dynamics

Governing equations	Fully compressible non-hydrostatic system
Spatial discretization	Finite Volume Method (Tomita et al. 2001,2002)
Horizontal grid configuration	Icosahedral grid
Vertical grid configuration	Lorenz grid
Topography	Terrain-following coordinate
Conservation	Total mass, total energy (Satoh 2002,2003)
Temporal scheme	Slow mode - explicit scheme ( RK2, RK3 ) Fast mode - Horizontal Explicit Vertical Implicit scheme
Advection scheme	Conservative and monotonic Miura (2004), Niwa et al.(2011)

## ■ Physics

Turbulence, surface flux	MYNN(Nakanishi and Niino 2004; Mellor & Yamada 2,2.5,3)/Louis(1979), Uno et al.(1995), Moon et al. (2007)
Radiation	MSTRNX (Sekiguchi and Nakajima, 2005)
Cloud physics	Kessler; Grabowski(1998); Lin et al.(1983); NSW6(Tomita 2008);NDW6(Seiki et al. 2011);WSM3-6 (Hong et al. 2004)
Subgrid convection	Prognostic AS, Kuo, Chikira (Chikira and Sugiyama 2010), Tiedtke (1989)

# **Global cloud-system resolving simulation of typhoon Fengshen (2008): comparison with ECMWF YOTC operational analysis data**

**Tomoe Nasuno**, Hiroyuki Yamada, Wataru Yanase,  
Akira T. Noda, and Masaki Satoh  
Email: [nasuno@jamstec.go.jp](mailto:nasuno@jamstec.go.jp)

## **Genesis of Typhoon Fengshen (2008) from an uptilted synoptic-scale disturbance: PALAU field experiment and global cloud-resolving simulation**


**Hiroyuki Yamada**, Tomoe Nasuno, Wataru Yanase,  
Ryuichi Shirooka, and Masaki Satoh  
Email: [yamada@jamstec.go.jp](mailto:yamada@jamstec.go.jp)

(2011, to be submitted)


12Z20JUN2008

NICAM JAMSTEC/AORI


NICAM 3.5 km mesh  
2008/06/20 12UTC



TC Fengshen



# Global cloud-resolving simulation of YOTC period #1



Horizontal grid spacing: **14 km, 3.5 km**

Vertical domain: 0 m ~ 38,000 m (40-levels)

Integration: **10 days from 00UTC 15 Jun 2008**


Initial conditions: **ECMWF YOTC Operational data**

NCEP final analysis (land surface, SST)

Boundary conditions: **slab ocean**



(nudging to Reynolds weekly SST)

➤ Fengshen formed on **17 Jun 2008**




# Genesis process of Typhoon Fengshen: mesoscale analysis


## observation (IR, radar)



## (a) GENESIS STAGE





## (b) INTENSIFICATION STAGE




# Impact of resolution on TC track

Fengshen track



Fengshen track is better simulated by 3.5 km NICAM than 14 km NICAM and JMA-GMS

Vertical velocity



Tendencies of vorticity

# Validation of cloud microphysical statistics simulated by a global cloud-resolving model with active satellite measurements

**Tempei Hashino<sup>1</sup>, Masaki Satoh<sup>1</sup>, Yuichiro Hagihara<sup>2</sup>,**  
**Takuji Kubota<sup>3</sup>, Toshihisa Matsui<sup>4</sup>, Tomoe Nasuno<sup>5</sup>, and**  
**Hajime Okamoto<sup>2</sup>**

Email: hashino@aori.u-tokyo.ac.jp

(2011, to be submitted)

<sup>1</sup>Atmosphere and Ocean Research Institute, The University of Tokyo

<sup>2</sup> Research Institute for Applied Mechanics, Kyushu University

<sup>3</sup>Japan Agency for Marine-earth Science and Technology


<sup>4</sup>NASA Goddard Space Flight Center

<sup>5</sup>Japan Agency for Marine-earth Science and Technology

# J-simulator (Joint Simulator for Satellite Sensors)


## by T. Hashino and the EarthCARE team

- Simulate EarthCARE (2014) observations from CRM outputs.
- Built on Satellite Data Simulator Unit (SDSU)  
Masunaga et al. (2010, BAMS)
- Extension at NASA/Goddard: Goddard-SDSU  
courtesy of T. Matsui & NASA GPM team




# Example 1: Tropical Cyclone

MTSAT IR  $T_b$  (10.8  $\mu\text{m}$ )



- OBS:
- Bright band exists.
  - High  $\beta_{532}$  above convective cores.
  - Overlap regions of C1 and C2 mask exist.



- NICAM:
- Bright band exists.
  - Low cloud top for C1, but high for C2.
  - Few overlap regions of C1 and C2 mask exist.

# The Athena Project

COLA, ECMWF, JAMSTEC, University of Tokyo, NICS, Cray

## List of experiments




Model/Exp.	Resolution	# Ensembles	Period	Notes
NICAM / Hindcasts	7 km	8	103 days	21 May - 30 Aug 2001 - 2009
IFS / Hindcasts	125 km 39 km 16 km	48	395 days	1 Nov - 30 Nov (following year) 1969 - 2007
IFS / Hindcasts	10 km	20		1 Nov - 30 Nov (following year) 1989 - 2007
IFS / Hindcasts	125 km 39 km 16 km 10 km	9	103 days	21 May - 30 Aug 2001 - 2009 NICAM analogs
IFS / Summer Ensembles	39 km 16 km	6	132 days	21 May - 30 Aug 2001 - 2009
IFS / Winter Ensembles	39 km 16 km	6	151 days	1 Nov - 30 Nov (following year)
IFS / AMIP	39 km 16 km	1	47 years	1961 - 2007
IFS / Time Slices	39 km 16 km	1	47 years	2071 - 2117

NICAM 7km  
dt=30sec

IFS 10km  
dt=450sec


<http://wxmaps.org/athena/home>,  
Kinter et al.(2011,submitted)

# Northward propagation of ISV in the Indian ocean



Time-latitude sections of anomalous 60–90E average surface precipitation (color) and zonal wind at 850 hPa (contour lines in the initial 52 days of NICAM (top) and IFS (middle) simulations in comparison with TRMM–3B42 and ERA–Interim data (bottom). The anomalies from the 8-year average (Fig. 2) are plotted. Contour intervals for zonal wind are  $2 \text{ m s}^{-1}$  (solid: positive, broken: negative). Zero contour lines are omitted.

# Most intensive Tropical cyclone structure





Distributions of 10m tangential wind (left panels; m s<sup>-1</sup>) and total column liquid water and ice (TCLWI; right panels; kg m<sup>-2</sup>) for the most intense TCs at the peak of their intensity from the NICAM simulation (panels a and d, labeled "NICAM"), the IFS 10--km simulation (panels b and e, labeled "T2047"), the IFS 39km simulation (panels c and f, labeled "T159"), respectively. Radius is 2° Contour interval is 3 m s<sup>-1</sup> for wind Dashed black contours in panels d, e and f show the radius of maximum winds for each case with respect to the center of the storm determined from the location of maximum vorticity at 925 hPa (1000 hPa for the IFS cases).

Kinter et al.(2011)

# Collaboration with JMA

- JMA's operational model is GSM, Global Spectral Model.
- Evolution of New Dynamics will be required for JMA's next-generation operational non-hydrostatic global model.
- Spectral model, grid-point model (Yin-Yang) and NICAM will be investigated at JMA.
- Comparison of NICAM and GSM is now conducted under collaboration with JMA.



# Testing NICAM and comparison with JMA-GSM





- NICAM  $dx=28\text{km}$ 
  - SR11000 10nodes
    - 150min for 72h forecast
    - Time step 150sec
- JMA-GSM  $dx=20\text{km}$ 
  - SR11000 60 nodes
    - 25min for 84h forecast
    - Time step 10min

# Sisters of NICAM: stretch-NICAM


(Tomita 2008; Satoh et al. 2008, J. Geophys.Res.)




Use of the stretched grid system



# Regional NICAM



# Plane-NICAM: double periodic model for LES



# Japan models application package

- Target Models
  - Climate models/Earth system models
    - MIROC: AORI/Univ. of Tokyo, JAMSTEC, NIES
    - MRI-GCM: MRI
  - Operational models
    - JMA-GSM: JMA
  - Global nonhydrostatic models
    - NICAM: AORI/Univ. of Tokyo, JAMSTEC, RIKEN/AICS
- Application package
  - Coupler: J-coupler, J-shell
  - Physics libraries

# Plan

## MIROC Atmosphere

- Climate
- Spectral, Grid (tracers)

## NICAM

- “Cloud-resolving”
- Geodesic, A-grid

A PetaFlops machine in Kobe, Japan




## Common atmos. core

- Climate/“Cloud-resolving”
- Geodesic, ZM-grid/A-grid(/Z-grid)

# Horizontal discretization


A



NICAM

- Geodesic
- A-grid
  - Tomita et al. (2001)
  - 2<sup>nd</sup>-order centered advection
  - piecewise linear method for tracers (Miura 2007)


Z



Common atmos. core (shallow water model at present)

- Geodesic
- ZM-grid(/Z-grid)
  - several changes from Ringler and Randall (2002)
  - 2<sup>nd</sup>-order upwind-biased advection for momentum
  - 3<sup>rd</sup>-order upwind-biased advection for mass
  - piecewise parabolic method for tracers


ZM




Courtesy of H. Miura (AORI)

# Grid staggering issues

- A-grid vs C-grid or others
  - A-grid: NICAM, NIM
  - C-grid: ICON, MPAS
  - Z-grid, ZM-grid, others
- Issues
  - Geostrophic flows
  - Divergent flows
  - Energy spectrum
  - Convergence
  - Numerical diffusions
- Personal view
  - Moist models will not converge until resolution becomes  $O(10m)$ .
  - If we stay around  $O(1km)$ , differences of grids are tuning or parameterization of cloud and turbulence schemes.
  - Differences between grids are resolvable scales. A vs C  $\sim 2:1$



# ppOpen-HPC: Open Source Infrastructure for Development and Execution of Large-Scale Scientific Applications on Post-Peta-Scale Supercomputers with Automatic Tuning (AT)



<http://www.jst.go.jp/kisoken/crest/en/areah22/2-03.html>

# Summary

- NICAM : 10 years history and beyond
- Good points of global nonhydrostatic models
  - Multiscale structure of cloud system
  - MJO, tropical cyclones
  - Cloud properties with cloud microphysics
  - KAKUSHIN & K-computer projects
- Physics
  - Use of satellite observations for evaluations
  - Sisters of NICAM(strech, regional, plane) are used for improvement of physics
  - Unified approach to global and regional models
- Japan models application package



**JMSJ Editorial Committee:**

**Chief Editor:** Satoh, Masaki;

**Co-Chief Editors:** Sato, Kaoru; Takemi, Tetsuya

**Editors:** Fujiwara, Masatomo; Inatsu, Masaru; Ishioka, Keiichi; Honda, Meiji; Horinouchi, Takeshi; Kato, Teruyuki; Kawashima, Masayuki; Kuji, Makoto; Kodama, Yasu-Masa; Masunaga, Hirohiko; Miyoshi, Takemasa; Nozawa, Toru; Shibata, Kiyotaka; Shige, Shoichi; Sudo, Kengo; Sugata, Seiji; Takegawa, Nobuyuki; Tanimoto, Youichi; Yukimoto, Seiji; Ueno, Kenichi; Wang, Yuqing; Watanabe, Masahiro

# Journal of the Meteorological Society of Japan (JMSJ)

JMSJ is an international English journal of the Meteorological Society of Japan. JMSJ publishes original research articles relating to meteorology. The articles are selected carefully by a thorough paper review based on the latest scientific knowledge.

The Meteorological Society of Japan was founded in 1882, for the wide information exchange of research, translation, introduction, and research plans relating to meteorology. The Society issued Series I of the Journal between 1882 and 1922. Since 1923 the society has been publishing Series II of the Journal.

*JMSJ, Since 1882*

visit <http://www.jstage.jst.go.jp/browse/jmsj>  
to submit papers and browse archives (*free access!*)