

Dynamical Exascale Entry Platform (DEEP)

Hendrik Merx
The Cyprus Institute
Max Planck Institute for Chemistry



DEEP Partners



The Cyprus Institute

Energy, Environment, and Water
Research Centre (EEWRC)

Science and Technology in Archaeology
Research Centre (STARC)

Computation-based Science and Technology
Research Centre (CaSToRC)

Scientific Applications

Seismic imaging

Brain simulation

Climate simulation

Space weather simulation


Computation fluid engineering

High-temperature superconductivity

Scientific Applications

Models of complex systems show
limited scalability for some processes
while being
highly-scalable for other processes

DEEP Hardware



Characteristics

Intel Xeon Cluster nodes

- provide I/O

- run fast mid-scalable model code parts

Intel Xeon Phi Booster nodes

- run standard Linux


- communicate directly through EXTOLL

- run self-contained full model code

- run highly-scalable model code parts

Dynamical creation of Cluster and Booster node groups

DEEP Software



ECHAM / MESSy

Operational Weather Forecast Model, Cycle 31

European Centre for Medium-Range Weather Forecast (ECMWF)
Reading, United Kingdom

ECMWF Model Hamburg (ECHAM)

Max Planck Institute for Meteorology (MPI-M), Hamburg, Germany

Modular Earth Subsystem Model (MESSy)

Max Planck Institute for Chemistry (MPI-C), Mainz, Germany

German Aerospace Centre (DLR), Oberpfaffenhofen, Germany

The Cyprus Institute (Cyl), Lefkosa, Cyprus

... and others

MESSy



MESSy

Links physical and chemical processes

Some models compute columns of air

Consumes ~ 80% of total run time

MESSy

bufly_physc	msbm_physc
d14co_physc	o3orig_physc
dradon_physc	photo_physc
e4chem_physc	plumegas_physc
gmxe_physc	psc_physc
h2o_physc	qbo_physc
hetchem_physc	satsims_physc
jval_physc	scav_physc
lnox_physc	sedi_physc
m7_physc	spe_physc
made_physc	spacenox_physc
meccal_physc	trexp_physc
mecca_physc	vahr_physc
mmforce_physc	

MECCA

Models homogeneous gas-phase chemistry

Solves a system of differential equations


System is integrated by adaptive integrator

Change in light intensity leads to stiff system

Run time varies by two orders of magnitude

Consumes ~ 80 % of MESSy run time

Atmospheric Chemistry



OmpSs

```
REAL(dp) :: zmr(ntrac_gp) ! tracer mixing ratio

level_loop: DO jk = 1, nlev

    !$omp task inout(pxtml(:,jk,:), pxtte(:,jk,:)) &
    !$omp      private(zmr) &
    !$omp      output(c(:,jk,:))

    kproma_loop: DO jp = 1, kproma

        CALL update_physc ! update rate coefficients

        ! estimate tracer mixing ratio before chemistry
        zmr(:) = pxtml(jp,jk,:) + pxtte(jp,jk,:) * time_step_len

        CALL kpp_integrate(time_step_len) ! works on x(jp,jk,:)
```

Advantages

Task off-loading provides an additional run-time degree of freedom for parallelisation

Increase threads for grid point calculations,
decrease tasks for spectral calculations,
improving implicit load balancing

From multi-core to many-core architectures
due to reduced memory constraints in multi-threaded parallelisation

Status



Thank you