

Cohort Definition Examples

Contents

1	Type 2 diabetes mellitus	1
2	Type 1 diabetes mellitus	3
3	Atrial Fibrillation	4

```
library(Capr)

# Function to test that cohort compiles
assertCohortCompiles <- function(cohortDefinition) {
  sql <- CirceR::cohortExpressionFromJson(as.json(ch))
  sql <- CirceR::buildCohortQuery(sql, options = CirceR::createGenerateOptions(generateStats = TRUE))

  stopifnot(nchar(sql) > 1)
  invisible(NULL)
}
```

This vignette provides a number of example cohorts built with Capr from the OHDSI *Phenotype Phebruary* event.

1 Type 2 diabetes mellitus

Persons with new type 2 diabetes mellitus at first diagnosis

<https://atlas-phenotype.ohdsi.org/#/cohortdefinition/88/definition>

```
library(Capr)

cs0 <- cs(descendants(443238, 201820, 442793),
          descendants(exclude(195771, 201254, 435216, 761051, 4058243, 40484648)),
          name = "Type 2 diabetes mellitus (diabetes mellitus excluding T1DM and secondary)")

ch <- cohort(
  entry = entry(
    conditionOccurrence(cs0),
    observationWindow = continuousObservation(priorDays = 365)
  ),
  exit = exit(
    endStrategy = observationExit()
  )
)
```

Persons with new type 2 diabetes and no prior T1DM or secondary diabetes

<https://atlas-phenotype.ohdsi.org/#/cohortdefinition/89/export>

```

library(Capr)

cs0 <- cs(descendants(443238, 201820, 442793),
          descendants(exclude(195771, 201254, 435216, 761051, 4058243, 40484648)),
          name = "Type 2 diabetes mellitus (diabetes mellitus excluding T1DM and secondary)")

cs1 <- cs(descendants(201254, 435216, 40484648),
          name = "Type 1 diabetes mellitus")

cs2 <- cs(descendants(195771),
          name = "Secondary diabetes mellitus")

ch <- cohort(
  entry = entry(
    conditionOccurrence(cs0),
    observationWindow = continuousObservation(priorDays = 365)
  ),
  attrition = attrition(
    't1d' = withAll(
      exactly(0, conditionOccurrence(cs1), duringInterval(eventStarts(-Inf, 0)))
    ),
    'secondaryDiabetes' = withAll(
      exactly(0, conditionOccurrence(cs2), duringInterval(eventStarts(-Inf, 0)))
    )
  ),
  exit = exit(
    endStrategy = observationExit()
  )
)

```

Persons with new type 2 diabetes mellitus at first dx rx or lab

<https://atlas-phenotype.ohdsi.org/#/cohortdefinition/90>

```

library(Capr)

cs0 <- cs(descendants(443238, 201820, 442793),
          descendants(exclude(195771, 201254, 435216, 761051, 4058243, 40484648)),
          name = "Type 2 diabetes mellitus (diabetes mellitus excluding T1DM and secondary)")

cs1 <- cs(descendants(201254, 435216, 40484648),
          name = "Type 1 diabetes mellitus")

cs2 <- cs(descendants(195771),
          name = "Secondary diabetes mellitus")

cs3 <- cs(descendants(4184637, 37059902),
          name = "Hemoglobin A1c (HbA1c) measurements")

cs4 <- cs(descendants(21600744),
          name = "Drugs for diabetes except insulin")

ch <- cohort(
  entry = entry(

```

```

conditionOccurrence(cs0),
drugExposure(cs4),
measurement(cs3, valueAsNumber(bt(6.5, 30)), unit("%")),
measurement(cs3, valueAsNumber(bt(48, 99)), unit("mmol/mol")),
observationWindow = continuousObservation(priorDays = 365)
),
attrition = attrition(
  'no T1D' = withAll(
    exactly(0, conditionOccurrence(cs1), duringInterval(eventStarts(-Inf, 0)))
  ),
  'no secondary diabetes' = withAll(
    exactly(0, conditionOccurrence(cs2), duringInterval(eventStarts(-Inf, 0)))
  )
),
exit = exit(
  endStrategy = observationExit()
)
)

```

2 Type 1 diabetes mellitus

Persons with new type 1 diabetes

<https://atlas-phenotype.ohdsi.org/#/cohortdefinition/92>

```

cs0 <- cs(descendants(195771),
  name = "Type 1 diabetes mellitus")

ch <- cohort(
  entry = entry(
    conditionOccurrence(cs0),
    observationWindow = continuousObservation(priorDays = 365)
  )
)

```

```
assertCohortCompiles(ch)
```

Persons with new type 1 diabetes and no prior T2DM or secondary diabetes

<https://atlas-phenotype.ohdsi.org/#/cohortdefinition/91>

```

cs0 <- cs(descendants(443238, 201820, 442793),
  descendants(exclude(195771, 201254, 435216, 761051, 4058243, 40484648)),
  name = "Type 2 diabetes mellitus (diabetes mellitus excluding T1DM and secondary)")

cs1 <- cs(descendants(201254, 435216, 40484648),
  name = "Type 1 diabetes mellitus")

cs2 <- cs(descendants(195771),
  name = "Secondary diabetes mellitus")

ch <- cohort(
  entry = entry(
    conditionOccurrence(cs1),
    observationWindow = continuousObservation(priorDays = 365)
  )
)

```

```

),
attrition = attrition(
  "no prior T2DM" = withAll(exactly(0, conditionOccurrence(cs0), duringInterval(eventStarts(-Inf, 0))),
  "no prior secondary T1DM" = withAll(exactly(0, conditionOccurrence(cs2), duringInterval(eventStarts
)
)
)

```

3 Atrial Fibrillation

Persons with atrial fibrillation per Wharton et al 2021

<https://atlas-phenotype.ohdsi.org/#/cohortdefinition/93>

```

cs0 <- cs(descendants(313217),
  name = "Atrial fibrillation")

ch <- cohort(conditionOccurrence(cs0))

```

Persons with atrial fibrillation per Subramanya et al 2021

<https://atlas-phenotype.ohdsi.org/#/cohortdefinition/94>

```

afib <- cs(descendants(313217),
  name = "Atrial fibrillation")

ip <- cs(descendants(262, 9201),
  name = "Inpatient or inpatient ER visit")

op <- cs(descendants(9202, 9203),
  name = "Outpatient or ER visit")

ch <- cohort(
  entry = entry(
    conditionOccurrence(afib,
      nestedWithAny(
        atLeast(1, visit(ip), duringInterval(eventStarts(-Inf, 0), eventEnds(0, Inf))),
        nestedWithAll(
          atLeast(1, visit(op, duringInterval(eventStarts(-Inf, 0), eventEnds(0, Inf))),
            nestedWithAll(
              atLeast(1, conditionOccurrence(afib, duringInterval(eventStarts(7, 365))),
                nestedWithAll(
                  atLeast(1, visit(op, duringInterval(eventStarts(-Inf, 0), eventEnds(0, Inf))))
                )
              )
            )
          )
        )
      )
    )
  )
)

```