

Building and evaluation of a PBPK model for antibody CDA1 in healthy adults

Version	1.0-OSP12.3
based on <i>Model Snapshot and Evaluation Plan</i>	https://github.com/Open-Systems-Pharmacology/CDA1-Model/releases/tag/v1.0
OSP Version	12.3
Qualification Framework Version	3.7

This evaluation report and the corresponding PK-Sim project file are filed at:

<https://github.com/Open-Systems-Pharmacology/OSP-PBPK-Model-Library/>

Table of Contents

- 1 Introduction
- 2 Methods
 - 2.1 Modeling Strategy
 - 2.2 Data
 - 2.2.1 In vitro / physico-chemical Data
 - 2.2.2 PK Data
 - 2.3 Model Parameters and Assumptions
- 3 Results and Discussion
 - 3.1 Final input parameters
 - 3.2 Diagnostics Plots
 - 3.3 Concentration-Time Profiles
- 4 Conclusion
- 5 References

1 Introduction

CDA1 is a human monoclonal antibody (IgG1) against the toxin A of *Clostridium difficile*.

CDA1 shows a pharmacokinetic behavior which is typical for an antibody without endogenous target. The plasma concentration–time profiles after i.v. infusion of 5, 10 and 20 mg/kg CDA1 in healthy adults (Taylor 2008) were used together with pharmacokinetic (PK) data from 5 other compounds to identify unknown parameters during the development of the generic large molecule physiologically based pharmacokinetic (PBPK) model in PK-Sim (Niederalt 2018).

The herein presented evaluation report evaluates the performance of the PBPK model for CDA1 in healthy adults for the PK data used for the development of the generic large molecule model in PK-Sim.

The presented CDA1 PBPK model as well as the respective evaluation plan and evaluation report are provided open-source (<https://github.com/Open-Systems-Pharmacology/CDA1-Model>).

2 Methods

2.1 Modeling Strategy

The development of the large molecule PBPK model in PK-Sim® has previously been described by Niederalt et al. ([Niederalt 2018](#)). In short, the model was built as an extension of the PK-Sim® model for small molecules incorporating (i) the two-pore formalism for drug extravasation from blood plasma to interstitial space, (ii) lymph flow, (iii) endosomal clearance and (iv) protection from endosomal clearance by neonatal Fc receptor (FcRn) mediated recycling.

For model development and evaluation, PK data were used from compounds with a wide range of solute radii and from different species. The PK data used for parameter estimation were from the following compounds: antibody–drug conjugate BAY 79-4620 in mice (Bayer in house data), antibody 7E3 in wild-type and FcRn knockout mice ([Garg 2007](#), [Garg2009](#)), domain antibody dAb2 in mice ([Sepp 2015](#)), antibodies MEDI-524 and MEDI-524-YTE in monkeys ([Dall'Acqua 2006](#)), and antibody CDA1 in humans ([Taylor 2008](#)). The PK data used for model evaluation were from inulin in rats ([Tsuji1983](#)) and tefibazumab in humans ([Reilly 2005](#)).

The PBPK model including the estimated physiological parameters as described by Niederalt et al. ([Niederalt 2018](#)) is available in the Open Systems Pharmacology Suite from version 7.1 onwards.

This evaluation report focuses on the PBPK model for the antibody CDA1.

Details about input data (physicochemical, *in vitro* and PK) can be found in [Section 2.2](#).

Details about the structural model and its parameters can be found in [Section 2.3](#).

2.2 Data

2.2.1 In vitro / physico-chemical Data

A literature search was performed to collect available information on physicochemical properties of CDA1. The obtained information from literature is summarized in the table below.

Parameter	Unit	Value	Source	Description
MW	g/mol	150000	Lobo 2004	Molecular weight
r	nm	5.34	Taylor 1984	Hydrodynamic solute radius
Kd (FcRn)	μM	0.63	Zhou 2003	Dissociation constant for binding of a human IgG1 antibody to human FcRn at pH 6

2.2.2 PK Data

Published clinical PK data on CDA1 in healthy adults were used.

Publication	Description
Taylor2008	The plasma concentration–time profiles after single i.v. infusion of 5, 10 and 20 mg/kg CDA1 in healthy adults were used. The data for the dosages 0.3 and 1 mg/kg were not used since the PK data could not be read with sufficient accuracy from the published figure.

2.3 Model Parameters and Assumptions

2.3.1 Absorption

There is no absorption process since CDA1 was administered intravenously.

2.3.2 Distribution

The standard lymph and fluid recirculation flow rates and the standard vascular properties of the different tissues (hydraulic conductivity, pore radii, fraction of flow via large pores) from PK-Sim were used. CDA1, among other compounds, has been used to identify these lymph and fluid recirculation flow rates used in PK-Sim ([Niederalt 2018](#)).

2.3.3 Metabolism and Elimination

The FcRn mediated clearance present in the standard PK-Sim model was used as only clearance process. The standard physiological parameters related to FcRn mediated clearance were used (rate constants for endosomal uptake and recycling, association rate constant for FcRn binding and concentration of FcRn in the endosomal space). CDA1, among other compounds, has been used to identify these parameters using literature values for the drug affinities to FcRn in the endosomal space ([Niederalt 2018](#)).

2.3.4 Automated Parameter Identification

No drug specific parameters were fitted. CDA1, among other compounds, has been used to develop the model for proteins and large molecules in PK-Sim ([Niederalt 2018](#)).

3 Results and Discussion

The PBPK model for CDA1 was evaluated with clinical PK data in healthy adults.

These PK data have been used together with PK data from 5 other compounds to simultaneously identify parameters during the development of the generic model for proteins and large molecules in PK-Sim ([Niederalt 2018](#)).

The next sections show:

1. the final model parameters for the building blocks: [Section 3.1](#).
2. the overall goodness of fit: [Section 3.2](#).
3. simulated vs. observed concentration-time profiles for the clinical studies used for model building and for model verification: [Section 3.3](#).

3.1 Final input parameters

The compound parameter values of the final PBPK model are illustrated below.

Compound: CDA1

Parameters

Name	Value	Value Origin	Alternative	Default
Solubility at reference pH	999 mg/l	Other-/Dummy value not used in the simulation	Measurement	True
Reference pH	7	Other-/Dummy value not used in the simulation	Measurement	True
Lipophilicity	-5 Log Units	Other-/Dummy value not used in the simulation	Measurement	True
Fraction unbound (plasma, reference value)	1	Other-Assumption	Measurement	True
Is small molecule	No			
Molecular weight	150000 g/mol	Publication-Lobo2004		
Plasma protein binding partner	Unknown			
Radius (solute)	5.34 nm	Publication-Taylor1984		
Kd (FcRn) in endosomal space	0.63 μ mol/l	Publication-Zhou2003		

Calculation methods

Name	Value
Partition coefficients	PK-Sim Standard
Cellular permeabilities	PK-Sim Standard

Processes

3.2 Diagnostics Plots

Below you find the goodness-of-fit visual diagnostic plots for the PBPK model performance of all data used presented in [Section 2.2.2](#).

The first plot shows observed versus simulated plasma concentration, the second weighted residuals versus time.

Table 3-1: GMFE for Goodness of fit plot for concentration in plasma

Group	GMFE
10 mg/kg IV	1.06
20 mg/kg IV	1.14
5 mg/kg IV	1.02
All	1.08




Figure 3-1: Goodness of fit plot for concentration in plasma




Figure 3-2: Goodness of fit plot for concentration in plasma

3.3 Concentration-Time Profiles

Simulated versus observed concentration-time profiles of all data listed in [Section 2.2.2](#) are presented below.




Figure 3-3: Plasma concentration - Dose 5 mg/kg (log scale)




Figure 3-4: Plasma concentration - Dose 5 mg/kg (linear scale)




Figure 3-5: Plasma concentration - Dose 10 mg/kg (log scale)




Figure 3-6: Plasma concentration - Dose 10 mg/kg (linear scale)




Figure 3-7: Plasma concentration - Dose 20 mg/kg (log scale)




Figure 3-8: Plasma concentration - Dose 20 mg/kg (linear scale)

4 Conclusion

The herein presented PBPK model overall adequately describes the pharmacokinetics of CDA1 in healthy humans. The initial plasma concentrations are slightly underestimated especially for the higher dose.

The PK data had been used during the development of the generic large molecule PBPK model in PK-Sim ([Niederalt 2018](#)) together with PK data from 5 other compounds (7E3, BAY 79-4620, dAb2, MEDI-524 & MEDI-524-YTE).

5 References

- Dall'Acqua 2006** Dall'Acqua WF, Kiener PA, Wu H. Properties of human IgG1s engineered for enhanced binding to the neonatal Fc receptor (FcRn). *J Biol Chem*. 2006 Aug; 281(33):23514-23524. doi: 10.1074/jbc.M604292200.
- Garg 2007** Garg A, Balthasar JP. Physiologically-based pharmacokinetic (PBPK) model to predict IgG tissue kinetics in wild-type and FcRn-knockout mice. *J Pharmacokinet Pharmacodyn*. 2007 Jul; 34(5):687-709. doi: 10.1007/s10928-007-9065-1.
- Garg 2009** Garg A, Balthasar J. Investigation of the influence of FcRn on the distribution of IgG to the brain. *AAPS J*. 2009 July; 11(3):553-557. doi: 10.1208/s12248-009-9129-9.
- Lobo 2004** Lobo ED, Hansen R J, Balthasar JP. Antibody pharmacokinetics and pharmacodynamics. *J Pharm Sci*. 2004 Nov;93(11):2645-2668. doi: 10.1002/jps.20178.
- Niederalt 2018** Niederalt C, Kuepfer L, Solodenko J, Eissing T, Siegmund HU, Block M, Willmann S, Lippert J. A generic whole body physiologically based pharmacokinetic model for therapeutic proteins in PK-Sim. *J Pharmacokinet Pharmacodyn*. 2018 Apr;45(2):235-257. doi: 10.1007/s10928-017-9559-4.
- Reilly 2005** Reilly S, Wenzel E, Reynolds L, Bennett B, Patti JM, Hetherington S. Open-label, dose escalation study of the safety and pharmacokinetic profile of tefibazumab in healthy volunteers. *Antimicrob Agents Chemother*. 2005 Mar;49(3):959–962. doi: 10.1128/AAC.49.3.959-962.2005.
- Sepp 2015** Sepp A, Berges A, Sanderson A, Meno-Tetang G. Development of a physiologically based pharmacokinetic model for a domain antibody in mice using the two-pore theory. *J Pharmacokinet Pharmacodyn*. 2015 Jan;42(2):97-109. doi: 10.1007/s10928-014-9402-0.
- Taylor 1984** Taylor AE, Granger DN. Exchange of macromolecules across the microcirculation. *Handbook of Physiology - Cardiovascular System. Microcirculation* (Eds. Renkin EM and Michel CC. Bethesda, MD, American Physiological Society). 1984; Vol. 4(Pt 2):467–520.
- Taylor 2008** Taylor CP, Tummala S, Molrine D, Davidson L, Farrell RJ, Lembo A, Hibberd PL, Lowy I, Kelly CP. Open-label, dose escalation phase I study in healthy volunteers to evaluate the safety and pharmacokinetics of a human monoclonal antibody to Clostridium difficile toxin A. *Vaccine*. 2008 Jun;26(27-28):3404–3409. doi: 10.1016/j.vaccine.2008.04.042.
- Tsuji 1983** Tsuji A, Yoshikawa T, Nishide K, Minami H, Kimura M, Nakashima E, Terasaki T, Miyamoto E, Nightingale CH, Yamana T. Physiologically based pharmacokinetic model for beta-lactam antibiotics I: tissue distribution and elimination in rats. *J Pharm Sci*. 1983 Nov;72(11):1239-1252. doi: 10.1002/jps.2600721103.
- Zhou 2003** Zhou J, Johnson JE, Ghetie V, Ober RJ, Ward ES. Generation of mutated variants of the human form of the MHC class I-related receptor, FcRn, with increased affinity for mouse immunoglobulin G. *J Mol Biol*. 2003 Sep;332(4):901-913. doi: 10.1016/s0022-2836(03)00952-5.