

8. BIBLIOGRAPHY

Over 300 references have been collected on remote sensing, seafloor mapping and habitat classification. Searches include online database and internets sources, published literature, technical reports and trade journals, and conference proceedings. Full references with abstracts (as available) are digitally archived in EndNote database.

- Able, K. W., *et al.* (1987). Sidescan sonar as a tool for detection of demersal fish habitats. *Fishery Bulletin* **85**(4): 725-737.
- Able, K. W., *et al.* (1993). Temporal and spatial variation in habitat characteristics of tilefish (*Lopholatilus chamaeleonticeps*) off the east coast of Florida. *Bulletin of Marine Science* **53**(3): 1013-1026.
- Able, K.W., Grimes, C.B., Twichell, and Jones, R.S., 1995. Side-scan sonar as a tool for determination of demersal fish habitat use patterns on the continental shelf: *in* O'Connell, T. and Wakefield, W., eds., Applications of side-scan sonar and laser-line systems in Fisheries Research: Alaska Dept. Fish and Game Special Publication No. 9, p. 8-9.
- Able, K.W., Twichell, D.C., Grimes, C.B., and Jones, R.S., 1987. Sidescan sonar as a tool for detection of demersal fish habitats: *Fish. Bull.*, U.S. 85: p. 725-736.
- Alais, P., *et al.* (1994). A high resolution sidescan sonar. *OCEANS, 94*(HELD IN CONJUNCTION WITH OSATES 94): RESERVATION-PROCEEDINGS.
- Alexandrou, D., & D. Pantzartzis (1990). Seafloor classification with neural networks. *OCEANS '90: ENGINEERING IN THE OCEAN ENVIRONMENT. CONFERENCE PROCEEDINGS.*
- Allen, J. L. R. (1984). Sedimentary structures: their character and physical basis. *Developments in Sedimentology*, vol. II, Amsterdam: Elsevier: 663 pp.
- Anger C.D., Mah, S., Babey, S.K., 1994, Technological enhancements to the compact airborne spectrographic imager (casi), in Proceedings of the First International Airborne Remote Sensing Conference and Exhibition, Strasbourg, France, Vol. II pp. 205-213
- Anger, C.D., S. K. Babey, and R. J. Adamson, 1990, A New Approach to Imaging Spectroscopy, SPIE Proceedings, Imaging Spectroscopy of the Terrestrial Environment, 1298: 72 - 86.
- Annual International Symposium of the North American Lake Management Society, Toronto, ON (Canada), 6-11 Nov 1995, 11(2): 185.
- Anon (1994). Interpreting sonar records. *International Underwater Systems Design [INT. UNDERWAT. SYST. DES.]*, **16**(3): 13-18.
- Anon (1995). Mapping fresh-water sediments with an acoustic bottom-discrimination system. *Lake and Reservoir Management [LAKE RESERV. MANAGE.]*

- Anon (1996). Seafloor mapping in the West, Southwest and South Pacific: Results and applications. Marine Geophysical Researches **18**: 2-4.
- Anstee, J.M., D.L.B. Jupp and G.T. Byrne, The Shallow Benthic Cover Map and Optical Water Quality of Port Phillip Bay, presented at the 4th International Conference on Remote Sensing for Marine and Coastal Environments, Orlando, Florida, USA, 1997.
- Apsey, B. F. (1998). Multibeam survey of damaged reef. University of Miami to pay for coral damage. Hydro International [Hydro Int.], **2**(1): 48-49.
- Augustin, J. M., *et al.* (1996). Contribution of the multibeam acoustic imagery to the exploration of the sea-bottom. Examples of SOPACMAPS 3 and ZoNeCo 1 cruises. SEAFLOOR MAPPING THE WEST, SOUTHWEST AND SOUTH PACIFIC: RESULTS AND APPLICATIONS, **18**: 2-4.
- Auster, P.J., 1991, Malatesta, R.J., LaRosa, S.C., Cooper, R.A. and Stewart, L.L., 1991. Microhabitat utilization by the megafaunal assemblages at a low relief outer continental shelf site - Middle Atlantic Bight, USA. *J. Northw. Atl. Fish. Sci.*, v. 11, p. 59-69
- Auster, P.J., Lewis, R.S., Wahle, L.C., Babb, I.G., and Malatesta, R.J., 1995. The use of side-scan sonar for landscape approaches to habitat mapping: *in* O'Connell, T. and Wakefield, W., eds., Applications of side-scan sonar and laser-line systems in Fisheries Research: Alaska Dept. Fish and Game Special Publication No. 9, p.1-7.
- Babey, S.K., Anger, C.D., 1989. A compact airborne spectrographic imager (casi), in Proceedings of IGARSS, Vol.2, pp. 1028-1031.
- Babey, S.K., Anger, C.D., 1993. Compact airborne spectrographic imager (casi): A progress review, in Proceedings of the SPIE Conference, Orlando, Florida, SPIE Vol. 1937 pp. 152-163.
- Baltzer, A.,P. Cochonat, & D. J. W. Piper (1994). In situ geotechnical characterization of sediments on the Nova Scotian slope, eastern Canadian continental margin. Marine Geology **120**(3-4): 291-308.
- Barans, C. A., & D. V. Holliday (1983). A practical technique for assessing some snapper/grouper stocks. Bulletin of Marine Science, **33**(1): 176-181.
- Barnhard, W. A., *et al.* (1998). Mapping the Gulf of Maine with side-scan sonar: A new bottom-type classification for complex seafloors. Journal of Coastal Research **14**(2): 646-659.
- Baudry, N., & S. Calmant (1996). Seafloor mapping from high-density satellite altimetry. SEAFLOOR MAPPING THE WEST, SOUTHWEST AND SOUTH PACIFIC: RESULTS AND APPLICATIONS, **18**: 2-4.
- Belderson, R. H., *et al.* (1972). Sonographs of the seafloor. Amsterdam: Elsevier: 185 pp.
- Belderson, R. H., *et al.* (1982). Bedforms. Offshore Tidal Sands, Process and Deposits, A.H. Stride: 27-57.

- Benaka, L. R. (ed) 1999. Fish Habitat: Essential Fish Habitat and Rehabilitation. American Fisheries Society Symposium 22. Proceedings of the Sea Grant Symposium on Fish Habitat: "Essential Fish Habitat" and Rehabilitation, Hartford, Connecticut, 26-27 August, 1998. 459 pp.
- Bird, R. T., *et al.* (1996). Merged GLORIA sidescan and hydrosweep pseudo-sidescan: Processing and creation of digital mosaics. Marine Geophysical Researches **18**(6): 651-661.
- Blackinton, J. G., (1991). Bathymetry resolution, precision, and accuracy considerations for swath bathymetry mapping sonar systems. IEEE Proc., : 550-556.
- Blake SG, Jupp DLB, Byrne GT. The potential use of high-resolution casi data to determine sedimentation patterns and primary productivity throughout the Whitsunday Islands region, Great Barrier Reef. Paper presented to 7th Australian Remote Sensing Conference. 1994.
- Blondel, P., Bramley J. M. 1997. Handbook of Seafloor Sonar Imagery. Wiley, New York. 314 pp.
- Bobbitt, A. M., *et al.* (1997). GIS analysis of remotely sensed and field observation oceanographic data. Marine Geodesy, **20**(2-3): 153-161.
- Bodur, M. N., & M. Ergin (1992). Holocene sedimentation patterns and bedforms in the wave-current-dominated nearshore waters of eastern Mersin Bay (Eastern Mediterranean). Marine Geology **108**(1): 73-93.
- Booth, J., Hay, D. E., Truscott, J. 1996. Standard methods for sampling resources and habitats in coastal subtidal regions of British Columbia: Part 1: Review of mapping and preliminary recommendations. Canadian Technical Report of Fisheries and Aquatic Sciences 2118. Fisheries and Oceans, Canada. 53 pp.
- Borstad, G. (1995). Compact Airborne Spectrographic Imager. Remote Sensing Techniques for Subtidal Classification, workshop, Institute of Ocean Sciences, Sydney. B. C.: 18-20.
- Borstad, G.A., 1992, Ecosystem surveillance and monitoring with a portable airborne imaging spectrometer system, in the Proceedings of the First Thematic Conference on Remote Sensing for Marine and Coastal Environments, New Orleans, Louisiana, pp. 883-892.
- Borstad, G.A., Hill, D.A., 1989, Using visible range imaging spectrometers to map ocean phenomena, in the proceedings of the SPIE Conference on Advanced Optical Instrumentation for Remote Sensing of the Earth's Surface from Space, Paris, France, Vol. 1129, pp. 130-136.
- Borstad, G.A., Kerr, R.C., Zacharias, M., 1994, Monitoring near shore water quality and mapping of coastal areas with a small airborne system and GIS, in the Proceedings of the Second Thematic Conference on Remote Sensing for Marine and Coastal Environments, New Orleans, Louisiana, Vol. II PP. 51- 56.

- Bourillet, J. F., *et al.* (1996). Swath mapping system processing: Bathymetry and cartography. SEAFLOOR MAPPING THE WEST, SOUTHWEST AND SOUTH PACIFIC: RESULTS AND APPLICATIONS, **18**: 2-4.
- Brekhovskikh, L. M., Yu. P. Lysanov (1991). Fundamentals of Ocean Acoustics. 2nd edition, Berlin: Springer-Verlag, 270 pp.
- Bretz, C.K., R.G. Kvitek, P.J. Iampietro. (1998). A case study in the application of spatial data acquisition tools and multimedia GIS for environmental monitoring and management in Antarctica. EOM.7:11-13.
- Broadus, M., *et al.* (1997). New survey systems and technology for the littoral environment. Marine Technology Society Journal [MAR. TECHNOL. SOC. J.], **31**(1): 21-30.
- Brown, J.L. Linnett, & I. McColl (1994). Synthesis of sonar data using three dimensional computer models. UNDERWATER ACOUSTICS. PROCEEDINGS OF THE 2ND EUROPEAN CONFERENCE ON UNDERWATER ACOUSTICS, **1**: 413-418.
- Bruce, E. M., I. G. Eliot, & D. J. Milton (1997). Method for assessing the thematic and positional accuracy of seagrass mapping. Marine Geodesy, **20**(2-3): 175-193.
- Caiti, A., & T. Parisini (1994). Mapping ocean sediments by RBF networks. IEEE Journal of Oceanic Engineering **19**(4): 577-582.
- Calmant, S., & N. Baudry (1996). Modeling bathymetry by inverting satellite altimetry data: A review. SEAFLOOR MAPPING THE WEST, SOUTHWEST AND SOUTH PACIFIC: RESULTS AND APPLICATIONS, **18**: 2-4.
- Carlson, R. F. (1992). Latest developments in geodetically correct multibeam bathymetry, side scan imagery, and sub-bottom profiling. MTS '92: GLOBAL OCEAN PARTNERSHIP. PROCEEDINGS., MTS, WASHINGTON, DC.
- Carr, H. A., & J. P. Fish (1992). The application of side scan sonar in active and passive modes to locate targets and generate high resolution sonograms. MTS '92: GLOBAL OCEAN PARTNERSHIP. PROCEEDINGS., MTS, WASHINGTON, DC.
- Carter, L., & K. Lewis (1995). Variability of the modern sand cover on a tide and storm driven inner shelf, South Wellington, New Zealand. New Zealand Journal of Geology and Geophysics [N.Z. J. GEOL. GEOPHYS.], **38**(4): 451-470.
- Cervenka, P., & C. de Moustier (1994). Postprocessing and corrections of bathymetry derived from sidescan sonar systems: Application with SeaMARC II. IEEE Journal of Oceanic Engineering **19**(4): 619-629.
- Cervenka, P., U. C. Herzfeld, & C. De Moustier (1994). Accuracy of the spatial representation of the seafloor with bathymetric sidescan sonars. Marine Geophysical Researches **16**(6): 407-425.
- Charlot, D., R. Girault, & B. Zerr (1994). Delph-sonar: a compact system for the acquisition and processing of side scan sonar images. OCEANS, **94(HELD IN CONJUNCTION WITH OSATES 94): RESERVATION-PROCEEDINGS**.

- Chavez, P. S., Jr., J. A. Anderson, & J. W. Schoonmaker, Jr. (1987). Underwater mapping using GLORIA and MIPS. PROCEEDINGS. OCEANS '87. THE OCEAN AN INTERNATIONAL WORKPLACE, **3**: 1202-1205.
- Chin, J. L., H. A. Karl, & N. M. Maher (1997). Shallow subsurface geology of the continental shelf, Gulf of the Farallones, California, and its relationship to surficial seafloor characteristics. Marine Geology **137**(3-4): 251-269.
- Chin-Yee, M. B., D. L. McKeown, & G. D. Steeves (1997). Proven equipment for selectively sampling the seafloor. PROCEEDINGS OF OCEANS '97, **1**(766).
- Churnside, J. H., & S. G. Hanson (1994). Effect of penetration depth and swell-generated on delta-k lidar performance. Appl. Optics, **33**(12): 2363-2367.
- Clarke, J. E. H., L. A. Mayer, & D. E. Wells (1996). Shallow-water imaging multibeam sonars: A new tool for investigating seafloor processes in the coastal zone and on the continental shelf. Marine Geophysical Researches **18**(6): 607-629.
- Clarke, J. H. (1994). Toward remote seafloor classification using the angular response of acoustic backscattering: A case study from multiple overlapping GLORIA data. IEEE Journal of Oceanic Engineering **19**(1): 112-127.
- Cobra, D. T., & H. Moraes (1994). Classification of side-scan sonar images through parametric modeling. OCEANS, **94**(HELD IN CONJUNCTION WITH OSATES 94): RESERVATION-PROCEEDINGS.
- Cobra, D. T., *et al.* (1992). Geometric distortions in side-scan sonar images: a procedure for their estimation and correction. IEEE Journal of Oceanic Engineering, **17**(3): 252-268.
- Cohen, L. M. (1994). Bathymetric data held at the National Geophysical Data Center. Marine Georesources & Geotechnology [MAR. GEORESOUR. GEOTECHNOL.], **12**(1): 53-60.
- Coles, B. (1997). Laser line scan systems as environmental survey tools. Ocean News & Technology [OCEAN NEWS TECHNOL.], **3**(4): 22-27.
- Coles, R. W. (1993). Asset researchers advanced side scan sonar. International Underwater Systems Design [INT. UNDERWAT. SYST. DES.], **15**(3): 5-12.
- Collins, W., R. Gregory, & J. Anderson (1996). A digital approach to seabed classification. Sea Technology **37**(8): 83-87.
- Cote, D., *et al.* (1998). A coded acoustic telemetry system for high precision monitoring of fish location and movement: Application to the study of nearshore nursery habitat of juvenile Atlantic cod (*Gadus morhua*). Marine Technology Society Journal [Mar. Technol. Soc. J.], **32**(1): 54-62.
- Coulter, M. C., *et al.* (1987). Mapping of wood stork foraging habitat with satellite data. Colonial Waterbirds [COLONIAL WATERBIRDS.], **10**(2): 178-180.

- Cowardian, L.M., Carter, V., Golet, F.C., and LaRoe, E.T., 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish and Wildlife Service, FWS/OBS-79/31.
- Czarnecki, M. F. An application of pattern recognition techniques to side scan sonar data. In: Oceans '79.
- Czarnecki, M., & J. M. Bergin (1985). Characteristics of ocean bottom roughness for several seamounts derived from multibeam bathymetric data. OCEANS '85 PROCEEDINGS: OCEAN ENGINEERING AND THE ENVIRONMENT, 1: 663-672.
- Daniel, S., et al. (1998). Side-Scan Sonar Image Matching. IEEE Journal of Oceanic Engineering 23(3): 245-259.
- De Groot, S. J. (1984). The impact of bottom trawling on benthic fauna of the North Sea. Ocean Management, vol. 9: 177-199.
- De Moustier, C. (1988). State of the art in swath bathymetry systems. Int. Hyd. Rev., vol 65, no. 2.
- de Moustier, C., & H. Matsumoto (1993). Seafloor acoustic remote sensing with multibeam echo-sounders and bathymetric sidescan sonar systems. Marine Geophysical Researches 15(1): 27-42.
- De Moustier, C., & J. F. Gettrust (1994). Advances in sea floor management. MTS, 94: 141.
- Dekker, A.G., Malthus, T.J.M., Hoogenboom H.J., 1994, Quantitative determination of chlorophyll a, cyanophycocyanin, seston dry weight, secchi depth transparency and the vertical attenuation coefficient in eutrophic surface waters by airborne imaging spectrometry in the Proceedings of the First International Airborne Remote Sensing Conference and Exhibition, Strasbourg, France, Vol. I pp. 141.
- Denbigh, P. N. (1981). Stereoscopic Visualization and Contour Mapping of the Sea Bed Using Bass. ELECTRONICS FOR OCEAN TECHNOLOGY, INST. OF ELECTRONIC AND RADIO ENG., LONDON, 51.
- Denbigh, P. N. (1989). Swath bathymetry: Principles of operation and an analysis of errors. IEEE Journal of Oceanic Engineering 14(4): 289-298.
- Denbigh, P. N. (1994). Signal processing strategies for a bathymetric sidescan sonar. IEEE Journal of Oceanic Engineering 19(3): 382-390.
- Dethier, M.N., 1992. Classifying marine and estuarine natural communities: an alternative to the Cowardian system. Natural Areas Journ., v. 12, p. 90-100.
- Diether, M. N. (1990). A marine and estuarine habitat classification system for Washington State. Washington Natural Heritage Program, Dept. of Natural Resources, Olympia, WA: 56 p.
- Douglas, B. L., H. Lee (1993). Motion compensation for improved sidescan sonar imaging. Oceans'93, vol. I: 378-383.

- Dziak, R. P., H. Matsumoto, & C. G. Fox (1993). Estimation of seafloor roughness spectral parameters from multi-beam sonar acoustic backscatter data: Axial seamount, Juan de Fuca Ridge. Geophysical Research Letters, **20**(17): 1863-1866.
- Edgerton, H. E. (1986). Sonar images. PRENTICE HALL, ENGLEWOOD CLIFFS, NJ, **304**.
- Edsall, T. A., *et al.* (1989). Side-scan sonar mapping of lake trout spawning habitat in northern Lake Michigan. North American Journal of Fisheries Management [N. AM. J. FISH. MANAGE.], **9**(3): 269-279.
- Eittreim, S. L., R. W. Embley, & W. R. Normark (1983). Monterey sea-valley meander revisited. EOS, Transactions, American Geophysical Union [EOS TRANS. AM. GEOPHYS. UNION.], **64**(45): 730.
- Elachi, C., *et al.* (1984). Spaceborne radar subsurface imaging in hyperarid regions. IEEE Trans. Geoscience and Remote Sensing, vol. GE-22: 383-388.
- Estep, L., J. Lillycrop, & L. Parson (1994). Estimation of maximum depth of penetration of a bathymetric lidar system using a Secchi depth database. REMOTE SENSING FOR MARINE AND COASTAL ENVIRONMENTS, **28**: 2.
- Fader, G. B. J. (1985). Surficial and bedrock geology of the Grand Banks. BIO REV., pp.
- Fader, G. B. J., *et al.* (1993). A comparison of swath bathymetric data and sidescan sonograms from the inner Scotian shelf off Halifax. Atlantic Geology, **29**(1): 79.
- Farmer, D. (1988). New directions in acoustic research. HYDROACOUSTICS WORKSHOP PROCEEDINGS: 1641.
- Farre, J. A. (1986). The importance of mass wasting processes on the continental slope. Dissertation Abstracts International Part B: Science and Engineering, **46**(11).
- Fish, J. P., H. A. Carr (1990). Sound underwater images. Orleans, MA: Lower Cape Publishing: 189 pp.
- Flemming, B. W. (1976). Side-scan sonar: a practical guide. Int. Hydrogr. Rev., **53**(1): 65-92.
- Forbes, D. L., & R. Boyd (1987). Gravel ripples on the inner Scotian Shelf. Journal of Sedimentary Petrology **57**(1): 46-54.
- Fox, C. G. (1996). Objective classification of oceanic ridge-crest terrains using two-dimensional spectral models of bathymetry: Application to the Juan de Fuca Ridge. Marine Geophysical Researches **18**(6): 707-728.
- Gann, J., *et al.* (1995). Characterization of rockfish (Scorpaenidae) habitats in two areas of complex topography in Southeast Alaska using side-scan sonar. TWENTY THIRD BENTHIC ECOLOGY MEETING.
- Gauldie, R. W., S. K. Sharma, & C. E. Helsley (1996). LIDAR applications to fisheries monitoring problems. Canadian Journal of Fisheries and Aquatic Sciences/Journal

- Canadien des Sciences Halieutiques et Aquatiques. Ottawa [CAN. J. FISH. AQUAT. SCI./J. CAN. SCI. HALIEUT. AQUAT.], 53(06): 1459-1468.
- Gerlotto, F., *et al.* (1994). Exhaustive observation of 3D school structure using multibeam side scan sonar: Potential use for school classification, biomass estimation and behaviour studies. ICES COUNCIL MEETING PAPERS, ICES, COPENHAGEN, 12.
- Goff, J. A., & T. H. Jordan (1989). Stochastic modeling of seafloor morphology: Resolution of topographic parameters by sea beam data. IEEE Journal of Oceanic Engineering 14(4): 326-337.
- Gonzalez, R. C., P. Wintz (1977). Digital Image Processing. Addison-Wesley, pp.:477.
- Gordon, A. (1993). Underwater laser line scan technology. Underwat. Syst. Des., 15(2): 26-27.
- Gordon, A. C. (1994). Turbid water test results of the SM2000 laser line scan system and low light level underwater camera tests. UNDERWATER INTERVENTION '94: MAN AND MACHINE UNDERWATER. CONFERENCE PROCEEDINGS., MARINE TECHNOLOGY SOCIETY, WASHINGTON, DC: 305-311.
- Gordon, D. C., Jr. (1997). Studies in eastern Canada. EFFECTS OF FISHING GEAR ON THE SEA FLOOR OF NEW ENGLAND, 8: 8.
- Gower, J.F.R., Borstad, G.A., 1993, Use of imaging spectroscopy to map solar-stimulated chlorophyll fluorescence, red tides and submerged vegetation, in the Proceedings of the 16th Canadian Symposium on Remote Sensing, Sherbrooke, Quebec, pp. 95-98.
- Gower, J.F.R., Borstad, G.A., Anger, C.D., Edel, H.R., 1992, CCD-based imaging spectroscopy for remote sensing: The FLI and casi Programs, in the Proceedings of the Canadian Journal of Remote Sensing, Toronto, Ontario, Vol. 18. No. 4 pp. 199-208.
- Green, Huntington and Rollings (1990). The Theory and Application of Digital Image Processing. Notes of a workshop presented at the 5th Australasian Remote Sensing Conference in Perth, October 1990.
- Green, M. O. (1986). Side-scan sonar mosaic of a sand ridge field: Southern Mid-Atlantic Bight. Geo-Marine Letters 6(1): 35-40.
- Greene, H.G., Stakes, D.S., Orange, D.L., Barry, J.P., and Robison, B.H., 1993. Application of an ROV in geologic mapping of Monterey Bay, California, USA: Amer. Acad. Underwater Sci., 13th Annual Scientific Diving Symposium, Pacific Grove, California, p. 17-32.
- Greene, H.G., Yoklavich, M.M., Barry, J.P., Orange, D.L., Sullivan, D.E., and Cailliet, G.M., 1994. Geology and related benthic habitats of Monterey Canyon, central California [abs.]: in EOS, Transactions, Amer. Geophys. Union Supplement v. 75, n. 3, San Diego, California, p. 203.
- Greene, H.G., Yoklavich, M.M., O'Connell, T. and Wakefield, W., Starr, R., D. Sullivan, (in press). Classification scheme for marine benthic habitats. Oceanologica Acta.

- Greene, H.G., Yoklavich, M.M., Sullivan, D., and Cailliet, G.M., 1995. A geophysical approach to classifying marine benthic habitats: Monterey Bay as a model: *in* O'Connell, T. and Wakefield, W., eds. Applications of side-scan sonar and laser-line systems in Fisheries Research: Alaska Dept. Fish and Game Special Publication No. 9, p.15-30.
- Greenstreet, S. P. R., *et al.* (1997). An assessment of the acoustic survey technique, RoxAnn, as a means of mapping seabed habitat. ICES journal of marine science. London [ICES J. MAR. SCI.], **54**(5): 939-959.
- Hagen, R. A., H. Vergara, & D. F. Naar (1995). Morphology of San Antonio submarine canyon on the central Chile forearc. Marine Geology **129**(3-4): 197-205.
- Hammerstad, E. (1997). Multibeam echo sounder for EEZ mapping. PROCEEDINGS OF OCEANS '97, **2**(787).
- Hampton, M. A., M. E. Torresan, & J. H. Barber, Jr. (1997). Sea-floor geology of a part of Mamala Bay, Hawai'i. Pacific Science **51**(1): 54-75.
- Haney, J. C. (1989). Remote characterization of marine bird habitats with satellite imagery. Colonial Waterbirds [COLONIAL WATERBIRDS.], **12**(1): 67-77.
- Haralick, R. M., *et al.* (1973). Textural features for image classification. IEEE Trans. Systems, Man, and Cybernetics, SMC-3: 610-621.
- Haralick, R. M., L. G. Shapiro (1985) Image Segmentation Techniques. Computer Vision, Graphics and Image Processing, vol. 29: 100-132.
- Hare, R. (1994). Calibrating Larsen-500 Lidar bathymetry in Dolphin and Union Strait using dense acoustic ground-truth. International hydrographic review. Monaco, **71**(1): 91-108.
- Harma, H. P., *et al.* (1990). A data structure for spatio-temporal databases. International Hydrographic Review, Monaco. LXVII(1).
- Harris, P. T., & M. B. Collins (1984). Side-scan sonar investigation into temporal variation in sand wave morphology: Helwick sands, Bristol Channel. Geo-Marine Letters **4**(2): 91-97.
- Head, M. E. M., *et al.* (1997). Applications of 3-D visualizations of oceanographic data bases. PROCEEDINGS OF OCEANS '97, **2**(787).
- Held, A.A., C. Field and N. Williams, Mapping Mangroves with AIRSAR, Proceedings 7th JPL Airborne Earth Science Workshop (in press), NASA's Jet Propulsion Laboratory in Pasadena, California, USA, January, 1998.
- Hellemn, J. E., T. J. Fredette, & D. A. Carey (1994). Underwater environmental survey operations using laser line scan technology. UNDERWATER INTERVENTION '94: MAN AND MACHINE UNDERWATER. CONFERENCE PROCEEDINGS., MARINE TECHNOLOGY SOCIETY, WASHINGTON, DC: 319-327.

- Hennings, I., *et al.* (1993). Sea surface signatures related to subaqueous dunes detected by acoustic and radar sensors. Continental Shelf Research, vol. 13, no. 8/9: 1023-1043.
- Herlihy, D. R., S. P. Matula, & C. Andreasen (1988). Swath mapping data management within the National Oceanic and Atmospheric Administration. International hydrographic review. Monaco, **65**(2): 55-74.
- Herzfeld, U.C. (1997). Geostatistical methods for evaluation of Seabeam bathymetric surveys: Case studies of Wegener Canyon, Antarctica, Marine Geology, 88: 83-95.
- Hobbs, C. H., III (1986). Side-scan sonar as a tool for mapping spatial variations in sediment type. Geo-Marine Letters **5**(4): 241-245.
- Holvland, M., A. G. Judd (1988). Seabed pockmarks and seepages: impact on geology, biology and the marine environment. London: Graham & Trotman: 293 pp.
- Howland, J. C., H. Singh, M. Marra and D. Potter (1999). Digital mosaicking of underwater imagery. Sea Technology, 40:65-69.
- Hudson, S.J., Moore, G.F., Bale, A.J., Dyer, K.R., Aiken, J.. 1994, An operational approach to determining suspended sediment distributions in the humber estuary by airborne multi-spectral imagery, in Proceedings of The First International Airborne Remote Sensing Conference and Exhibition, Strasbourg, France, Vol. III pp. 10.
- Huff, L. C., H. Orlinsky, & S. Matula (1994). Using near-shore multibeam systems for survey investigations. Hydrographic journal. Plymouth **74**: 9-15.
- Hughes Clarke, J. E., *et al.* (1983). Processing and interpretation of 95-kHz backscatter data from shallow-water multibeam sonars. IEEE Oceans'93 Proc., vol. II,: 437-442.
- Huhnerbach, V. (1996). Hierro debris avalanche and the upper part of the Canary Slide: Sidescan sonar imaging and its geological interpretation. Unpublished report, University of Kiel: 69 pp.
- Irish, J. L., & W. J. Lillycrop (1997). Monitoring New Pass, Florida, with high density lidar bathymetry. Journal of Coastal Research **13**(4): 1130-1140.
- Irish, J. L., L. E. Parson, & W. J. Lillycrop (1995). Detailed bathymetry of four Florida inlets. SAND WARS, SAND SHORTAGES & SAND HOLDING STRUCTURES. PROCEEDINGS OF THE: 243-258.
- Jaffe, J. S., G. Chandran, & E. Reuss (1995). High frequency acoustic imaging in the ocean. PROC ICASSP IEEE INT CONF ACOUST SPEECH SIGNAL PROCESS, IEEE, PISCATAWAY, NJ, **5**: 2793-2796.
- Jameson, S. C. (1981). Biological zonation. KEY LARGO CORAL REEF NATIONAL MARINE SANCTUARY DEEP WATER RESURVEY., NOAA/OCZM, WASHINGTON, DC.

- Jenkins, C. (1997). Building offshore soils databases. Software converts geologic materials into numerics, producing detailed seabed materials maps of Australian EEZ. Sea Technology **38**(12): 25-28.
- Johnson, A. E., & M. Hebert (1996). Seafloor map generation for autonomous underwater vehicle navigation. UNDERWATER ROBOTS., KLUWER ACADEMIC PUBLISHERS, BOSTON, MA: 71-94.
- Johnson, D. (1991). Sidescan sonar imagery analysis techniques. Proc. IEEE Oceans'91, vol. 2: 935-941.
- Johnson, H. P., M. Helferty (1990). The geological interpretation of side-scan sonar. Rev. of Geophysics, vol. 28, no. 4: 357-380.
- Jorgen, N., & P. Steenstrup (1998). Beach nourishment in the North Sea. New multibeam system helps plan and evaluate beach nourishment projects. Hydro International [Hydro Int.], **2**(1): 26-29.
- Jupp, D., Kirk, J.T.O., Harris, G.P., 1994, Detection, identification and mapping cyanobacteria - using remote sensing to measure the optical quality of turbid inland waters, Australian Journal Mar. Freshwater Res.,**45**, 801-28.
- Karl, H. A., *et al.* (1994). Acoustic mapping as an environmental management tool: 1. Detection of barrels of low-level radioactive waste, Gulf of the Farallones National Marine Sanctuary, California. Ocean & Coastal Management [OCEAN COAST. MANAGE.], **22**(3): 201-227.
- Keeton, J. A., & R. C. Searle (1996). Analysis of Simrad EM12 multibeam bathymetry and acoustic backscatter data for seafloor mapping, exemplified at the Mid-Atlantic Ridge at 45 degree N. Marine Geophysical Researches **18**(6): 663-688.
- Kenyon, N. H. (1987). Mass-wasting features on the continental slope of Northwest Europe. Marine Geology **74**(1-2): 57-77.
- Kielland, P., & M. Dagbert (1997). Third generation electronic charts: What they will provide users and how to produce them. PROCEEDINGS OF THE SEVENTH, **4**: 305-312.
- Kieninger, K. (1991). Multibeam echosounding for coastal and shallow waters. Sea Technology **32**(6): 67-69.
- Kirlin, R. L., *et al.* (1993). Towfish orientation and position estimation. IEEE Journal of Oceanic Engineering **18**(3): 319-326.
- Kleinrock, M. C. (1991) Overview of sidescan sonar systems and processing. IEEE Proc. : 77-83.
- Kleinrock, M. C., *et al.* (1992). Practical geological comparison of some seafloor survey instruments. Geophys. Res. Lett. vol. 19, no. 13: 1407-1410.

- Knebel, H. J., & R. C. Circe (1995). Seafloor environments within the Boston Harbor-Massachusetts Bay sedimentary system: A regional synthesis. Journal of Coastal Research **11**(1): 230-251.
- Knebel, H. J., et al. (1996). Seafloor environments in Cape Cod Bay, a large coastal embayment. Marine Geology **133**(1-2): 11-33.
- Kraeutner, P. H., & J. S. Bird (1995). Seafloor scatter induced angle of arrival errors in swath bathymetry sidescan sonar. OCEANS '95: CHALLENGES OF OUR CHANGING GLOBAL ENVIRONMENT, MTS/IEEE, NEW YORK, NY, **2**: 975-980.
- Krekova, M. M., et al. (1994). Numerical evaluation of the possibilities of remote laser sensing of fish schools. Applied Optics **33**(24): 5715-5720.
- Kronman, M. (1992). LIDAR: Light detecting and ranging. Laser technology could revolutionize aerial fish finding. Natl. Fisherman, **72**(12): 40-42.
- Krost, P., et al. (1990). Otter trawl tracks in Kiel Bay(Western Baltic) mapped by side-scan sonar. Meeresforschungen, vol. 32: 344-353.
- Kuons, R. R., & R. D. Cardwell (1981). Significant areas for certain species of food fish and shellfish in Puget Sound. TECH. REP. WASH. DEP. FISH., WDF, OLYMPIA, WA, **49**.
- Kuwahara, R. H., & R. H. Poeckert (1989). Accurate seafloor mapping using precise navigation and side-scan sonar. OCEANS '89: THE GLOBAL OCEAN, **4**: 1148-1152.
- Kvitek, R. G., K.E. Conlan, and P. Iampietro. 1998. Black pools of death: Hypoxic, brine-filled ice gouge depressions become lethal traps for benthic organisms in a shallow Arctic embayment. Marine Ecology Progress Series. **162**: 1-10.
- Le Bas, T. P., & D. C. Mason (1997). Automatic registration of TOBI side-scan sonar and multi-beam bathymetry images for improved data fusion. Marine Geophysical Researches **19**(2): 163-176.
- Le Bas, T. P., D. C. Mason, & N. C. Millard (1995). TOBI image processing -- the state of the art. IEEE Journal of Oceanic Engineering **20**(1): 85-93.
- Le Gal, L., & C. Edy (1997). CARAIBES: An integrated software for multibeam echosounder and sidescan sonar data mapping. PROCEEDINGS OF OCEANS '97, **2**(787).
- LeBlanc, L. R., et al. (1992). Marine sediment classification using the chirp sonar. Journal of the Acoustical Society of America [J. ACOUST. SOC. AM.], **91**(1): 107-115.
- Lenink, C. J., & R. D. Jones, Jr. (1977). Characterization of coastal habitat for migratory birds. In: Environmental assessment of the Alaskan continental shelf, **5**: 413-451.
- Levine, E. R., et al. (1995). Oceanographic mapping with Navy's large-diameter UUV. Sea Technology **36**(6): 49-58.

- Li, R. (1992). Correction of pixel locations of sidescan sonar images using bathymetric data acquired separately. Marine Geodesy, 15(2/3): 211-213.
- Liang, Y., B. Lu, & K. How (1989). Acoustic discrimination of seafloor properties. OCEANS '89: THE GLOBAL OCEAN, 5: 1541-1548.
- Lillycrop, W. J., & L. L. Estep (1995). Generational advancements in coastal surveying, mapping. Sea Technology 36(6): 10-16.
- Lillycrop, W. J., J. L. Irish, & L. E. Parson (1997). SHOALS [Scanning Hydrographic Operational Airborne Lidar] system. Sea Technology 38(6): 17-25.
- Linnett, L. M., et al. (1996). Tools for sonar processing and seabed mapping. Sea Technology 37(6): 87-93.
- Lockwood, M. (1991). Seafloor image maps on CD-ROM. Sea Technology 32(9): 53-57.
- Lockwood, M., & G. W. Hill (1989). 1989 Exclusive Economic Zone symposium: Summary and recommendations. Marine Geodesy 13(4): 347-350.
- Loncarevic, B. D., B. M. Scherzinger (1994). Compensation of ship attitude for multibeam sonar surveys. Sea Technology: 10-15.
- Longhurst, A. R. (1981). Analysis of marine ecosystems. Publ. by: Academic Press.
- Loomis, T. J., & D. J. Kall (1989). Sounding data thinning techniques: A knowledge-based approach to modeling. OCEANS '89: THE GLOBAL OCEAN, 5: 1691-1694.
- Lough, R.G., Valentine, P.C., Brown, C.L., and Michaels, W.L., 1992. Maps showing the distribution of juvenile cod in relation to the sedimentary environment of eastern Georges Bank: U.S. Geol. Survey Open-File Report 92-566.
- Lough, R.G., Valentine, P.C., Potter, D.C., Auditore, P.J., Bolz, G.R., Neilson, J.D., and Perry, R.I., 1989. Ecology and distribution of juvenile cod and haddock in relation to sediment type and bottom currents on eastern Georges Bank. Marine Ecology Progress Series, v. 56, p. 1-12.
- Luyendyk, B. P., E. J. Hajic, & D. S. Simonett (1978). Image enhancement of side-scan sonar mapping. Proc. Int. Symp. Remote Sensing Environ, Publ. by : Environmental Research Institute of Michigan, 1: 1025-1026.
- Luyendyk, B. P., E. J. Hajic, & D. S. Simonett (1983). Side-scan sonar mapping and computer-aided interpretation in the Santa Barbara Channel, California. Marine Geophysical Researches 5(4): 365-388.
- MacDonald, I. R., et al. (1997). NR-1: Deep-ocean introduction of new laser line scanner. Sea Technology 38(2): 59-64.
- MacLeod, W., Stanton-Gray, R., Dyk, A., Farrington, G., 1992, Discrimination of substrate type using airborne remotely sensed data, Bay of Quinte, Ontario, Canada, in the Proceedings of the 15th Canadian Symposium on Remote Sensing, Toronto, Ontario, pp. 13-16.

- Magorrian, B. H., M. Service, & W. Clarke (1995). An acoustic bottom classification survey of Strangford Lough, Northern Ireland. Journal of the Marine Biological Association of the United Kingdom. Plymouth [J. MAR. BIOL. ASSOC. U.K.], **75**(4): 987-992.
- Marks, R. L., S. M. Rock, & M. J. Lee (1995). Real-time video mosaicking of the ocean floor. IEEE Journal of Oceanic Engineering **20**(3): 229-241.
- Masson, D. G., B. J. Bett, & K. G. Birch (1997). Atlantic margin environmental survey. Sea Technology **38**(10): 52-59.
- Masson, D. G., B. Van Niel, & P. P. E. Weaver (1997). Flow processes and sediment deformation in the Canary Debris Flow on the NW African Continental Rise. Sedimentary Geology [SEDIMENT. GEOL.], **110**(3-4): 163-179.
- Matthew, K.R., and Richards, L.R., 1991. Rockfish (Scorpaenidae) assemblages of trawlable and untrawlable habitats off Vancouver Island, British Columbia: North Amer. Journ. Fish. Management, v. 11, p. 312-318.
- Matthews, A.M., Boxall, S.R., 1994, Novel algorithms for the determination of phytoplankton concentration and maturity, in the Proceedings of the Second Thematic Conference on Remote Sensing for Marine and Coastal Environments, New Orleans, Louisiana, Vol I pp. 173-180.
- McGregor, B. A., & G. W. Hill (1989). Seafloor image maps of the US Exclusive Economic Zone. Hydrographic journal. Plymouth [HYDROGR. J.], no, **53**: 9-13.
- McKeown, D. L., & D. E. Heffler (1997). Precision navigation for benthic surveying and sampling. PROCEEDINGS OF OCEANS '97, **1**(766).
- Meinesz, A., *et al.* (1991). First evaluation of the suitability of Spot satellite imagery for the cartography of shallow epibenthic communities in the Western Mediterranean. Oceanologica acta. Paris [OCEANOL. ACTA.], **14**(3): 299-307.
- Michalopoulou, Z. H., D. Alexandrou, & C. De Moustier (1995). Application of neural and statistical classifiers to the problem of seafloor characterization. IEEE Journal of Oceanic Engineering **20**(3): 190-197.
- Miller, J. E., *et al.* (1996). Shallow water multibeam hydrography to IHO standards. Sea Technology **37**(6): 81-86.
- Miller, J. E., J. S. Ferguson, & J. S. Byrne (1995). Shallow water multibeam sonar performance, results from a modern hydrographic survey system. OCEANS '95: CHALLENGES OF OUR CHANGING GLOBAL ENVIRONMENT., MTS/IEEE, NEW YORK, NY, **2**: 985-990.
- Miller, J., R. Tyce, & R. Edwards (1987). Sea Beam mapping systems. PROCEEDINGS, INTERNATIONAL SYMPOSIUM ON MARINE POSITIONING, "POSITIONING THE FUTURE", INSMAP, **86**: 7-16.

- Miller, S. P. (1990). 3-D bathymetric imaging: State of the art. OCEANS '90: ENGINEERING IN THE OCEAN ENVIRONMENT. CONFERENCE PROCEEDINGS.
- Miller, S. P. (1991). 3-D bathymetric imaging: State of the art visualization. Sea Technology **32**(6): 27-32.
- Mitchell, N. C., M. L. Somers (1989) Quantitative backscatter measurements with a long-range sidescan sonar. IEEE J. Oceanic Engineering, vol. 14, no. 4:368-374.
- Moose, P. H. (1980). Acoustical techniques in fisheries remanagement. Publ. Wash. Sea Grant., Publ. by: Washington Sea Grant Program, **1980**: 7.
- Morang, A., & R. L. McMaster (1980). Nearshore bedform patterns along Rhode Island from side-scan sonar surveys. J. Sediment. Petrol., **50**(3): 831-840.
- Mullen, L. J., P. R. Herczfeld, & V. M. Contarino (1996). Progress in hybrid lidar-radar for ocean exploration. Sea Technology **37**(3): 45-52.
- Nelson, C. H., K. R. Johnson, & J. H. Barber, Jr. (1987). Gray whale and walrus feeding excavation on the Bering Shelf, Alaska. Journal of Sedimentary Petrology, **57**(3): 419-430.
- Newton, R. S., A. Stefanon (1975). Application of sidescan sonar in marine biology. Marine Biology, vol. 31: 287-291.
- Niblack, W. (1986). An introduction to digital image processing. Englewood Cliffs, NJ: Prentice Hall: 215 pp.
- Nichols, B., et al. (1993). Correlation of swath bathymetric images with metrics derived from seismic and sidescan data. Atlantic Geology **29**(1): 93.
- O'Connell, V.M. and Wakefield, W.W. (eds.), 1995 Application of side-scan sonar and laser-line systems in Fisheries research: Alaska Dept. Fish and Game Special Publication, 50 p.
- O'Connell, V.M., Carlile, D.C., Wakefield, W.W. and Greene, H.G., 1997. Habitat-based assessments of a long-lived species in the Gulf of Alaska: implications for commercial fisheries management (abs.): Marine Benthic Habitats Conference, Program and Abstracts, Nouméa, New Caledonia, (IFREMER/ORSTOM), p. 86.
- O'Connell, V.M., Wakefield, W.W., and Greene, H.G., 1998. The use of a No-Take Marine Reserve in the eastern Gulf of Alaska to protect essential fish habitats: In Yoklavich, M.M. (ed.), Marine Harvest Refugia for West Coast Rockfish: A Workshop. Nat. Oceanogr. Atmospheric Admin. Tech. Memo. NMFS (NOAA-TM-NMFS-SWFSC-245, p. 127-134.
- O'Connell, V.M. and Carlile, D.W., 1993. Habitat-specific density of adult yelloweye rockfish *Sebastodes ruberrimus* in the eastern Gulf of Alaska: Fishery Bull., U.S. v. 91, p. 304-309.

- Okyar, M., & V. Ediger (1997). Sea-floor sediments and bedforms around Turkey, revealed by side-scan sonar imagery. *Oceanologica acta. Paris [Oceanol. Acta]*, **20**(5): 673-685.
- Olbert, C., Singh, R.P., Schaale, M., Furrer, R., Monitoring of water with an airborne spectrographic imager, in Proceedings of The First International Airborne Remote Sensing Conference and Exhibition , Strasbourg, France, Vol. III pp. 141.
- Ostlund, C., Lindell, T., Sorensen, K., 1994, Experiences from a casi session over Norwegian waters, in Proceedings of The First International Airborne Remote Sensing Conference and Exhibition, Strasbourg, France.
- Pace, N. G., & C. M. Dyer (1979). Machine classification of sedimentary sea bottoms. *IEEE Trans. Geosci. Electron.*, **GE17**, *56*(1979).
- Pace, N. G., & H. Gao (1988). Swathe seabed classification. *IEEE Journal of Oceanic Engineering* **13**(2): 83-90.
- Papatheodorou, G. (1997). Seafloor remote sensing techniques and their applications on fisheries research. [5th Hellenic Symposium on Oceanography and Fisheries. Kavala, Greece, April, 2: 2.
- Parker, G., & E. Saade (1998). Multiple sensor FOCUS-1500. *Sea Technology* **39**(2): 112-116.
- Parrott, D. R., & G. B. J. Fader (1988). Geological interpretation of sidescan sonar, subbottom profiler and SWATH sounding data. *HYDROACOUSTICS WORKSHOP PROCEEDINGS*: 1641.
- Parrott, R., *et al.* (1997). Multibeam bathymetric data for regional assessment of marine geology. *PROCEEDINGS OF OCEANS '97*, **2**(787).
- Parson, L. E., *et al.* (1997). Use of lidar technology for collecting shallow water bathymetry of Florida Bay. *Journal of Coastal Research* **13**(4): 1173-1180.
- Patterson, D. R., & J. Pope (1983). Coastal applications of side scan sonar. *Dock & Harbour Authority*, **64**(755): 132-134.
- Penvenne, L. J., & J. Penvenne (1994). Isis -- versatile sonar data acquisition. *Sea Technology* **35**(6): 59-65.
- Permenter, R. W., W. L. Stubblefield, & D. J. P. Swift (1975). Substrate mapping by sidescan sonar. *Fla. Sci.*, **38**(Suppl.1): 13-14.
- Perugini, N. E., T. R. Waddington, & G. T. Noll (1991). Implementation of the hydrographic data acquisition and processing system (HDAPS) in the NOAA fleet. *International hydrographic review. Monaco*, **68**(1): 127-139.
- Pettersson, L.H., Johannessen, O.M., Frette, O., 1993, Norwegian remote sensing spectrometry for mapping and monitoring of algal blooms and pollution, 89' Proceedings of an ESA JRC workshop ESA-SP 360.

- Phillips, N. W., D. A. Gettleson, & K. D. Spring (1990). Benthic biological studies of the southwest Florida shelf. American Zoologist **30**(1): 65-75.
- Platts, W. S. (1980). A plea for fishery habitat classification. Fisheries, **5**(1): 2-6.
- Poehner, F., & E. Hammerstad (1991). Combining bathymetric mapping, seabed imaging. Sea Technology **32**(6): 17-25.
- Populus, J., Guilloumont, B., Ruiz, O., Tallec, P., 1994, Biomass assessment of green algae proliferations with high resolution airborne images, in Proceedings of The First International Airborne Remote Sensing Conference and Exhibition, Strasbourg, France, Vol. III pp. 153.
- Prager, B. (1995). ISAH-S and QTC View Bottom Classification. Remote Sensing Techniques for Subtidal Classification, workshop, Institute of Ocean Sciences, Sydney, B. C.: 7-8.
- Pratson, L. F., & M. H. Edwards (1996). Introduction to advances in seafloor mapping using sidescan sonar and multibeam bathymetry data. Marine Geophysical Researches **18**(6): 601-605.
- Pratson, L. F., & W. B. F. Ryan (1996). Automated drainage extraction in mapping the Monterey submarine drainage system, California margin. Marine Geophysical Researches **18**(6): 757-777.
- Pratson, L. F., & W. F. Haxby (1996). What is the slope of the U.S. continental slope? Geology, **24**(1): 3-6.
- Prior, D. B., J. M. Coleman, & H. H. Roberts (1981). Mapping With Side-Scan Sonar. Offshore (Houston) **41**(4): 151-161.
- Prior, D. B., J. M. Coleman, & L. E. Garrison (1979). Digitally acquired undistorted side-scan sonar images of submarine landslides, Mississippi River delta. Geology, **7**(9): 423-425.
- Provencher, L., M. Giguere, & P. Gagnon (1997). [Seabed characteristics between the 10 and 50 meter isobaths around the Magdalen Islands measured by hydroacoustics and sediment sampling. Evaluation of acoustic RoxAnn USP system]. Canadian technical report of fisheries and aquatic sciences/Rapport technique canadien des sciences halieutiques et aquatiques. Imprint varies [CAN. TECH. REP. FISH. AQUAT. SCI./RAPP. TECH. CAN. SCI. HALIEUT. AQUAT.], **49**.
- Quinn, R. (1995). Sidescan sonar. Remote Sensing Techniques for Subtidal Classification, workshop, Institute of Ocean Sciences, Sydney, B. C.: 12-13.
- Rees, E. I. S. (1975). Appendix E. Evaluation of side-scan sonar for locating muddy deposits in Liverpool Bay. SEWAGE SLUDGE DISPOSAL IN LIVERPOOL BAY. RESEARCH INTO EFFECTS, **2**: 100-112.
- Reid, D. G. (1995). Relationships between herring school distribution and sea bed substrate derived from RoxAnn [sea bed classification system]. ICES INTERNATIONAL SYMPOSIUM ON FISHERIES AND PLANKTON ACOUSTICS HELD IN ABERDEEN.

- Rimski-Korsakov, N. A., Y. S. Russak, & R. B. Pavlov (1994). Simple digital system for side scan sonar data imaging. OCEANS, **94**(HELD IN CONJUNCTION WITH OSATES 94): RESERVATION-PROCEEDINGS.
- Ripley, H.T., Thruscitt, A.M., Jones, W., 1994, Acquisition and analysis of airborne multispectral data for the detection and mapping of submerged logs on the vaga river, Russia, in Proceedings of The First International Airborne Remote Sensing Conference and Exhibition, Strasbourg, France, Vol. III pp. 1.
- Riyait, V. S., *et al.* (1995). A review of the acid synthetic aperture sonar and other sidescan sonar systems. International hydrographic review. Monaco, **72**(1): 115-123.
- Robinson, C.L.K., Hay, D. E., Booth, J. Truscott, J. 1996. Standard methods for sampling resources and habitats in coastal subtidal regions of British Columbia: Part 2: Review of sampling with preliminary recommendations. Canadian Technical Report of Fisheries and Aquatic Sciences 2119. Fisheries and Oceans Canada. 118 pp.
- Rochon, G. L., *et al.* (1997). Three-dimensional visualization of oceanographic data: A case study of the Gulf of Mexico's coastal interface with Texas and Louisiana. PROCEEDINGS OF OCEANS '97, **2**(787).
- Rollings (1994) Casi and Port Phillip Bay - Victoria Australia. Proceedings of the Workshop on Water Quality, Melbourne, 28th February.
- Rollings (1995) CASI and Water Quality Mapping. Presentation to Australian Map Circle Conference, 6 -9 Feb, Armidale NSW.
- Rollings, N., Doblin, M. and Chifings, A. (in peer review) A Comparison of CASI, SPOT and Aerial Photography for Benthic Mapping in Port Phillip Bay.
- Rollings, N.M., Light, B.R., Doblin, M.A. and Chiffings, T. (1993). An evaluation of remote sensing and associated field techniques for mapping the distribution of benthic habitats in Port Phillip Bay. Port Phillip Bay Environmental Study Task G2.1 Final report, November 1993.
- Roush, B., G. Adams, & R. New (1980). Bibliographic overview of Ocean Acoustic Remote Sensing (OARS). Publ. Wash. Sea Grant, Publ. by: Washington Sea Grant Program, **1980**: 17-1.
- Rubin, D. M., D. S. McCulloch, & H. R. Hill (1983). Sea-floor-mounted rotating side-scan sonar for making time-lapse sonographs. Continental shelf research. Oxford, New York NY, **1**(3): 295-301.
- Rukavina, N. A. (1995). Mapping contaminated sediments with an acoustic bottom-discrimination system. PROCEEDINGS OF THE 38TH CONFERENCE OF THE INTERNATIONAL ASSOCIATION OF GREAT LAKES RESEARCH, INTERNATIONAL ASSOCIATION FOR GREAT LAKES RESEARCH, **2099**(USA): 63.

- Ryan, W. B., & A. Malinverno (1990). Acquisition of stereo photography for seafloor characterization. Source, **3**.
- Saade, E., G. Parker, & R. Morton (1997). Multi-sensor instrumentation on a FOCUS-1500 search and survey system. PROCEEDINGS OF OCEANS '97, **2**(787).
- Saunders, J.F., Jupp, D.L.B., Harris, G.P., Hawkins, P.R., Byrne, G., Hutton, P.G., 1994, Mapping optical water quality of the Hawkesbury River using casi airborne spectrometer data, 7th Australasian Remote Sensing Conference, Melbourne, Australia.
- Schenke, H. W., & G. Kuhn (1994). Small-scale structures from multibeam bathymetry in the Weddell Sea. ANNU. REP. AWI POLAR MAR. RES., no.
- Schlagintweit, G. (1995). Discussion of (RoxAnn) operations. Remote Sensing Techniques for Subtidal Classification, workshop, Institute of Ocean Sciences, Sydney, B. C.: 5-7.
- Schock, S. G., L. R. Leblanc, & S. Panda (1994). Spatial and temporal pulse design considerations for a marine sediment classification sonar. IEEE Journal of Oceanic Engineering **19**(3): 406-415.
- Schoen, C., & J. Sibert (1996). Feasibility of dual mode lidar for pelagic fish surveys. JIMAR Contrib.
- Schulkin, M. (1980). NOAA workshop on Ocean Acoustic Remote Sensing, Seattle, Washington - January 21-24, 1980. Volume 1, summary report. Publ. by: WSG, **45**.
- Schulkin, M. (1980). NOAA Workshop on ocean acoustic remote sensing. Volume 2: presentations and working group reports. Publ. by: WSG, **423**: ubl.
- Schulkin, M. (1980). Ocean acoustic remote sensing: workshop overview and remarks. Publ. Wash. Sea Grant, Publ. by: Washington Sea Grant Program, **1980**: 16-1.
- Seibold, E., W. H. Berger (1993). The sea floor: an introduction to marine geology. 2nd edn., Springer Verlag: pp. 356.
- Service, M., & B. H. Magorrian (1997). The extent and temporal variation of disturbance to epibenthic communities in Strangford Lough, Northern Ireland. Journal of the Marine Biological Association of the United Kingdom. Plymouth [J. Mar. Biol. Assoc. U.K.], **77**(4): 1151-1164.
- Shapiro, J., & M. Boule Coastal aquatic area management in the Puget Sound region. .
- Shapiro, L. H., & P. W. Barnes (1991). Correlation of nearshore ice movement with seabed ice gouges near Barrow, Alaska. Journal of Geophysical Research. C. Oceans, **96**(C9): 16,979-989.
- Shaw, J., D. R. Parrott, & J. Hughes-Clarke (1996). Use of multibeam bathymetry to determine seabed impacts at the Argentia Naval Base, Newfoundland. PARTNERSHIP IN COASTAL ZONE MANAGEMENT., SAMARA PUBLISHING LTD., CARDIGAN: 499-506.
- Shepard, F.P., 1963. Submarine Geology (3rd ed), New York, Harper and Row.

- Sheppard, C. R. C., *et al.* (1995). Habitat mapping in the Caribbean for management and conservation: Use and assessment of aerial photography. Aquatic Conservation: Marine and Freshwater Ecosystems [AQUAT. CONSERV.: MAR. FRESHWAT. ECOSYST.], **5**(4): 277-298.
- Shippey, G., A. Bolinder, & R. Finndin (1994). Shade correction of side-scan sonar imagery by histogram transformation. OCEANS, 94(HELD IN CONJUNCTION WITH OSATES 94): RESERVATION-PROCEEDINGS.
- Simons, J. D., *et al.* (1992). An improved method for mapping oyster bottom using a global positioning system and an acoustic profiler. AQUACULTURE '92: GROWING TOWARD THE 21st CENTURY, **207**: 207.
- Smith, D. K., & P. R. Shaw (1989). Using topographic slope distributions to infer seafloor patterns. IEEE Journal of Oceanic Engineering **14**(4): 338-347.
- Somers, M. L. (1993) Sonar imaging of the seabed: Techniques, performance, applications. Acoustic Signal Processing for Ocean Exploration, Canadian Govt. : 355-369.
- Somers, M. L. (1995) Resolving the issue: a look at resolution and related topics in sonar. Man-made objects on the seafloor, Society for Underwater Technology, U. K.: 41-58.
- Somers, M. L., A. R. Stubbs (1984). Sidescan sonar. IEE Proceedings, vol. 131, Part F, no. 3: 243-256.
- Sotheran, I. S., R. L. Foster-Smith, & J. Davies (1995). Mapping of marine benthic habitats using image processing techniques within a raster-based geographic information system. SCIENCE FOR MANAGEMENT IN COASTAL AND ESTUARINE WATERS: PROCEEDINGS OF THE 25TH ANNUAL SYMPOSIUM OF THE ECSA. PART, 44(SUPPL).
- Spencer, P. (1997). Bathymetric measurement -- a data quality standard. Hydrographic journal. Plymouth [Hydrogr. J.], no, **85**: 9-15.
- Spiess, F. N. (1980). Ocean acoustic remote sensing of the sea floor. Publ. Wash. Sea Grant., Publ. by: Washington Sea Grant Program, **1980**: 11.
- St.C.Wright, A. (1997). Deep-towed sidescan sonars. Sea Technology **38**(6): 31-38.
- Staenz, K., 1992, A decade of imaging spectrometry in Canada, in the Proceedings of the Canadian Journal of Remote Sensing, Toronto, Ontario, Vol. 18 No. 4 - PP. 187-197.
- Staenz, K., 1992, Development of imaging spectrometry in Canada, in the Proceedings of the International Symposium on Spectral Sensing Research (ISSSR), Maui, Hawaii.
- Stanic, S., *et al.* (1988). Shallow-water high-frequency bottom scattering off Panama City, Florida. Journal of the Acoustical Society of America [J. ACOUST. SOC. AM.], **83**(6): 2134-2144.

- Stea, R. R., *et al.* (1994). Morphology and seismic stratigraphy of the inner continental shelf off Nova Scotia, Canada: Evidence for a -65 m lowstand between 11,650 and 11,250 C super(14) yr B.P. Marine Geology **117**(1-4): 135-154.
- Stein, D.L., Tissot, M.A., Hixon, M.A. and Barss, W., 1992. Fish-habitat associations on a deep reef at the edge of the Oregon continental shelf. Fishery Bull., U.S. 90; p. 540-551.
- Steinvall, O., & J. Banic (1997). Airborne laser hydrography: A vital part of future coastal mapping. Hydro International [HYDRO INT.], **1**(3): 25-27.
- Stewart, W. K. (1990). A model-based approach to 3-D imaging and mapping underwater. Transactions of the ASME. Journal of Offshore Mechanics and Arctic [J. OFFSHORE MECH. ARCT. ENG.], **112**(4): 352-356.
- Stewart, W. K. S. (1990). Subsea Data Processing Standards, RIDGE Workshop Report.
- Stewart, W. K., *et al.* (1994). Quantitative seafloor characterization using a bathymetric sidescan sonar. IEEE Journal of Oceanic Engineering **19**(4): 599-610.
- Stone, D.J., Freemantle, J.R., Shepherd, P.R., Miller, J.R., 1993, Total water vapour amount determined from casi imagery, in the Proceedings of the 16th Canadian Symposium on Remote Sensing, Sherbrooke, Quebec, pp. 805-809.Talukdar, K. K., & R. C. Tyce (1990). Digital processing of sidescan images from bottom backscatter data collected by sea beam. Marine Geodesy, **14**(2): 81-100.
- Tamsett, D. (1993). Sea-bed characterisation and classification from the power spectra of side-scan sonar data. Marine Geophysical Researches **15**(1): 43-64.
- Thorpe, S. A., & A. J. Hall (1993). Nearshore side-scan sonar studies. Journal of Atmospheric and Oceanic Technology [J. ATMOS. OCEAN. TECHNOL.], **10**(5): 778-783.
- Thorpe, S. A., *et al.* (1982). Wave-produced bubbles observed by side-scan sonar. Nature, vol. 296: 636-638.
- Tsai, J. J. (1984). Acoustic remote sensing of waste disposal in the ocean. NOAA TECH. MEMO., **108**.
- Tsemahman, A. S., W. T. Collins, & B. T. Prager (1997). Acoustic seabed classification and correlation analysis of sediment properties by QTC view. PROCEEDINGS OF OCEANS '97, **2**(787).
- Tuck, I. D., S. J. Hall, & D. G. Reid (1995). Identification of benthic disturbance by fishing gear using RoxAnn [sea bed classification system]. ICES INTERNATIONAL SYMPOSIUM ON FISHERIES AND PLANKTON ACOUSTICS HELD IN ABERDEEN.
- Twichell, D. C., & D. G. Roberts (1982). Morphology, Distribution, and Development of Submarine Canyons on the United States Atlantic Continental Slope between Hudson and Baltimore Canyons. Geology, **10**(8): 408-412.

- Twichell, D.C. and Able, K.W., 1993. Bathymetry, sidescan sonar image, and surficial geological interpretation of the inner shelf off Little Egg Inlet, New Jersey. U.S. Geol. Survey Misc. Field Studies Map, MF-2221.
- Uchida, K., *et al.* (1995). Evaluation of settled position of component reef blocks in an area with some large-scaled artificial reef groups detected by side scan sonar. Fisheries engineering (Japan)/Suisan Kogaku (Japan). Ibaraki [FISH. ENG. (JAPAN)/SUISAN KOGAKU (JAPAN)], **32**(1): 13-22.
- Udintsev, G. B., Y. Odinokov, & V. M. Golod (1987). Experience of and prospects for application of multibeam echosounding in marine geomorphology. Okeanologiya/Oceanology (Mosc.), **27**(3): 497-506.
- Valentine, P.C., and Lough, R.G., 1991. The sea floor environment and the fishery of eastern Georges Bank—the influence of geologic and oceanographic environmental factors on the abundance and distribution of fisheries resources of the northeastern United States continental shelf. U.S. Geol. Survey Open-File Report 91-439.
- Valentine, P.C., and Schmuck, C.A., 1995. Geological mapping of biological habitats on Georges Bank and Stellwagen Bank, Gulf of Maine region: *in* O'Connell, T. and Wakefield, W., eds. Applications of side-scan sonar and laser-line systems in Fisheries Research: Alaska Dept. Fish and Game Special Publication No. 9, p. 31-40.
- Voulgaris, G., M. B. Collins (1991). Linear features on side-scan sonar images: an algorithm for the correction of angular distortion. Marine Geology, **96**(1/2): 187-190.
- Wakefield, W.W., Greene, H.G., O'Connell, V.M. and Gann, J., 1996. The offshore Edgecumbe lava field, Southeast Alaska: geological and habitat characterization of a commercial fishing ground. Poster, Amer. Geophys. Union, ASLO Ocean Sciences Meeting, San Diego, EOS, Transaction, AGU v. 76, n. 3, p. 95.
- Warner, L. S. (1986). Mapping the unmappable; use of geographic information systems in fisheries management. PROCEEDINGS OF TENTH NATIONAL CONFERENCE. ESTUARINE AND COASTAL MANAGEMENT: TOOLS OF THE TRADE. NEW ORLEANS, LOUISIANA, **2**: 705.
- Warner, L. S. (1987). Delineation of marine habitat: A study of spatial error. Dissertation Abstracts International Part B: Science and Engineering, **47**(11).
- Wiedicke, M. H., & M. E. Weber (1996). Small-scale variability of seafloor features in the northern Peru Basin: Results from acoustic survey methods. Marine Geophysical Researches **18**(5): 507-526.
- Williams, D. G. (1994). Marine habitats of the Cocos (Keeling) Islands. ECOLOGY AND GEOMORPHOLOGY OF THE COCOS(406): 399-414.
- Williams, S. J. (1982). Use of high resolution seismic reflection and side-scan sonar equipment for offshore surveys. COASTAL ENG. TECH. AID, U.S. ARMY COAST. ENG. RES. CENT., CERC, FORT BELVOIR, VA, **25**.

- Williamson, M. E. (1993). Evaluation of seafloor properties using co-registered swath bathymetry, sidescan sonar imagery, and high resolution seismic profiles. MTS '93: TECHNOLOGY REQUIREMENTS IN THE NINETIES. PROCEEDINGS., MTS, WASHINGTON, DC.
- Wilson, A.K., 1990, The NERC 1989 compact airborne spectrographic imager (casi) campaign", in the Proceedings of the Natural Environment Research Council Symposium on Airborne Remote Sensing, British Geological Survey, Keyworth, Nottingham, U.K., pp. 259-283.
- Woodward-Clyde Consultants, L. A. C. A. (1979). Eastern Gulf of Mexico marine habitat study. Volume 1. Publ. by: W CC, 109.
- Woodworth-Linas, C. M. T. (1983). The relative age of ice scours using cross-cutting relationships. C CORE publication. St. John's NF [C CORE PUBL.], no.
- Woodworth-Lynas, C. M. T., & J. V. Barrie (1985). Iceberg scouring frequencies and scour degradation on Canada's eastern shelf areas using sidescan mosaic remapping techniques. C CORE publication. St. John's NF [C CORE PUBL.], no.
- Yoklavich, M.M., 1997. Applications of side-scan sonar and in-situ submersible survey techniques to marine fisheries habitat research: *In* Boehlert, G.W. and J.D. Schumacher, eds. Changing Oceans and Changing Fisheries: Environmental data for fisheries research and management. NOAA Tech. Memo. NOAA-TM-NMFS-SWFSC-239, p. 140-141.
- Yoklavich, M.M., Cailliet, G.M., Greene, H.G., and Sullivan, D., 1995. Interpretation of side-scan sonar records for rockfish habitat analysis: examples from Monterey Bay: *in* O'Connell, T. and Wakefield, W., eds. Applications of side-scan sonar and laser-line systems in Fisheries Research: Alaska Dept. Fish and Game Special Publication No. 9, p. 11-14.
- Yoklavich, M.M., Greene, H.G., Moreno, G., Cailliet, G.M., and Sullivan, D., 1992. The importance of small-scale refugia to deep water rockfishes (*Sebastodes sp.*) - a pilot study in Soquel Canyon, Monterey Bay, California [abs.]: *in* EOS, Transactions, Amer. Geophys. Union, v. 73. no 43, p. 318.
- Yoklavich, M.M., Starr, R. Steger, J., Greene, H.G., Schwing, F. and Malzone, C., 1997. Mapping benthic habitats and ocean currents in the vicinity of central Big Creek Ecological Reserve. NOAA Tech. Memo. NOAA-TM-NMFS-SWFSC-245, 52 p.
- Zacharias, M., Niemann, O., Borstad, G., 1992, An assessment and classification of a multispectral bandset for the remote sensing of intertidal seaweeds, in the Proceedings of the Canadian Journal of Remote Sensing, Toronto, Ontario, Vol 18 No. 4 - pp. 263-274.

9. ACKNOWLEDGEMENTS

This work would not have been possible without the generous contributions from numerous colleagues and partner institutions sharing our interest in and commitment to exploring and mapping nearshore marine habitats. Our thanks to: S. Blasco, G. Greene, M. Yoklavich, D. Ventresca, M. Gingras, J. Paris, B. Woodruff, Geological Survey of Canada, C. Malzone, Triton-Elics International, T. Anderson, R. Anyma, B. Appelgate, B. Apsy, R. Armstrong, J. Borgeld, G. Borstad, D. Bray, M. Buckingham, J. Burns, M. Caballero, A. Cameron, F. Campbell, P. Campell, D. Carlson, C. Carpenter, M. Carr, J. Castro, D. Caulfield, T. Cawood, P.S. Chavez Jr., D. Clague, M. Clancy, J. Lynch, G. Cochrane, E. Coffman, I. Colburn, B.W. Coles, L. Cooper, R. Cosgrove, T. Curan, P. Dartnell, J. Dasler, G. Davis, C. DeMoustier, W. Dietrich, C. Dorman, E. Druffel, M. Dupuis, S. Eittreim, C. Elliot, L. Estip, J.H. Evans, R. Evans, P. Fischer, V. Fisher, C. Flynn, L. Ford, J. Foster, D. Francis, J. Gardner, J. Garlik, B. Garwood, M. Giaramita, J.A. Goff, J. Greiner, G. Griggs, C. Grimes, K. Grove, C. Gutmacher, B. Hall, D. Hallard, C. Hancock, R. Hanson, G. Hatcher, A. 339Held, B. Hoffman, T. Janitess, S. Johnson, M. Kennedy, D. Keyes, J. Kilgore, J. Klasik, A. Kleiner, G. Kozak, W. Krabil, P. Kronfield, K. Kuittinen, F. Kyte, B. Layborn, C. Lee, D. Leedy, N. Lesnikowski, S. Lewis, J. Lilycrop, J. Lima, E. Livingston, K. Long, M. Love, B. Luyendyk, N. Maher, M. Mattox, J. McClain, R. McClain, R. McConnaughey, S. McGill, T. Mckenzie, S. Miller, A. Mitsche, M. Moline, C. Moore, J.C. Moore, M. Morrison, D. Moss, M. Muirhead, H. Murphy, R. Myers, R. Negrini, J. Oakden, G. O'Keefe, D. Peyton, R. Piper, D. Reed, F. Rezac, B. Rosenthal, F. Schwing, A. Shak, P. Sharfstein, N. Shaw, C. Shue, D. Sigurdson, J. Snow, J. Stamm, W. Stroup, T. Sullivan, S. Talbott, J. Tamplin, J. Thayer, C. Thormeyer, J. Van Coutis, B. Van Wagenen, P. Veisze, R. Villa, B. Virden, D. Wagner, B. Waltzenberger, R. Warnken, J. Wilson, F. Wolf, R. Wright.