

# Allosteric activation or inhibition of PI3Ky mediated through conformational changes in the p110 $\gamma$ helical domain

## Reviewed Preprint

Revised by authors after peer review.

[About eLife's process](#)

## Reviewed preprint version 2

June 12, 2023 (this version)

## Reviewed preprint version 1



May 23, 2023

## Posted to bioRxiv

April 12, 2023

## Sent for peer review

April 12, 2023

Noah J Harris, Meredith L Jenkins, Sung-Eun Nam, Manoj K Rathinaswamy, Matthew AH Parson, Harish Ranga-Prasad, Udit Dalwadi, Brandon E Moeller, Eleanor Sheekey, Scott D Hansen, Calvin K Yip , John E Burke 

Department of Biochemistry and Microbiology, University of Victoria • Department of Biochemistry and Molecular Biology, The University of British Columbia, Vancouver • Department of Chemistry and Biochemistry, Institute of Molecular Biology, University of Oregon

 ([https://en.wikipedia.org/wiki/Open\\_access](https://en.wikipedia.org/wiki/Open_access))

 (<https://creativecommons.org/licenses/by/4.0/>)

## Abstract

PI3Ky is a critical immune signaling enzyme activated downstream of diverse cell surface molecules, including Ras, PKC $\beta$  activated by the IgE receptor, and G $\beta\gamma$  subunits released from activated GPCRs. PI3Ky can form two distinct complexes, with the p110 $\gamma$  catalytic subunit binding to either a p101 or p84 regulatory subunit, with these complexes being differentially activated by upstream stimuli. Here using a combination of cryo electron microscopy, HDX-MS, and biochemical assays we have identified novel roles of the helical domain of p110 $\gamma$  in regulating lipid kinase activity of distinct PI3Ky complexes. We defined the molecular basis for how an allosteric inhibitory nanobody potently inhibits kinase activity through rigidifying the helical domain and regulatory motif of the kinase domain. The nanobody did not block either p110 $\gamma$  membrane recruitment or Ras/G $\beta\gamma$  binding, but instead decreased ATP turnover. We also identified that p110 $\gamma$  can be activated by dual PKC $\beta$  helical domain phosphorylation leading to partial unfolding of an N-terminal region of the helical domain. PKC $\beta$  phosphorylation is selective for p110 $\gamma$ -p84 compared to p110 $\gamma$ -p101, driven by differential dynamics of the helical domain of these different complexes. Nanobody binding prevented PKC $\beta$  mediated phosphorylation. Overall, this work shows an unexpected allosteric regulatory role of the helical domain of p110 $\gamma$  that is distinct between p110 $\gamma$ -p84 and p110 $\gamma$ -p101 and reveals how this can be modulated by either phosphorylation or allosteric inhibitory binding partners. This opens possibilities of future allosteric inhibitor development for therapeutic intervention.

### eLife assessment

This study presents **fundamental** new insight into the regulatory apparatus of PI3Kgamma; an **important** kinase in signaling pathways that control the immune response and cancer. A suite of biophysical and biochemical approaches provide **convincing** evidence for new sites of allosteric control over enzyme activity. The rigorous findings provide structure and dynamic information that may be exploited in efforts to control PI3Kgamma activity in a therapeutic setting.

## Introduction

The class I phosphoinositide 3 kinases (PI3Ks) are master regulators of myriad functions through their generation of the lipid signalling molecule phosphatidylinositol 3,4,5-trisphosphate (PIP<sub>3</sub>) downstream of cell surface receptors (Burke and Williams, 2015; Rathinaswamy and Burke, 2019; Vanhaesebroeck et al., 2021; Vasan and Cantley, 2022). The class I PI3Ks can be further subdivided into the class IA and class IB subfamilies, with class IB PI3Ks being critical in immune signalling, and are composed of a single p110γ catalytic subunit that can bind to either a p101 or p84 regulatory subunit (Hawkins and Stephens, 2015; Lanahan et al., 2022; Okkenhaug, 2013). The two PI3Kγ complexes (either p110γ-p84 or p110γ-p101) play essential and independent roles in both the adaptive and innate immune systems. PI3Kγ has shown promise as a therapeutic target, primarily as an immunomodulator of the tumor microenvironment leading to enhanced anti-tumor immune responses (De Henau et al., 2016; Kaneda et al., 2016b). Multiple isoform selective small molecule ATP competitive inhibitors of p110γ are in clinical trials for multiple forms of human cancers (Li et al., 2021). However, all inhibitors currently developed towards p110γ act as ATP competitive inhibitors, showing equal potency against both p110γ-p84 or p110γ-p101 complexes.

Detailed experiments on the role of p110γ in mice show that knockout of both p101 and p84 leads to PIP<sub>3</sub> levels that are equivalent to knockout of p110γ, showing that all cellular PI3Kγ activity requires the presence of either a p84 or p101 regulatory subunit (Rynkiewicz et al., 2020). The two complexes are differentially activated by membrane localised receptors, including G-protein coupled receptors (Li et al., 2000; Stephens et al., 1997), Ras (Jin et al., 2020; Kurig et al., 2009), toll like receptors (TLRs) (Luo et al., 2018), and the IgE antigen receptor (Laffargue et al., 2002; Walser et al., 2013). This leads to the different complexes driving unique immune responses, with p110γ-p101 involved in chemotaxis in neutrophils (Bohnacker et al., 2009; Deladeriere et al., 2015), and p110γ-p84 involved in reactive oxide production. Differential activation of unique PI3Kγ complexes downstream of GPCRs and Ras is caused by the ability of p101 to directly bind to Gβγ subunits downstream of activated GPCRs, with this being lost in p84, making p110γ-p84 activation by Gβγ dependent on Ras mediated membrane recruitment (Rathinaswamy et al., 2023; Kurig et al., 2009; Rynkiewicz et al., 2020). Activation of PI3Kγ downstream of the IgE antigen receptor is driven by calcium mediated activation of protein kinase Cβ, leading to the selective phosphorylation and activation of p110γ at S582 (Walser et al., 2013), with this putatively only occurring in p110γ-p84 and not p110γ-p101. The full molecular mechanisms underlying how phosphorylation of p110γ is selective for different p84 or p101 complexes, and how it activates lipid kinase activity are poorly understood.

Extensive biophysical approaches including cryo electron microscopy (cryo-EM), X-ray crystallography, and hydrogen deuterium exchange mass spectrometry (HDX-MS) have provided extensive insight into the molecular underpinnings of how p110 $\gamma$  associates with both p101 and p84, how they are differentially activated by Ras and GPCR signals, and how they can be activated on lipid membranes (Pacold et al., 2000; Walker et al., 1999; Rathinaswamy et al., 2021c; 2021a; Gangadhara et al., 2019; Vadas et al., 2013; Rathinaswamy et al., 2023). The p110 $\gamma$  catalytic subunit is composed of an adaptor binding domain (ABD), a Ras binding domain (RBD), a C2 domain, a helical domain, and a bi-lobal kinase domain (Rathinaswamy et al., 2021a; Walker et al., 1999). A set of helices positioned C-terminal to the activation loop in the kinase domain play a critical role in regulating activity, with this region referred to as the regulatory motif (Rathinaswamy et al., 2021c). The p110 $\gamma$  isoform is unique in that it is inhibited in the absence of a regulatory subunit, with this driven by an autoinhibitory conformation of the regulatory motif, that is proposed to require membrane association to disrupt (Gangadhara et al., 2019). The regulatory motif is a common site of activating mutations in the other class I PI3K isoforms (Jenkins et al., 2023), with p110 $\gamma$  having rare activating oncogenic mutants in this region (Rathinaswamy et al., 2021c). The p110 $\gamma$  subunit interacts with both p84 and p101 at an interface composed of the C2 domain, and the linkers between the RBD-C2 and C2-helical domains (Rathinaswamy et al., 2023; 2021a). The p110 $\gamma$ -p84 complex forms a more dynamic complex compared to p110 $\gamma$ -p101 (Rathinaswamy et al., 2023; Shymanets et al., 2013), however, no clear unique regulatory role of this difference in dynamics has been identified.

The fundamental roles of p110 $\gamma$  in inflammatory processes has made it a therapeutic target in many pathological conditions, including asthma (Campa et al., 2018), arthritis (Camps et al., 2005), obesity (Becattini et al., 2011; Breasson et al., 2017), and cancer (De Henau et al., 2016; Kaneda et al., 2016a; 2016b). There are significant side effects from non-isoform selective PI3K inhibitors (Bohnacker et al., 2017; Vanhaesebroeck et al., 2021), which has driven the development of highly p110 $\gamma$  selective small molecule inhibitors (Bell et al., 2012; Evans et al., 2016; Gangadhara et al., 2019). However, all p110 $\gamma$  inhibitors will target both p110 $\gamma$ -p101 and p110 $\gamma$ -p84, so there is a potential for the development of allosteric inhibitors outside of the ATP binding site. Initial promise has been reported for the development of class IA p110 $\alpha$  oncogene specific allosteric inhibitors. However, further investigation of the molecular mechanisms underlying p110 $\gamma$  regulation will be required for the discovery of regions that can be targeted for allosteric inhibitor development.

Here we report critical roles of the helical domain of p110 $\gamma$  in both activation and inhibition of lipid kinase activity. We characterised an allosteric inhibitory nanobody (NB7) that potently inhibits p110 $\gamma$  activity. Cryo-EM was used to define the inhibitory interface, which is composed of the helical domain, the ABD-RBD linker, and the regulatory motif of the kinase domain of p110 $\gamma$ . The region that the nanobody binds to is in close spatial proximity to a previously identified PKC $\beta$  phosphorylation site (S582) in the helical domain, and oncogenic activating mutants in the regulatory motif. We fully characterised the activity and dynamics of stoichiometrically PKC $\beta$  phosphorylated p110 $\gamma$ , leading to the discovery of a novel additional phosphorylation site (either S594 or S595). PKC $\beta$  phosphorylation was highly selective for p110 $\gamma$  and p110 $\gamma$ -p84, with limited phosphorylation of p110 $\gamma$ -p101. Hydrogen deuterium exchange mass spectrometry (HDX-MS) analysis showed that phosphorylation of p110 $\gamma$  leads to unfolding of the N-terminal region of the helical domain, and increased kinase activity. The presence of the inhibitory nanobody significantly blocks PKC $\beta$  phosphorylation, while phosphorylation of p110 $\gamma$  prevented binding to NB7. Overall, this work provides unique insight into the critical role of the helical domain in controlling p110 $\gamma$  activity, and how phosphorylation and binding partners can modify this regulation. It also reveals a unique binding site located at the interface of the helical and kinase domain that can be targeted for future allosteric inhibitor design.

## Results

### Molecular mechanism of nanobody inhibition of p110 $\gamma$


We previously identified multiple nanobodies that inhibited the activity of p110 $\gamma$ . One from this group (denoted NB7 throughout the manuscript) potently inhibited the membrane mediated activation of p110 $\gamma$ -p84 by both Ras and G $\beta\gamma$ , with HDX-MS experiments mapping the NB7 binding interface to the RBD, helical and kinase domains (Rathinaswamy et al., 2021b). We originally hypothesized that NB7 worked by sterically inhibiting Ras binding to the RBD domain of p110 $\gamma$ . To further explore the molecular mechanism of inhibition we purified all complexes of p110 $\gamma$  (p110 $\gamma$  apo, p110 $\gamma$ -p84, p110 $\gamma$ -p101) along with the NB7 nanobody. The SDS-PAGE of all proteins utilised in this study are shown in the source data file included in the supplemental information.

To define the mechanism concerning how NB7 inhibits PI3K activity we analyzed how this nanobody inhibited all class IB PI3K complexes (p110 $\gamma$ , p110 $\gamma$ -p84, p110 $\gamma$ -p101) upon activation by lipidated G $\beta\gamma$  subunits. Intriguingly, we found that all three forms of p110 $\gamma$  were potently inhibited by NB7 (Fig. 1A). While the IC<sub>50</sub> measured for the three complexes was different, this is likely mainly due to the dramatic difference in protein required to measure lipid kinase activity *in vitro* (~300 nM for p110 $\gamma$  apo/p110 $\gamma$ -p84, and ~10 nM for p110 $\gamma$ -p101, respectively). This suggested that the mechanism of inhibition was not driven by a steric block of Ras association through the RBD, as previously proposed (Rathinaswamy et al., 2021b). We examined the binding of this nanobody to all complexes using biolayer interferometry (BLI). The nanobody bound equivalently tightly to all complexes, with ~2 nM potency for p110 $\gamma$ , p110 $\gamma$ -p84, and p110 $\gamma$ -p101 (Fig. 1B). We also tested binding of the nanobody to all class IA PI3Ks, and there was no detectable binding to p110 $\alpha$ , p110 $\beta$ , and p110 $\delta$  (Fig. 1C).

**Figure 1.**

**The inhibitory nanobody NB7 binds tightly to all p110 $\gamma$  complexes and inhibits kinase activity, but does not prevent membrane binding**

**A.** Cartoon schematic depicting nanobody inhibition of activation by lipidated G $\beta\gamma$  (1.5  $\mu$ M final concentration) on 5% PIP<sub>2</sub> membrane (5% phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>), 95% phosphatidylserine (PS)) activation. Lipid kinase assays showed a potent inhibition of lipid kinase activity with increasing concentrations of NB7 (3-3000 nM) for the different complexes. Experiments are carried out in triplicate (n=3) with



each replicate shown. The y-axis shows lipid kinase activity normalised for each complex activated by G $\beta\gamma$  in the absence of nanobody. Concentrations of each protein were selected to give a lipid kinase value in the detectable range of the

ATPase transcriber assay. The protein concentration of p110 $\gamma$  (300 nM), p110 $\gamma$ -p84 (330 nM) and p110 $\gamma$ -p101 (12 nM) was different due to intrinsic differences of each complex to be activated by lipidated G $\beta$  $\gamma$  and is likely mainly dependent for the difference seen in NB7 response.

**B.** Association and dissociation curves for the dose response of His-NB7 binding to p110 $\gamma$ , p110 $\gamma$ -p84 and p110 $\gamma$ -p101 (50 – 1.9 nM) is shown. A cartoon schematic of BLI analysis of the binding of immobilized His-NB7 to p110 $\gamma$  is shown on the left. Dissociation constants ( $K_D$ ) were calculated based on a global fit to a 1:1 model for the top three concentrations and averaged with error shown. Error was calculated from the association and dissociation value ( $n=3$ ) with standard deviation shown. Full details are present in the source data.

**C.** Association and dissociation curves for His-NB7 binding to p110 $\gamma$ , p110 $\alpha$ -p85 $\alpha$ , p110 $\beta$ -p85 $\beta$ , and p110 $\delta$ -p85 $\beta$ . Experiments were performed in duplicate with a final concentration of 50 nM of each class I PI3K complex.

**D.** Effect of NB7 on PI3K recruitment to supported lipid bilayers containing H-Ras (GTP) and farnesyl-G $\beta$  $\gamma$  as measured by Total Internal Reflection Fluorescence Microscopy (TIRF-M). DY647-p84/p110 $\gamma$  displays rapid equilibration kinetics and is insensitive to the addition of 500 nM nanobody (black arrow, 250 sec) on supported lipid bilayers containing H-Ras (GTP) and farnesyl-G $\beta$  $\gamma$ .


**E.** Kinetics of 50 nM DY647-p84/p110 $\gamma$  membrane recruitment appears indistinguishable in the absence and presence of nanobody. Prior to sample injection, DY647-p84/p110 $\gamma$  was incubated for 10 minutes with 500 nM nanobody.

**F.** Representative TIRF-M images showing the localization of 50 nM DY647-p84/p110 $\gamma$  visualized in the absence or presence of 500 nM nanobody (+NB7). Membrane composition for panels C-E: 93% DOPC, 5% DOPS, 2% MCC-PE, Ras (GTP) covalently attached to MCC-PE, and 200 nM farnesyl-G $\beta$  $\gamma$ .

To further understand the mechanism by which this nanobody blocked lipid kinase activity we measured the bulk membrane recruitment dynamics of fluorescently labeled Dy647-p84-p110 $\gamma$  on supported lipid bilayers (SLBs) using Total Internal Reflection Fluorescence (TIRF) Microscopy. We found that the nanobody had no effect on membrane recruitment of p110 $\gamma$ -p84 on bilayers containing membrane-tethered Ras (GTP) and G $\beta$  $\gamma$  (Fig. 1D-F). Membrane binding was not affected when the nanobody was spiked into samples containing membrane associated Dy647-p84-p110 $\gamma$  (Fig 1D). Similarly, pre-incubation of Dy647-p84-p110 $\gamma$  with 500 nM NB7 did not perturb membrane association of the kinase when flowed over a supported membrane (Fig. 1E).

We wanted to define the molecular mechanism of how nanobody NB7 was a potent allosteric inhibitor of lipid kinase activity. We purified the complex of nanobody NB7 bound to p110 $\gamma$ -p84 to homogeneity by gel filtration. Using this sample, we obtained a cryo-EM reconstruction of the complex of nanobody (NB7)-bound p110 $\gamma$  at 3.0-Å overall resolution from 149,603 particles (Figs. 2A-D, S1-S2 and supplemental table 1). The density map was of sufficient quality to perform automated and manual construction of the p110 $\gamma$ -NB7 complex, with unambiguous building of the interfacial contacts between NB7 and p110 $\gamma$  (Fig. S2). Nanobody binding did not induce any large-scale conformational changes of p110 $\gamma$ , as the structure of p110 $\gamma$  bound to NB7 was similar to the apo p110 $\gamma$  crystal structure or p110 $\gamma$ -p101 cryo-EM structure (Fig. S3). The lowest local resolution was in the ABD domain, with increased B-factors of the ABD in the p110 $\gamma$ -NB7 structure compared to p110 $\gamma$ -p101 (Fig. S3). This is consistent with the concept that ABD flexibility plays an important role in class I PI3K regulation (Liu et al., 2022).





**Figure 2.**

## Structure of p110 $\gamma$ bound to inhibitory nanobody NB7

**A.** Domain schematics of p110 $\gamma$  with helical domain (blue), activation loop (orange), and regulatory motif (green) of p110 $\gamma$  annotated.

**B.** Cryo EM density of the p110 $\gamma$ -NB7 complex colored according to the schematic in **A**.

**C.** Cartoon model of the structure of p110 $\gamma$  bound to NB7 colored according to **A**.

**D.** Schematic depicting the key features of p110 and the nanobody binding site, colored according to panel **A**.

**E.** Domain schematic of NB7 CDR regions and their sequences.

**F.** Zoom in on the binding interface of NB7, with the CDRs colored as in panel E, and the electron density of the CDR regions contoured at 3 $\sigma$  (blue mesh).

The interface between NB7 and p110 $\gamma$  was extensive, with ~1200 Å<sup>2</sup> of buried surface area, with interactions of the ABD-RBD linker, N-terminus of the helical domain, and the regulatory motif at the turn between  $\alpha$ 8- $\alpha$ 9 (1022-1026aa). This location in the regulatory motif is where both activating oncogenic (R1021C) and inhibitory loss of function mutants have been identified (R1021P) (Takeda et al., 2019), as well as a putative inhibitory phosphorylation site (T1024) (Perino et al., 2011). The resolution was sufficient to unambiguously build the three complementarity determining region (CDR) loops of NB7 that mediate target selectivity (Fig. 2E-F). The interface is primarily hydrophobic, with only 8 hydrogen bonds, and 1 electrostatic interaction among the 33 interfacial residues of NB7. A pocket formed between the helical domain and the ABD-RBD linker forms the majority of the interface, with extensive interactions with the long CDR3. The CDR1 loop packed up against the N-terminal section of the helical domain, with the CDR2 loop forming the interface with the regulatory motif. Previous study of oncogenic mutants in the regulatory motif of p110 $\gamma$  showed that increased dynamics mediated by these mutants increased kinase activity, putatively by breaking the autoinhibitory tryptophan lock in  $\alpha$ 12 of the regulatory motif (Rathinaswamy et al., 2021c). Therefore, rigidifying the regulatory motif likely explains the molecular basis for how it prevents kinase activity. The nanobody interface is distinct from the predicted G $\beta$  $\gamma$  interface (Rathinaswamy et al., 2023) and the experimentally resolved Ras interface (Pacold et al., 2000), explaining why it can still be membrane recruited by these stimuli.

The interface of NB7 with p110 $\gamma$  is distant from both the putative membrane binding surface, as well as the catalytic machinery of the kinase domain. To further understand how

this nanobody could potentially inhibit PI3K activity we examined any other potential modulators of PI3K activity localised in this region. There are two regulatory phosphorylation sites in the helical (Walser et al., 2013) and kinase domain (Perino et al., 2011) localised at the NB7 interface. This is intriguing as helical domain phosphorylation is activating, and kinase domain phosphorylation is inhibitory. This suggested a critical role in the regulation of p110 $\gamma$  is the dynamics of this kinase-helical interface. To fully define the role of NB7 in altering the dynamics of the helical domain we needed to study other modulators of helical domain dynamics.

## p110 $\gamma$ activation by helical domain phosphorylation


To further understand the potential role of helical domain dynamics in controlling p110 $\gamma$  activity we examined the role of S582 helical domain phosphorylation by the protein kinase PKC $\beta$ II (encoded by the gene *PRKCB2*, referred to as PKC $\beta$  for simplicity throughout the manuscript) (Walser et al., 2013). S582 is located on the interior of the helical domain and would not be expected to be exposed when the N-terminal region of the helical domain is folded (Fig. 3A). To understand this better at a molecular level, we purified a catalytic fragment of PKC $\beta$  and performed protein phosphorylation reactions on p110 $\gamma$  apo, p110 $\gamma$ -p84, and p110 $\gamma$ -p101. We identified a phosphorylated peptide containing S582, and surprisingly, we found an additional p110 $\gamma$  phosphorylation site at either S594/S595 (Fig. 3B, Fig. S4). The modification at this site results in a single phosphorylation event, but due to CID MS/MS fragmentation we cannot determine which site is modified, and we will refer it as S594/S595 throughout the manuscript. The S594/S595 site is also located in the N-terminal region of the helical domain, and is even more buried than S582, and would not be expected to be exposed when this region is folded (Fig. 3A). Dose response curves of PKC $\beta$  treatment was carried out for p110 $\gamma$  (Fig. 3C), p110 $\gamma$ -p84 (Fig. 3D), and p110 $\gamma$ -p101 (Fig. 3E). Both p110 $\gamma$  and p110 $\gamma$ -p84 showed similar dose response curves for PKC $\beta$  treatment, with similar curves for S582 and S594/S595. The p110 $\gamma$ -p101 complex was only very weakly phosphorylated, with <100-fold lower levels compared to p110 $\gamma$  and p110 $\gamma$ -p84 (Fig. 3E). This is consistent with the helical domain in p110 $\gamma$  being more rigid when bound to p101, compared to either bound to p84 or p110 $\gamma$  alone.

**Figure 3.**

**PKC $\beta$  leads to dual phosphorylation of internal sites in the helical domain, with selectivity for apo p110 $\gamma$  and p110 $\gamma$ -p84 over p110 $\gamma$ -p101**

**A.** Putative phosphorylation sites mapped on the structure of p110 $\gamma$  (PDB: 7MEZ) and cartoon schematic. The regions are colored based on domain schematics featured in Fig 2A.


**B.** Raw MS spectra of the unphosphorylated and phosphorylated peptide for a region spanning 579-592 (RYESLKHPKAYPKL) and 593-607



(FSSVKWGQQEIVAKT). The putative phosphorylation sites in the sequence are shown in red, with the  $m/z$  theoretical ( $m/z$  t) and  $m/z$  experimental ( $m/z$  e) shown below each sequence.

**C-E.** Extracted traces and ratios of the intensity of extracted ion traces of different phosphorylation site peptides (Top to bottom: S594/S595 and S582) from (C) p110 $\gamma$ , (D) p110 $\gamma$ /p84 or (E) p110 $\gamma$ /p101 samples treated with increasing concentration of PKC $\beta$  according to the legend. The black traces in the ratio graphs are the intensity of the non-phosphorylated peptide, and the red traces in the ratio graphs are the intensity of the phosphorylated peptide.

To provide additional insight into the molecular mechanisms underlying p110 $\gamma$  phosphorylation we carried out hydrogen deuterium exchange mass spectrometry (HDX-MS) experiments on p110 $\gamma$  and phosphorylated p110 $\gamma$  (90.8% phosphorylated S594/595, 92% phosphorylated S582) (Fig. 4A). The full data underlying the experiment is available in the source data, and data processing information is in supplemental table 2. We have previously observed that the N-terminal region of the helical domain of apo p110 $\gamma$  (residues spanning 557-630aa) shows isotope profiles that are consistent with EX1 H/D exchange kinetics (Rathinaswamy et al., 2021b; 2021c; Vadas et al., 2013; Walser et al., 2013). This is indicative of cooperative unfolding of extended protein regions, with H/D exchange occurring faster than the refolding event. This region is where the PKC $\beta$  phosphorylation sites are located and may explain how the buried residues S582 and S594/S595 can be exposed to PKC $\beta$ . This is compatible with the observation that p110 $\gamma$ -p101 is protected from phosphorylation, as it does not show EX1 kinetics in this region, whereas both p110 $\gamma$  and p110 $\gamma$ -p84 do (Rathinaswamy et al., 2021a).



**Figure 4.**

## Activating phosphorylation at the helical domain leads to opening of the regulatory motif

**A.** HDX-MS comparing apo and phosphorylated p110 $\gamma$ . Significant differences in deuterium exchange are mapped on to the structure and cartoon of p110 $\gamma$  according to the legend (PDB: 7MEZ).

**B.** The graph of the #D difference in deuterium incorporation for p110 $\gamma$ , with each point representing a single peptide. Peptides colored in red are those that had a significant change in the mutants (greater than 0.4 Da and 5% difference at any timepoint, with a two tailed t-test  $p < 0.01$ ). Error bars are S.D. ( $n=3$ ).

**C.** Representative bimodal distribution (EX1 kinetics) observed in the helical domain peptides of p110 $\gamma$ .



**D.** Representative p110 $\gamma$  peptides displaying increases in exchange in the phosphorylated state are shown. For all panels, error bars show SD ( $n=3$ )


**E.** Measurement of ATP to ADP conversion of phosphorylated and non-phosphorylated p110 $\gamma$  (1000 nM final concentration) ATPase activity in the absence (left) and presence of PIP<sub>2</sub> membranes (5% phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>), 95% phosphatidylserine (PS)) activation (right). Significance is indicated by \*\*(<0.001%), and \*\*\*(<0.0001%).

When we compared phosphorylated p110 $\gamma$  (>90.8% as measured by mass spectrometry at both sites) to unphosphorylated p110 $\gamma$  we observed extensive increases in dynamics in the C2, helical domain and kinase domain (Fig. 4A-D). The largest increases in exchange upon phosphorylation were located in the N-terminal region of the helical domain, with the peptides directly adjacent to the phosphorylation site showing almost complete deuterium incorporation at the earliest time points of exchange. This is indicative of significant disruption of the alpha helical secondary structure in this region. When we examined the exchange profiles in this region, they still underwent EX1 kinetics (Fig. 4C), however, phosphorylated p110 $\gamma$  was enriched in the more fully deuterated species. In addition to the regions in the helical domain, a portion of the regulatory motif of the kinase domain also showed increased deuterium exposure. This included the  $\alpha 9$ - $\alpha 12$  helices that surround the activation loop of p110 $\gamma$ . These increases in exchange were similar to those we had observed in a R1021C oncogenic activating mutant of *PIK3CG* (Rathinaswamy et al., 2021c).

To further explore the potential role of phosphorylation in mediating p110 $\gamma$  activity, we examined the kinase activity of p110 $\gamma$  under two conditions: basal ATP turnover, and with PIP<sub>2</sub> containing lipid membranes. The experiments in the absence of PIP<sub>2</sub> measure turnover of ATP into ADP and phosphate and is a readout of basal catalytic competency. Experiments with PIP<sub>2</sub>, measured ATP consumed in the generation of PIP<sub>3</sub>, as well as in non-productive ATP turnover. The p110 $\gamma$  enzyme in the absence of stimulators is very weakly active towards PIP<sub>2</sub> substrate with only ~2 fold increased ATP turnover compared to in the absence of membranes. This is consistent with very weak membrane recruitment of p110 $\gamma$  complexes in the absence of lipid activators (Rathinaswamy et al., 2023). PKC $\beta$ -mediated phosphorylation enhanced the ATPase activity of p110 $\gamma$  ~2-fold in both the absence and presence of membranes (Fig. 4E). This suggests that the effect of phosphorylation is to change the intrinsic catalytic efficiency of phosphorylated p110 $\gamma$ , with limited effect on membrane binding.

## Nanobody decreases p110 $\gamma$ phosphorylation

As NB7 bound at the interface of the helical and kinase domains that is exposed upon PKC $\beta$  phosphorylation of p110 $\gamma$  we hypothesized that the nanobody would likely alter phosphorylation. We carried out PKC $\beta$  phosphorylation of p110 $\gamma$ , p110 $\gamma$ -p101 and p110 $\gamma$  bound to NB7. The presence of NB7 showed a significant decrease in p110 $\gamma$  phosphorylation at both sites (Fig. 5A-B). We also wanted to determine whether p110 $\gamma$  phosphorylation reciprocally perturbed NB7 binding. BLI experiments showed that there was no detectable binding of NB7 to phosphorylated p110 $\gamma$  (Fig. 5C-D), consistent with phosphorylation disrupting the N-terminal region of the p110 $\gamma$  helical domain. In addition, lipid kinase assays using phosphorylated p110 $\gamma$  showed no detectable difference in activity when measured in the absence and presence nanobody (Fig. 5E), consistent with NB7 being unable to bind to phosphorylated p110 $\gamma$ .



**Figure 5.**

## Nanobody NB7 blocks PKCβ phosphorylation, and phosphorylation prevents nanobody binding

**A.** Extracted ion chromatograms for p110γ, p110γ-p101, and p110γ bound to NB7 are shown for the S594 or S595 phosphorylation site in p110γ. A bar graph showing the intensities of phosphorylated and non-phosphorylated p110γ peptide (593-607) for p110γ (black), p110γ with NB7 (red) and p110γp101 (purple) are shown to the right of the extracted ion chromatograms (n=3, right). In all experiments in panels **A+B**, PKCβ was present at 500 nM. Significance is indicated by \*\*\*(<0.0001%).

**B.** Extracted ion chromatograms for p110γ, p110γ-p101, and p110γ bound to NB7 are shown for the S582 phosphorylation site in p110γ. A bar graph showing the intensities of phosphorylated and non-phosphorylated p110γ peptide (579-592) p110γ (black), p110γ with NB7 (red) and p110γ-p101 (purple) are shown to the right of the extracted ion chromatograms (n=3, right). Significance is indicated by \* (<0.01%), and \*\*\*(<0.0001%). The putative phosphorylation site is shown in red in the sequence above the bar graphs for both panel **A+B**.

**C.** Cartoon schematic of BLI analysis of the binding of immobilized His-NB7 to phosphorylated and non-phosphorylated p110γ.

**D.** Association curves for phosphorylated and non-phosphorylated p110γ (25nM) binding to His-NB7 are shown (n=3).


**E.** ATPase kinase activity assays comparing the activation/inhibition of phosphorylated and non-phosphorylated p110γ (1000 nM) with or without nanobody (3000 nM final) in the absence of PIP<sub>2</sub> membranes. Significance is indicated by \* (<0.05%), and NS (>0.05%).

## Discussion

Here we find that the helical domain is a central regulator of the p110γ catalytic subunit of class IB PI3K, with modulation of helical dynamics through binding partners or PTMs able to

either increase or decrease lipid kinase activity. These results expand on previous work defining the helical domain as a central regulator of class IA PI3Ks, where the nSH2 domain of the p85 regulatory subunits makes inhibitory interactions that significantly inhibit lipid kinase activity of all class IA catalytic subunits (p110 $\alpha$ , p110 $\beta$ , and p110 $\delta$ ) (Mandelker et al., 2009; Miled et al., 2007; Burke and Williams, 2013; Burke, 2018). This inhibitory interaction in class IA PI3Ks is disrupted in human cancers (helical hotspot mutations in *PIK3CA*) (Samuels et al., 2004) and immune disorders (helical mutations in *PIK3CD* in APDS1) (Angulo et al., 2013; Lucas et al., 2014). Class IB PI3Ks are unique compared to class IA PI3Ks, as they are not inhibited by p101 and p84 regulatory subunits, but instead potentiate GPCR activation. This lack of inhibition is due to the distinct binding interface of class IB PI3K regulatory subunits compared to class IA regulatory subunits, with only class IA regulatory subunits making direct inhibitory interactions with the kinase and helical domains of p110 catalytic subunits (Rathinaswamy et al., 2021a). Here we show that a unique surface at the interface of the helical and kinase domains of p110 $\gamma$  is a potential site for the development of novel allosteric inhibitors that modulate p110 $\gamma$  activity.

The previously identified inhibitory nanobody (NB7) (Rathinaswamy et al., 2021b) bound with high affinity and inhibited all complexes of p110 $\gamma$ . The nanobody interface is distinct from how the nSH2 inhibits class IA PI3K activity, as its binding site is on the opposite face of the helical domain (Fig. 6A). The mechanism of inhibition is also distinct, as the nSH2-helical interaction plays a critical role in preventing membrane recruitment of inhibited class I PI3Ks, with removal of this interface either through pYXXM motif binding, or oncogenic mutations leading to increased membrane recruitment (Burke et al., 2012; 2011; Zhang et al., 2011). Analysis of the nanobody binding site compared to the structure of HRas-p110 $\gamma$  or the HDX-MS supported AlphaFold-multimer prediction of G $\beta$  $\gamma$ -p110 $\gamma$  (Rathinaswamy et al., 2023) shows that nanobody binding does not sterically block complex formation (Fig. 6B). This is consistent with it not blocking membrane recruitment by Ras/G $\beta$  $\gamma$ . The nanobody inhibited ATP turnover both in solution and on membranes, suggesting that it prevents formation of a catalytically competent conformation of p110 $\gamma$ , but still allows for membrane recruitment. Further development of small molecule allosteric binding partners in this allosteric pocket between the kinase and helical domain may reveal the specific molecular interactions in this pocket that mediate inhibition.



**Figure 6.**

## Comparison of nanobody binding site compared to p85 inhibition of class IA PI3Ks and class IB activation sites

**A.** Comparison of the nanobody NB7 binding site in p110 $\gamma$  compared to the nSH2 inhibitory site in p110 $\alpha$  (PDB: 3HHM) (Mandelker et al., 2009)

**B.** Comparison of the nanobody NB7 binding site in p110 $\gamma$  compared to the X-ray structure of the Ras binding site (PDB: 1HE8) (Pacold et al., 2000) and the AlphaFold model of G $\beta$  $\gamma$  bound to p110 $\gamma$  (Rathinaswamy et al., 2023).

**C.** Oncogenic mutations and post-translational modifications in

spatial proximity to the nanobody binding site.

Oncogenic mutations are frequent in the class IA PI3K $\alpha$  encoded by *PIK3CA*, with this being the 2<sup>nd</sup> most frequently mutated gene in human cancer (Lawrence et al., 2014). Mutations in p110 $\gamma$  encoded by *PIK3CG* in cancer are less frequent, however, they can still provide insight into regulatory mechanisms that control activity. Oncogenic mutations in the kinase domain (R1021C) and helical domain (E581K) are in close proximity to the nanobody binding site, and both would be expected to disrupt the stability of the helical domain or regulatory motif of the kinase domain (Fig. 6C). In addition to these mutations there are also multiple post-translational modifications that occur in this region, including inhibitory phosphorylation at T1024 (Perino et al., 2011), and activating phosphorylation at S582 (Walser et al., 2013). PKC $\beta$  is activated downstream of the IgE receptor in mast cells (Walser et al., 2013), but the full details of how this activates PI3K has been unclear. We identified an additional PKC $\beta$  phosphorylation site located in the helical domain (S594/S595) (Fig. 6C). Both the S582 and S594/S595 sites are not surface accessible and would require a transient opening of the helical domain for kinase accessibility. HDX-MS analysis of the helical domain of p110 $\gamma$  has shown that it is more dynamic than other class I PI3K isoforms (Burke and Williams, 2013; Walser et al., 2013), with the presence of the p101 regulatory subunit dramatically decreasing helical domain dynamics (Vadas et al., 2013). This putative mechanism of helical domain dynamics driving PKC $\beta$  phosphorylation is consistent with our observation that p101 subunits decreased p110 $\gamma$  phosphorylation >100-fold. PKC $\beta$  phosphorylation of p110 $\gamma$  leads to increased dynamics in both the helical and kinase domains with increased kinase activity, although only weakly compared to full activation by either membrane localised Ras or G $\beta$  $\gamma$ . This increase was observed with both membrane and soluble substrate, so likely is not driven by altered membrane recruitment.

Overall, our biophysical and biochemical analysis of modulators of helical domain dynamics reveal the critical role of this domain in regulating class IB PI3K activity. This raises possibilities for development of small molecule modulators that may either increase or decrease helical domain dynamics, leading to either activation or inhibition. The high-resolution structure of an allosteric inhibitor nanobody provide initial insight into which pockets can specifically be targeted. Multiple ATP competitive p110 $\gamma$  selective inhibitors are in clinical trials for human cancers (Li et al., 2021), with many having significant side effects. The identification of novel inhibitory strategies provides new opportunities for targeting p110 $\gamma$  dysregulation in human disease.

## Acknowledgements

J.E.B. is supported by the Canadian Institute of Health Research (CIHR, 168998), and the Michael Smith Foundation for Health Research (MSFHR, scholar 17686). C.K.Y. is supported by CIHR (FDN-143228, PJT-168907) and the Natural Sciences and Engineering Research Council of Canada (RGPIN-2018-03951). S.D.H. is supported by an NSF CAREER Award (MCB-2048060). This research project was supported in part by the UBC High Resolution Macromolecular Cryo-Electron Microscopy Facility (HRMEM). A portion of this research was supported by NIH grant U24GM129547 and performed at the PNCC at OHSU and accessed through EMSL (grid.436923.9), a DOE Office of Science User Facility sponsored by the Office of Biological and Environmental Research. We appreciate help from Theo Humphreys and Rose Marie Haynes with data collection at PNCC. Competing Interests: The authors declare that they have no competing interests.



## Conflict of Interest statement

JEB reports personal fees from Scorpion Therapeutics, Reactive therapeutics and Olema Oncology; and research grants from Novartis. Other authors declare no competing interests.

## Methods

Resources table	SOURCE	IDENTIFIER
<b>Bacterial and virus strains</b>		
E.coli XL10-GOLD KanR Ultracompetent Cells	Agilent	200317
E.coli DH10EMBacY Competent Cells	Geneva Biotech	DH10EMBacY
<b>Chemicals, peptides, and recombinant proteins</b>		
Deuterium Oxide 99.9%	Sigma	151882
Guanosine 5'-diphosphate (GDP) sodium salt hydrate	Sigma	G7127-100MG
Guanosine 5'-triphosphate (GTP) sodium salt hydrate	Sigma	G8877-250MG
Sodium deoxycholate	Sigma	D6750
Polyoxyethylene (10) lauryl ether	Sigma	P9769
CHAPS, Molecular Biology Grade	EMD Millipore	220201
Phosphatidylserine (Porcine Brain)	Avanti	840032C
Phosphatidylethanolamine (Egg yolk)	Sigma	P6386
Cholesterol	Sigma	47127-U
Phosphatidylcholine (Egg yolk)	Avanti	840051C
Phosphatidylinositol-4,5-bisphosphate (Porcine Brain)	Avanti	840046
Sphingomyelin (Egg yolk)	Sigma	S0756
1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC)	Avanti	850375C
1,2-dioleoyl-sn-glycero-3-phospho-L-serine (18:1, DOPS)	Avanti	840035C
1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-[4-(p-maleimidomethyl)cyclohexane-carboxamide] (18:1 MCC-PE)	Avanti	780201C
10 mg/mL beta casein solution	ThermoFisher	37528
10x PBS [pH 7.4]	Corning	46-013-CM
glucose oxidase from <i>Aspergillus niger</i> (225 U/mg)	Biophoretics	B01357.02
catalase	Sigma	C40-100MG Bovine Liver
Trolox	Cayman	10011659

	Chemicals	
Dyomics 647 maleimide dye	Dyomics	647P1-03
Coenzyme A	Sigma	C3019
Sulfuric acid	Sigma	58105-2.5L-PC
<b>Critical commercial assays</b>		
Transcreener ADP2 FI Assay (1,000 Assay, 384 Well)	BellBrook Labs	3013-1K
<b>Deposited data</b>		
PDB coordinate file for p110 $\gamma$ -NB7 structure	PDB	8DP0
EM density file for p110 $\gamma$ -NB7 complex	EMD	EMD-27627
HDX-MS and phosphorylation proteomics data	PRIDE	PXD040765
<b>Oligonucleotides</b>		
Fwd primer for amplifying KD of PKC $\beta$ II GTATTTTCAGGGCgcccgtaccACGACCAACTGTCT CCAAATTTG	Sigma	MR51F
Rvs primer for amplifying KD of PKC $\beta$ II gactcgagcggccgcTTATAGCTCTTGACTTCGGGTTTAA AAAATTCAG	Sigma	MR51R
Fwd primer for amplifying N term of PKC $\beta$ II CCATCACggatctggcgtagtATGGCTGACCCGGCTGCG	Sigma	MR52F
Rvs primer for amplifying N term of PKC $\beta$ II GCCCTGAAAATACAGGTTTTCTTTTCTTCCGGGAC CTTGGTTCCC	Sigma	MR52R
Fwd primer for adding stop codon to PKC $\beta$ II AGTCAAGAGCTAAgcccgcgctcagcttagagcctgc	Sigma	MR56F
Rvs primer for adding stop codon to PKC $\beta$ II gactcgagcggccgcTTAGCTCTTGACTTCGGGTTTAA AATTCAG	Sigma	MR56R
<b>Recombinant DNA</b>		
pMultiBac-G $\beta$ 1/G $\gamma$ 2	PMID:34452907	pOP737
pACEBac1-hsp110 $\gamma$	PMID:34452907	MR30
pMultiBac-hsp110 $\gamma$ -ssp101	PMID:34452907	MR22
pMultiBac-hsp110 $\gamma$ -mmp84	PMID:34452907	MR24
pFastBac HRas G12V	PMID:34452907	BS9
biGBac hsp110 $\gamma$ /ybbr-hsp84	PMID:36842083	HP28
biGBac hsp110 $\gamma$ /ybbr-hsp101	PMID:36842083	HP29
his6-GST-PrescissionProtease-SNAP-RBD(K65E)	PMID:34452907	pSH936
his6TEV-HRas(1-184aa) C118S, C181S	PMID:34452907	pSH414
his6-G $\gamma$ 2, SNAP-G $\beta$ 1 (DUAL FastBac)	PMID:34452907	pSH651
pACEBAC-PKC $\beta$ II (internal tev cleavage site)	This paper	pMR56
pFASTBac p110 $\alpha$	PMID: 28515318	pOV1181
pFASTBac p110 $\beta$	PMID: 28515318	pOV1182
pFASTBac p110 $\delta$	PMID: 28515318	pOV1183
pFASTBac p85 $\beta$	This paper	EX21
<b>Software and algorithms</b>		
COOT-0.9.4.1	CCP4	<a href="https://www2.mrc-lmb.cam.ac.uk/personal/pemsley/coot/">https://www2.mrc-lmb.cam.ac.uk/personal/pemsley/coot/</a>

Phenix-1.19.1	Open source	<a href="https://www.phenix-online.org/">https://www.phenix-online.org/</a>
PDBEPIISA (Proteins, Interfaces, Structures and Assemblies)	EMBL-EBI	<a href="https://www.ebi.ac.uk/pdbe/pisa/pistart.html">https://www.ebi.ac.uk/pdbe/pisa/pistart.html</a>
ESPrnt 3.0	Robert et al NAR 2014	<a href="https://esprnt.ibcp.fr">https://esprnt.ibcp.fr</a>
HDEaminer	Sierra Analytics	<a href="http://massspec.com/hdexaminer">http://massspec.com/hdexaminer</a>
GraphPad Prism 7	GraphPad	<a href="https://www.graphpad.com">https://www.graphpad.com</a>
PyMOL	Schroedinger	<a href="http://pymol.org">http://pymol.org</a>
Compass Data Analysis	Bruker	<a href="https://www.bruker.com">https://www.bruker.com</a>
ChimeraX	UCSF	<a href="https://www.rbvi.ucsf.edu/chimera/">https://www.rbvi.ucsf.edu/chimera/</a>
ImageJ/Fiji	ImageJ	<a href="https://imagej.net/software/fiji/">https://imagej.net/software/fiji/</a>
Nikon NIS elements	Nikon	<a href="https://www.microscope.healthcare.nikon.com/products/software/nis-elements">https://www.microscope.healthcare.nikon.com/products/software/nis-elements</a>
cryoSPARC v.3.3.2	Structura Biotechnology	<a href="https://cryosparc.com/">https://cryosparc.com/</a>
<b>Other</b>		
Sf9 insect cells for expression	Expression Systems	94-001S
Insect cell media	Expression Systems	96-001-01
Hellmanex III cleaning solution	Fisher	14-385-864
6-well sticky-side chamber	IBIDI	80608
C-Flat 2/2-T grids	Electron Microscopy Sciences	CFT-223C

## Plasmid Generation

Plasmids encoding Homo sapiens p110 $\gamma$  (human), Mus musculus p84 (mouse), Sus scrofa p101 (porcine), and G $\beta$ y were used as previously described (Rathinaswamy et al., 2023; 2021a). The plasmids encoding the class IA PI3Ks were also used as previously described (Dornan et al., 2017; Siempelkamp et al., 2017). The pDONR223-PRKCB2 (PKC $\beta$ II, uniprot identifier: P05771-2) was a gift from William Hahn & David Root (Addgene plasmid #23746; <http://n2t.net/addgene:23746>; RRID:Addgene\_23746) (Johannessen et al., 2010). The PKC $\beta$ II construct contains an internal TEV site that cleaves the catalytic domains from the C1/C2 regulatory domains (TEV site inserted between residues 320 and 321 of PKC $\beta$ ). This construct was subcloned into a pACEBAC Sf9 expression vector for Sf9 protein production. All constructs were cloned to include a 10 $\times$  histidine tag, a 2 $\times$  strep tag, and a tobacco etch virus protease cleavage site on the N terminus. For p110 $\gamma$  and PKC $\beta$ II this tag was included at the N-terminus, with this tag included at the N-terminus of either p84 or p101 for purification of p110 $\gamma$ -p101 and p110 $\gamma$ -p84. Full details of the plasmids are included in the resource table.

## Virus Generation and Amplification

The plasmids encoding genes for insect cell expression were transformed into DH10MultiBac cells (MultiBac, Geneva Biotech) to generate bacmid containing the genes of interest. Successful generation was identified by blue-white colony screening and the bacmid was purified using a standard isopropanol-ethanol extraction method. Bacteria were grown overnight (16 hours) in 3-5 mL 2xYT (BioBasic #SD7019). Cells were spun down and the pellet was resuspended in 300  $\mu$ L of 50 mM Tris-HCl, pH 8.0, 10 mM EDTA, 100 mg/mL RNase A. The pellet was lysed by the addition of 300  $\mu$ L of 1% sodium dodecyl sulfate (SDS) (W/V), 200 mM NaOH, and the reaction was neutralized by addition of 400  $\mu$ L of 3.0 M potassium acetate, pH 5.5. Following centrifugation at 21130 RCF and 4  $^{\circ}$ C (Rotor #5424 R), the supernatant was

mixed with 800  $\mu$ L isopropanol to precipitate bacmid DNA. Following centrifugation, the pelleted bacmid DNA was washed with 500  $\mu$ L 70% Ethanol three times. The pellet was then air dried for 1 minute and re-suspended in 50  $\mu$ L Buffer EB (10 mM Tris-Cl, pH 8.5; All buffers from QIAprep Spin Miniprep Kit, Qiagen #27104). Purified bacmid was then transfected into Sf9 cells. 2 mL of Sf9 cells at  $0.6 \times 10^6$  cells/mL were aliquoted into a 6-well plate and allowed to attach to form a confluent layer. Transfection reactions were prepared mixing 8-12  $\mu$ g of bacmid DNA in 100  $\mu$ L 1xPBS and 12  $\mu$ g polyethylenimine (Polyethylenimine “Max” MW 40,000, Polysciences #24765, USA) in 100  $\mu$ L 1xPBS and the reaction was allowed to proceed for 20-30 minutes before addition to an Sf9 monolayer containing well. Transfections were allowed to proceed for 5-6 days before harvesting virus containing supernatant as a P1 viral stock.

Viral stocks were further amplified by adding P1 to Sf9 cells at  $\sim 2 \times 10^6$  cells/mL (2/100 volume ratio). This amplification was allowed to proceed for 4-5 days and resulted in a P2 stage viral stock that was used in final protein expression. Harvesting of P2 viral stocks was carried out by centrifuging cell suspensions in 50 mL Falcon tubes at 2281 RCF (Beckman GS-15). To the supernatant containing virus, 5-10% inactivated fetal bovine serum (FBS; VWR Canada #97068-085) was added and the stock was stored at 4°C.

## Expression and purification of PI3Ky, PI3K $\alpha/\beta/\delta$ and PKC $\beta$ constructs

PI3Ky and PKC $\beta$  constructs were expressed in Sf9 insect cells using the baculovirus expression system. Following 55 hours after infection with P2 viral stocks, cells were harvested by centrifuging at 1680 RCF (Eppendorf Centrifuge 5810 R) and the pellets were snap-frozen in liquid nitrogen. The complex was purified through a combination of nickel affinity, streptavidin affinity and size exclusion chromatography.

Frozen insect cell pellets were resuspended in lysis buffer (20 mM Tris pH 8.0, 100 mM NaCl, 10 mM imidazole pH 8.0, 5% glycerol (v/v), 2 mM  $\beta$ ME), protease inhibitor (Protease Inhibitor Cocktail Set III, Sigma) and sonicated for 2 minutes (15s on, 15s off, level 4.0, Misonix sonicator 3000). Triton-X was added to the lysate to a final concentration of 0.1% and clarified by spinning at 15,000 RCF at 4°C for 45 minutes (Beckman Coulter JA-20 rotor). The supernatant was loaded onto a 5 mL HisTrap™ FF crude column (GE Healthcare) equilibrated in NiNTA A buffer (20 mM Tris pH 8.0, 100 mM NaCl, 20 mM imidazole pH 8.0, 5% (v/v) glycerol, 2 mM  $\beta$ ME). The column was washed with high salt NiNTA A buffer (20 mM Tris pH 8.0, 1 M NaCl, 20 mM imidazole pH 8.0, 5% (v/v) glycerol, 2 mM  $\beta$ ME), NiNTA A buffer, 6% NiNTA B buffer (20 mM Tris pH 8.0, 100 mM NaCl, 250 mM imidazole pH 8.0, 5% (v/v) glycerol, 2 mM  $\beta$ ME) and the protein was eluted with 100% NiNTA B. The eluent was loaded onto a 5 mL StrepTrap™ HP column (GE Healthcare) equilibrated in gel filtration buffer (20mM Tris pH 8.5, 100 mM NaCl, 50 mM Ammonium Sulfate and 0.5 mM TCEP). To purify PI3K $\alpha/\beta/\delta$ , the purification protocol was performed as described above but instead the protein was eluted in PI3K $\alpha$  gel filtration buffer (20mM HEPES 7.5, 150mM NaCl, 0.5mM TCEP). The column was washed with the corresponding gel filtration buffer and loaded with tobacco etch virus protease. After cleavage on the column overnight, the PI3Ky protein constructs were eluted in gel filtration buffer. The protein was concentrated in a 50,000 MWCO Amicon Concentrator (Millipore) to <1 mL and injected onto a Superdex™ 200 10/300 GL Increase size-exclusion column (GE Healthcare) equilibrated in gel filtration buffer. After size exclusion, the protein was concentrated, aliquoted, frozen, and stored at -80°C. For PKC $\beta$ , the protein was eluted from the strep column in gel filtration buffer, and the eluate was then loaded on a 1ml HisTrap™ FF column to remove his tagged LipTev. The flowthrough was collected, and the column was washed with 2ml of gel filtration buffer. These fractions were pooled and concentrated and stored at -80°C.



To purify phosphorylated p110 $\gamma$ , the purification protocol as described above was performed but PKC $\beta$  was added to the strep column at a molar ratio of 1:3 (PKC $\beta$ :p110 $\gamma$ ) along with LipTEV, 20 mM MgCl<sub>2</sub> and 1mM ATP and allowed to incubate on ice for 4 hours. The protein was eluted by adding 7 ml of gel filtration buffer and treated with a second dose of PKC $\beta$  (same ratio as above) and allowed to incubate on ice for another 3 hours. For non-phosphorylated p110 $\gamma$ , same protocol was followed with the exception in the addition of PKC $\beta$ . Both the proteins were concentrated in a 50,000 MWCO Amicon Concentrator (Millipore) to <1 mL and injected onto a Superdex™ 200 10/300 GL Increase size-exclusion column (GE Healthcare) equilibrated in gel filtration buffer. The final phosphorylation level of the two sites was characterised by mass spectrometry, with these values being 92% and 90.8%, for S582 and S594/S595 respectively. After size exclusion, the protein was concentrated, aliquoted, frozen, and stored at -80°C.

## Expression and Purification of lipidated G $\beta\gamma$ for kinase activity assays

Full length, lipidated human G $\beta\gamma$  (G $\beta$ 1 $\gamma$ 2) was expressed in Sf9 insect cells and purified as described previously. After 65 hours of expression, cells were harvested, and the pellets were frozen as described above. Pellets were resuspended in lysis buffer (20 mM HEPES pH 7.7, 100 mM NaCl, 10 mM  $\beta$ ME, protease inhibitor (Protease Inhibitor Cocktail Set III, Sigma)) and sonicated for 2 minutes (15s on, 15s off, level 4.0, Misonix sonicator 3000). The lysate was spun at 500 RCF (Eppendorf Centrifuge 5810 R) to remove intact cells and the supernatant was centrifuged again at 25,000 RCF for 1 hour (Beckman Coulter JA-20 rotor). The pellet was resuspended in lysis buffer and sodium cholate was added to a final concentration of 1% and stirred at 4°C for 1 hour. The membrane extract was clarified by spinning at 10,000 RCF for 30 minutes (Beckman Coulter JA-20 rotor). The supernatant was diluted 3 times with NiNTA A buffer (20 mM HEPES pH 7.7, 100 mM NaCl, 10 mM Imidazole, 0.1% C12E10, 10mM  $\beta$ ME) and loaded onto a 5 mL HisTrap™ FF crude column (GE Healthcare) equilibrated in the same buffer. The column was washed with NiNTA A, 6% NiNTA B buffer (20 mM HEPES pH 7.7, 25 mM NaCl, 250 mM imidazole pH 8.0, 0.1% C12E10, 10 mM  $\beta$ ME) and the protein was eluted with 100% NiNTA B. The eluent was loaded onto HiTrap™ Q HP anion exchange column equilibrated in Hep A buffer (20 mM Tris pH 8.0, 8 mM CHAPS, 2 mM Dithiothreitol (DTT)). A gradient was started with Hep B buffer (20 mM Tris pH 8.0, 500 mM NaCl, 8 mM CHAPS, 2 mM DTT) and the protein was eluted in ~50% Hep B buffer. The eluent was concentrated in a 30,000 MWCO Amicon Concentrator (Millipore) to < 1 mL and injected onto a Superdex™ 75 10/300 GL size exclusion column (GE Healthcare) equilibrated in Gel Filtration buffer (20 mM HEPES pH 7.7, 100 mM NaCl, 10 mM CHAPS, 2 mM TCEP). Fractions containing protein were pooled, concentrated, aliquoted, frozen and stored at -80 °C.

## Expression and purification of nanobody

Nanobody NB7-PIK3CG with a C-terminal 6X His tag was expressed from a pMESy4 vector in the periplasm of WK6 *E. coli*. A 1L culture was grown to OD600 of 0.7 in Terrific Broth containing 0.1% glucose and 2mM MgCl<sub>2</sub> in the presence of 100  $\mu$ g/mL ampicillin and was induced with 0.5 mM isopropyl- $\beta$ -D-thiogalactoside (IPTG). Cells were harvested the following day by centrifuging at 2500 RCF (Eppendorf Centrifuge 5810 R) and the pellet was snap-frozen in liquid nitrogen. The frozen pellet was resuspended in 15 mL of TES buffer containing 200 mM Tris pH 8.0, 0.5mM ethylenediaminetetraacetic acid (EDTA) and 500 mM Sucrose and was mixed for 45 minutes at 4°C. To this mixture, 30 mL of TES buffer diluted four times in water was added and mixed for 45 minutes at 4°C to induce osmotic shock. The lysate was clarified by centrifuging at 14,000 rpm for 15 minutes (Beckman Coulter JA-20

rotor). Imidazole was added to the supernatant to final concentration of 10mM loaded onto a 5 mL HisTrap™ FF crude column (GE Healthcare) equilibrated in NiNTA A buffer (20 mM Tris pH 8.0, 100 mM NaCl, 20 mM imidazole pH 8.0, 5% (v/v) glycerol, 2 mM  $\beta$ -mercaptoethanol ( $\beta$ ME)). The column was washed with high salt NiNTA A buffer (20 mM Tris pH 8.0, 1 M NaCl, 20 mM imidazole pH 8.0, 5% (v/v) glycerol, 2 mM  $\beta$ ME), followed by 100% NiNTA A buffer, then a 6% NiNTA B wash buffer (20 mM Tris pH 8.0, 100 mM NaCl, 250 mM imidazole pH 8.0, 5% (v/v) glycerol, 2 mM  $\beta$ ME) and the protein was eluted with 100% NiNTA B. The eluent was concentrated in a 10,000 MWCO Amicon Concentrator (Millipore) to <1 mL and injected onto a Superdex™ 75 10/300 GL Increase size-exclusion column (GE Healthcare) equilibrated in gel filtration buffer (20mM Tris pH 8.5, 100 mM NaCl, 50 mM Ammonium Sulfate and 0.5 mM tris(2-carboxyethyl) phosphine (TCEP)). Following size exclusion, the protein was concentrated, frozen and stored at  $-80^{\circ}\text{C}$ .

## Lipid vesicle preparation for kinase activity assays

Lipid vesicles containing 5% brain phosphatidylinositol 4,5-bisphosphate (PIP<sub>2</sub>), and 95% brain phosphatidylserine (PS), were prepared by mixing the lipids solutions in organic solvent. The solvent was evaporated in a stream of argon following which the lipid film was desiccated in a vacuum for 45 minutes. The lipids were resuspended in lipid buffer (20 mM HEPES pH 7.0, 100 mM NaCl and 10 % glycerol) and the solution was vortexed for 5 minutes followed by sonication for 15 minutes. The vesicles were then subjected to ten freeze thaw cycles and extruded 11 times through a 100-nm filter (T&T Scientific: TT-002-0010). The extruded vesicles were sub-aliquoted and stored at  $-80^{\circ}\text{C}$ . Final vesicle concentration was 2 mg/mL.

## Kinase Assays

All kinase assays were done using Transcreener ADP2 Fluorescence Intensity (FI) assays (Bellbrook labs) which measures ADP production. All assays contained ATP at a final concentration of 100  $\mu\text{M}$ , and those with membranes used vesicles containing 5% phosphatidylinositol 4,5-bisphosphate (PI(4,5)P<sub>2</sub>), and 95% phosphatidylserine (PS) at a final concentration of 0.5 mg/mL.

For assays measuring the inhibition by nanobody, 4X kinase (final concentration: 330 nM for p110 $\gamma$ , 300nM for p110 $\gamma$ /p84 and 12nM for p110 $\gamma$ /p101) was mixed with varying 4X concentrations of nanobody (final concentration: 2 $\mu\text{M}$  – 2.7nM) or kinase buffer (20mM HEPES pH 7.5, 100mM NaCl, 3mM MgCl<sub>2</sub>, 0.03% CHAPS, 2mM TCEP, and 1mM EGTA) and allowed to sit on ice for 15 minutes. 2  $\mu\text{L}$  of protein mix was mixed with 2  $\mu\text{L}$  of lipid solution containing G $\beta\gamma$  (1 $\mu\text{M}$  final concentration), ATP (100  $\mu\text{M}$ ), PIP<sub>2</sub> lipid vesicles (0.5mg/ml final concentration), and lipid buffer (25mM HEPES pH 7, 5% Glycerol, and 100mM NaCl) and incubated at  $20^{\circ}\text{C}$  for 60 minutes.

For assays comparing the difference in activation between phosphorylated and non-phosphorylated p110 $\gamma$ , 2X kinase (final concentrations: 1 $\mu\text{M}$ ) was mixed with 2X lipid solutions containing ATP (100  $\mu\text{M}$ ), and lipid buffer and either nanobody (3  $\mu\text{M}$  final concentration), PIP<sub>2</sub> lipid vesicles (0.5mg/ml final concentration) or both nanobody and lipid. The reaction was incubated at  $20^{\circ}\text{C}$  for 60 minutes.

After the 60-minute incubation, all reactions were stopped with 4  $\mu\text{L}$  of 2X stop and detect solution containing Stop and Detect buffer (20mM HEPES, 0.02% Brij-35, 400mM 40mM EDTA pH 7.5), 8 nM ADP Alexa Fluor 594 Tracer and 93.7  $\mu\text{g/mL}$  ADP2 Antibody IRDye QC-1, covered and incubated at  $20^{\circ}\text{C}$  for 1 hr before reading the fluorescence. The fluorescence intensity was measured using a SpectraMax M5 plate reader at excitation 590 nm and emission 620 nm. All data was normalized against the appropriate measurements obtained

for 100  $\mu\text{M}$  ATP and 100  $\mu\text{M}$  ADP with no kinase. The percent ATP turnover was interpolated using a standard curve (0.1–100  $\mu\text{M}$  ADP). Interpolated values were then used to calculate the specific activity of the enzyme.

## Biolayer interferometry

All Biolayer interferometry experiments were performed using the Octet K2 instrument (Fortebio Inc.). For all experiments His-tagged nanobody (500 nM) was immobilized on an Anti-Penta-His biosensor for 600s, and the sensor was dipped into varying concentrations of the protein complex being measured. A dose response was carried out for p110 $\gamma$ , p110 $\gamma$ -p84, and p110 $\gamma$ -p101 (50 nM – 1.9 nM), with association occurring for 600s, followed by a 1200s dissociation in Octet Buffer (20 mM tris pH 8.5, 100 mM NaCl, 50 mM ammonium sulfate, 0.1% bovine serum albumin, and 0.02% Tween 20). Experiments comparing class IA PI3K versus class IB PI3K used 50 nM of each class IA PI3K.

When comparing nanobody binding to phosphorylated and unphosphorylated p110 $\gamma$ , we used a final concentration of 25 nM for both phosphorylated and non-phosphorylated p110 $\gamma$  with association occurring for 600s, followed by a 600s dissociation. The  $K_D$  (dissociation constant) for the different p110 $\gamma$  complexes was calculated from the binding curves based on their global fit to a 1:1 binding model using ForteBio data analysis 12.0 (Fortebio Inc.).

## Supported lipid bilayer TIRF microscopy experiments

The membrane binding dynamics of Dy647-p84-p110 $\gamma$  were measured in the absence and presence of nanobody 7 (NB7) using TIRF microscopy. As previously described (Rathinaswamy et al., 2023), supported lipid bilayers were formed using 50 nm extruded small unilamellar vesicles (SUVs) containing the following lipids: 1,2-dioleoyl-sn-glycero-3-phosphocholine (18:1 DOPC, Avanti # 850375C), 1,2-dioleoyl-sn-glycero-3-phospho-L-serine (18:1 DOPS, Avanti # 840035C), 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-[4-(p-maleimidomethyl)cyclohexane-carboxamide] (18:1 MCC-PE, Avanti # 780201C). Lipid compositions reported in figure legends represent the molar percentage of each lipid species.

To create SLBs, a total concentration of 0.25 mM lipids was solvated in 1x PBS [pH 7.4] and deposited on Piranha etched glass coverslips (25 x 75 mm) adhered to an IBIDI chamber. After a 30-minute incubation, membranes were washed with 4 mL of 1x PBS [pH 7.4] and then blocked for 10 minutes with 1 mg/mL beta casein (Thermo FisherSci, Cat# 37528) in 1x PBS [pH 7.4] (Corning, Cat# 46-013-CM). To conjugate H-Ras to maleimide lipids (MCC-PE), blocked membranes were incubated with 30  $\mu\text{M}$  H-Ras (GDP) in buffer containing 1x PBS [pH 7.4], 1 mM  $\text{MgCl}_2$ , 50  $\mu\text{M}$  GDP, and 0.1 mM TCEP for 2 hours. The membrane conjugation reaction was terminated after 2 hours with 1x PBS [pH 7.4] containing 5 mM  $\beta$ -mercaptoethanol ( $\beta$ ME). Membranes were then washed and stored in 1x PBS [pH 7.4] until performing the TIRF-M membrane binding experiments. H-Ras was purified as previously described (Rathinaswamy et al., 2023).

To perform the TIRF-M membrane binding assays, 200 nM farnesyl- $G\beta\gamma$  was equilibrated into the supported membranes for 30 minutes. In parallel, nucleotide exchange of H-Ras (GDP) was performed by adding 50 nM guanine nucleotide exchange factor (SosCat) in 1x PBS [pH 7.4], 1 mM  $\text{MgCl}_2$ , 50  $\mu\text{M}$  GDP. To measure membrane binding, Dy647-p84-p110 $\gamma$  was diluted into the following buffer: 20 mM HEPES [pH 7.0], 150 mM NaCl, 50  $\mu\text{M}$  GTP, 1 mM ATP, 5 mM  $\text{MgCl}_2$ , 0.5 mM EGTA, 20 mM glucose, 200  $\mu\text{g/mL}$  beta casein (ThermoScientific, Cat# 37528), 20 mM BME, 320  $\mu\text{g/mL}$  glucose oxidase (Serva, #22780.01 *Aspergillus niger*), 50  $\mu\text{g/mL}$  catalase (Sigma, #C40-100MG Bovine Liver), and 2 mM Trolox. Trolox was prepared as

previously described (Hansen et al., 2019). Perishable reagents (i.e., glucose oxidase, catalase, and Trolox) were added 10 minutes before image acquisition.

TIRF-M experiments were performed using an inverted Nikon Ti2 microscope with a 100x Nikon (1.49 NA) oil immersion objective. The x-axis and y-axis positions were controlled using a Nikon motorized stage. Dy647-p84-p110y was excited with a 637 nm diode laser (OBIS laser diode, Coherent Inc. Santa Clara, CA) controlled with an acousto-optic tunable filter (AOTF) and laser launch built by Vortran (Sacramento, CA). The power output measured through the objective for single particle imaging was 1-3 mW. Excitation light passing through quad multi-pass dichroic filter cube (Semrock). Fluorescence emission passed through Nikon emission filter wheel containing the following 25 mm ET700/75M emission filters (Semrock) before being detected on iXion Life 897 EMCCD camera (Andor Technology Ltd., UK). All TIRF-M experiments were performed at room temperature (23°C). Microscope hardware was controlled using Nikon NIS elements. Data analysis was performed using ImageJ/Fiji and Prism graphing program.

## Cryo-EM Sample Preparation and Data Collection

3  $\mu$ L of purified nanobody-bound p110y at 0.45 mg/ml was adsorbed onto C-Flat 2/2-T grids that were glow discharged for 25 s at 15 mA. Grids were then plunged into liquid ethane using a Vitrobot Mark IV (Thermo Fisher Scientific) with the following settings: -5 blot force, 1.5 s blot time, 100% humidity and 4 °C. Vitrified specimens were screened for ice and particle quality at the UBC High resolution macromolecular electron microscopy (HRMEM) facility using a 200-kV Glacios transmission electron microscope equipped with a Falcon 3EC direct electron detector (DED). Clipped grids were sent to the Pacific Northwest Cryo-EM Center (PNCC) where 7,322 movies were collected using a Titan Krios equipped with a Gatan K3 DED and a BioQuantum K3 energy filter with a slit width of 20 eV. The movies were collected at a physical pixel size of 0.830 Å/pix and a total dose of 50e<sup>-</sup>/Å<sup>2</sup> over 50 frames.

## Cryo-EM image analysis

The data were processed using cryoSPARC v.3.3.2 (Punjani et al., 2017). The movies were pre-processed by patch motion correction using default settings except Fourier-cropping by a factor of 2, followed by patch CTF estimation using default settings. A 3D map of PI3K p110y-p101 complex (EMD-23808) was used to create 2D projections for use as templates to auto-pick 1,463,553 particles. Particles were extracted with a box size of 380 pixels, Fourier cropped to a box size of 96 pixels and subjected to 2D classification. After discarding classes with obvious noise and no features, 795,162 particles were used for multiple rounds of *ab initio* reconstruction and heterogeneous refinement using 4 or 5 classes. 365,178 particles, which generated the two best 3D reconstruction, were used to carry out Per-particle local-motion correction with 760 pixels box size later downsized to 380 pixels followed by several rounds of *ab initio* reconstruction and heterogeneous refinement using 3 or 5 classes. 149,603 from best class were further refined by homogeneous refinement and a final Non-Uniform (NU)-refinement which generated a reconstruction with an overall resolution of 3.02 Å based on the Fourier shell correlation (FSC) 0.143 criterion.

## Building the structural model of p110y-NB7

The previous structural model of full length p110y from the complex of p110y-p101(PDB: 7MEZ) (Rathinaswamy et al., 2021a) was fit into the map using Chimera (Pettersen et al., 2004). A model of the nanobody was generated using AlphaFold2 using the Colabfold v1.5.2 server (Mirdita et al., 2022). The CDR loops were removed from this initial model, and the remaining nanobody was fit into the map using Chimera. The final structure was built by



iterative rounds of automated model building in Phenix, manual model building in COOT (Emsley et al., 2010), and refinement in Phenix.real\_space\_refine using realspace, rigid body, and adp refinement with tight secondary structure restraints (Afonine et al., 2012). This allowed for unambiguous building of the CDRs of the nanobody, and their interface with p110 $\gamma$ . The full refinement and validation statistics are shown in Supplemental table 1.

## Phosphorylation analysis

For the dose–response phosphorylation of p110 $\gamma$ , p110/p84, and p110/p101, each protein or complex (750nM) was mixed with ATP (200  $\mu$ M), GFB (20mM Tris pH 8.5, 100 mM NaCl, 50 mM Ammonium Sulfate and 0.5 mM TCEP), MgCl<sub>2</sub> (20mM) and various amounts of PKC $\beta$  (4  $\mu$ g, 800 ng, 160 ng, 32 ng, 6.4 ng, and 0 ng). Reactions were incubated for three hours on ice and quenched with 50  $\mu$ L of ice-cold acidic quench buffer (0.7 M guanidine-HCl, 1% formic acid), followed by immediate freezing using liquid nitrogen and storage at  $-80^{\circ}\text{C}$ .

For the experiment studying the effect of nanobody on phosphorylation, p110 $\gamma$  or p110/p101, (500nM) was mixed with ATP (1 mM), GFB (20mM Tris pH 8.5, 100 mM NaCl, 50 mM Ammonium Sulfate and 0.5 mM TCEP), MgCl<sub>2</sub>(20mM), with nanobody and PKC $\beta$  present at 1200 nM and 500nM, respectively. Reactions were incubated for one hour at room temperature and quenched with 54  $\mu$ L of ice-cold acidic quench buffer (0.7 M guanidine-HCl, 1% formic acid) followed by immediate freezing using liquid nitrogen and storage at  $-80^{\circ}\text{C}$ .

Phosphorylation of all proteins was confirmed using Mass spectrometry and PEAKS7 analysis. The LC-MS analysis of these samples was carried out using the same pipeline as used in the HDX-MS section. The phosphorylated and non-phosphorylated peptide ratios were determined by generating extracted ion chromatograms for each phosphorylated or non-phosphorylated peptide using their molecular formula and charge state in the Bruker Compass Data Analysis software. The area under each extracted curve was then extracted. The full MS quantification of each of the phosphorylated and non-phosphorylated peptide is provided in the source data.

## Hydrogen Deuterium eXchange Mass Spectrometry

Exchange reactions to assess differences in p110 $\gamma$  upon phosphorylation were carried out at  $20^{\circ}\text{C}$  in 10  $\mu$ L volumes with final concentrations of 1.6  $\mu$ M for both apo and phosphorylated p110 $\gamma$ . A total of two conditions were assessed: p110 $\gamma$  apo and PKC $\beta$  phosphorylated p110 $\gamma$ . The hydrogen-deuterium exchange reaction was initiated by the addition of 8  $\mu$ L D<sub>2</sub>O buffer (94.3% D<sub>2</sub>O, 100 mM NaCl, 20 mM HEPES pH 7.5) to the 2  $\mu$ L protein for a final D<sub>2</sub>O concentration of 75.4%. Exchange was carried out over five time points (3s on ice, and 3s, 30s, 300s and 3000s at  $20^{\circ}\text{C}$ ) and the reaction was quenched with addition of 60  $\mu$ L of ice-cold acidic quench buffer (0.7 M guanidine-HCl, 1% formic acid). After quenching, samples were immediately frozen in liquid nitrogen and stored at  $-80^{\circ}\text{C}$ . All reactions were carried out in triplicate.

## Protein Digestion and MS/MS Data Collection

Protein samples for both HDX-MS and phosphorylation analysis were analyzed using the same LC-MS setup. Samples were rapidly thawed and injected onto an integrated fluidics system containing a HDx-3 PAL liquid handling robot and climate-controlled chromatography system (LEAP Technologies), a Dionex Ultimate 3000 UHPLC system, as well as an Impact HD QTOF Mass spectrometer (Bruker). The protein was run over two immobilized pepsin columns (Applied Biosystems; Poroszyme™ Immobilized Pepsin Cartridge, 2.1 mm x 30 mm; Thermo-Fisher 2L3131L00; at  $10^{\circ}\text{C}$  and  $2^{\circ}\text{C}$  respectively) at 200

$\mu\text{L}/\text{min}$  for 3 minutes. The resulting peptides were collected and desalted on a C18 trap column [Acquity UPLC BEH C18 1.7 mm column (2.1 x 5 mm); Waters 186003975]. The trap was subsequently eluted in line with an ACQUITY 1.7  $\mu\text{m}$  particle, 100 x 1 mm<sup>2</sup> C18 UPLC column (Waters 186002352), using a gradient of 3-35% B (buffer A, 0.1% formic acid; buffer B, 100% acetonitrile) over 11 min immediately followed by a gradient of 35-80% B over 5 minutes. MS experiments acquired over a mass range from 150 to 2200 mass/charge ratio ( $m/z$ ) using an electrospray ionization source operated at a temperature of 200°C and a spray voltage of 4.5 kV.

## Peptide Identification

Peptides were identified using data-dependent acquisition following tandem MS/MS experiments (0.5 s precursor scan from 150L2000  $m/z$ ; twelve 0.25 s fragment scans from 150-2000  $m/z$ ). MS/MS datasets were analyzed using PEAKS7 (PEAKS), and a false discovery rate was set at 0.1% using a database of purified proteins and known contaminants. The same approach was used to identify phosphorylated and non-phosphorylated peptides for our in-vitro phosphorylation experiments, with variable phosphorylation of STY residues was added to the search. The search parameters were set with a precursor tolerance of 20 parts per million, fragment mass error 0.02 Da, and charge states from 1 to 8, with a selection criterion of peptides that had a  $-10\log P$  score of  $>24.03$  for phosphorylated and  $>23.05$  for non-phosphorylated. The MS/MS spectra of the PKC $\beta$  phosphorylated peptides are included in Fig S4.

## Mass Analysis of Peptide Centroids and Measurement of Deuterium Incorporation

HDExaminer Software (Sierra Analytics) was used to automatically calculate the level of deuterium incorporation into each peptide. All peptides were manually inspected for correct charge state, correct retention time, and appropriate selection of isotopic distribution. Deuteration levels were calculated using the centroid of the experimental isotope clusters. HDX-MS results are presented with no correction for back exchange shown in the Source data, with the only correction being applied correcting for the deuterium oxide percentage of the buffer used in the exchange (75.4%). Changes in any peptide at any time point greater than specified cut-offs (5% and 0.45 Da) and with an unpaired, two-tailed t-test value of  $p < 0.01$  was considered significant. A number of peptides in the helical domain showed isotope distributions consistent with EX1 H/D exchange. Attempts to define the relative percentages of each population using HDExaminer were extremely noisy, so representative EX1 profiles are shown in Fig. 4C. The raw peptide deuterium incorporation graphs for a selection of peptides with significant differences are shown in Fig. 4D, with the raw data for all analysed peptides in the source data. To allow for visualization of differences across all peptides, we utilized number of deuterium difference (#D) plots (Fig. 4B). These plots show the total difference in deuterium incorporation over the entire H/D exchange time course, with each point indicating a single peptide. The data analysis statistics for all HDX-MS experiments are in Supplemental Table 2 according to the guidelines of (Masson et al., 2019). The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository (Perez-Riverol et al., 2022) with the dataset identifier PXD040765.

## References

- Afonine PV , Grosse-Kunstleve RW , Echols N , Headd JJ , Moriarty NW , Mustyakimov M , Terwilliger TC , Urzhumtsev A , Zwart PH , Adams PD (2012) **Towards automated crystallographic structure refinement with phenix.refine** *Acta Crystallogr D Biol Crystallogr* **68**:352–367  
<https://doi.org/10.1107/S0907444912001308>
- Angulo I , Vadas O , Garçon F , Banham-Hall E , Plagnol V , Leahy TR , Baxendale H , Coulter T , Curtis J , Wu C , Blake-Palmer K , Perisic O , Smyth D , Maes M , Fiddler C , Juss J , Cilliers D , Markelj G , Chandra A , Farmer G , Kielkowska A , Clark J , Kracker S , Debré M , Picard C , Pellier I , Jabado N , Morris JA , Barcenas-Morales G , Fischer A , Stephens L , Hawkins P , Barrett JC , Abinun M , Clatworthy M , Durandy A , Doffinger R , Chilvers ER , Cant AJ , Kumararatne D , Okkenhaug K , Williams RL , Condliffe A , Nejentsev S (2013) **Phosphoinositide 3-kinase  $\delta$  gene mutation predisposes to respiratory infection and airway damage** *Science* **342**:866–871  
<https://doi.org/10.1126/science.1243292>
- Becattini B , Marone R , Zani F , Arsenijevic D , Seydoux J , Montani J , Dulloo A , Thorens B , Preitner F , Wymann M , Solinas G (2011) **PI3Kgamma within a nonhematopoietic cell type negatively regulates diet-induced thermogenesis and promotes obesity and insulin resistance** *Proc Natl Acad Sci USA* **108**  
<https://doi.org/10.1073/pnas.1106698108/-/DCSupplemental>
- Bell K , Sunose M , Ellard K , Cansfield A , Taylor J , Miller W , Ramsden N , Bergamini G , Neubauer G (2012) **SAR studies around a series of triazolopyridines as potent and selective PI3Ky inhibitors** *Bioorg Med Chem Lett* **22**:5257–5263  
<https://doi.org/10.1016/j.bmcl.2012.06.049>
- Bohnacker T , Marone R , Collmann E , Calvez R , Hirsch E , Wymann M (2009) **PI3Kgamma adaptor subunits define coupling to degranulation and cell motility by distinct PtdIns(3,4,5)P3 pools in mast cells** *Sci Signal* **2**  
<https://doi.org/10.1126/scisignal.2000259>
- Bohnacker T , Prota AE , Beaufils F , Burke JE , Melone A , Inglis AJ , Rageot D , Sele AM , Cmiljanovic V , Cmiljanovic N , Bargsten K , Aher A , Akhmanova A , Díaz JF , Fabbro D , Zvelebil M , Williams RL (2017) **Steinmetz MO Wymann MP**
- Deconvolution of Buparlisib's mechanism of action defines specific PI3K and tubulin inhibitors for therapeutic intervention (1970) **Deconvolution of Buparlisib's mechanism of action defines specific PI3K and tubulin inhibitors for therapeutic intervention. Nat Commun** **8**:14683. doi:10.1038/ncomms14683 *Nat Commun* **8**  
<https://doi.org/10.1038/ncomms14683>
- Breasson L , Becattini B , Sardi C , Molinaro A , Zani F , Marone R , Botindari F , Bousquenaud M , Ruegg C , Wymann MP , Solinas G (2017) **PI3Ky activity in leukocytes promotes adipose tissue inflammation and early-onset insulin resistance during obesity** *Sci Signal* **10**  
<https://doi.org/10.1126/scisignal.aaf2969>

Burke JE (2018) **Structural Basis for Regulation of Phosphoinositide Kinases and Their Involvement in Human Disease** *Mol Cell* **71**:653–673  
<https://doi.org/10.1016/j.molcel.2018.08.005>

Burke JE , Perisic O , Masson GR , Vadas O , Williams RL (2012) **Oncogenic mutations mimic and enhance dynamic events in the natural activation of phosphoinositide 3-kinase p110α (PIK3CA)** *Proc Natl Acad Sci USA* **109**:15259–15264  
<https://doi.org/10.1073/pnas.1205508109>

Burke JE , Vadas O , Berndt A , Finegan T , Perisic O , Williams RL (2011) **Dynamics of the phosphoinositide 3-kinase p110δ interaction with p85α and membranes reveals aspects of regulation distinct from p110α** *Structure* **19**:1127–1137  
<https://doi.org/10.1016/j.str.2011.06.003>

Burke JE , Williams RL (2015) **Synergy in activating class I PI3Ks** *Trends in Biochemical Sciences* **40**:88–100  
<https://doi.org/10.1016/j.tibs.2014.12.003>

Burke JE , Williams RL (2013) **Dynamic steps in receptor tyrosine kinase mediated activation of class IA phosphoinositide 3-kinases (PI3K) captured by H/D exchange (HDX-MS)** *Adv Biol Regul* **53**:97–110  
<https://doi.org/10.1016/j.jbior.2012.09.005>

Campa CC , Silva RL , Margaria JP , Pirali T , Mattos MS , Kraemer LR , Reis DC , Grosa G , Copperi F , Dalmarco EM , Lima-Júnior RCP , Aprile S , Sala V , Dal Bello F , Prado DS , Alves-Filho JC , Medana C , Cassali GD , Tron GC , Teixeira MM , Ciruolo E , Russo RC , Hirsch E (2018) **Inhalation of the prodrug PI3K inhibitor CL27c improves lung function in asthma and fibrosis** *Nat Commun* **9**:5232–16  
<https://doi.org/10.1038/s41467-018-07698-6>

Camps M , Rückle T , Ji H , Ardisson V , Rintelen F , Shaw J , Ferrandi C , Chabert C , Gillieron C , Françon B , Martin T , Gretener D , Perrin D , Leroy D , Vitte P-A , Hirsch E , Wymann MP , Cirillo R , Schwarz MK , Rommel C (2005) **Blockade of PI3Kγ suppresses joint inflammation and damage in mouse models of rheumatoid arthritis** *Nat Med* **11**:936–943  
<https://doi.org/10.1038/nm1284>

De Henau O , Rausch M , Winkler D , Campesato LF , Liu C , Cymerman DH , Budhu S , Ghosh A , Pink M , Tchaicha J , Douglas M , Tibbitts T , Sharma S , Proctor J , Kosmider N , White K , Stern H , Soglia J , Adams J , Palombella VJ , McGovern K , Kutok JL , Wolchok JD , Merghoub T. (2016) **Overcoming resistance to checkpoint blockade therapy by targeting PI3Kγ in myeloid cells** *Nature* **539**:443–447  
<https://doi.org/10.1038/nature20554>

Deladeriere A , Gambardella L , Pan D , Anderson KE , Hawkins PT , Stephens LR (2015) **The regulatory subunits of PI3Kγ control distinct neutrophil responses** *Sci Signal* **8**  
<https://doi.org/10.1126/scisignal.2005564>



Dornan GL , Siempelkamp BD , Jenkins ML , Vadas O , Lucas CL , Burke JE (2017) **Conformational disruption of PI3K $\delta$  regulation by immunodeficiency mutations in PI3KCD and PIK3R1** *Proc Natl Acad Sci USA* **114**:1982–1987  
<https://doi.org/10.1073/pnas.1617244114>

Emsley P , Lohkamp B , Scott WG , Cowtan K (2010) **Features and development of Coot** *Acta Crystallogr D Biol Crystallogr* **66**:486–501  
<https://doi.org/10.1107/S0907444910007493>

Evans CA , Liu T , Lescarbeau A , Nair SJ , Grenier L , Pradeilles JA , Glenadel Q , Tibbitts T , Rowley AM , DiNitto JP , Brophy EE , O’Hearn EL , Ali JA , Winkler DG , Goldstein SI , O’Hearn P , Martin CM , Hoyt JG , Soglia JR , Cheung C , Pink MM , Proctor JL , Palombella VJ , Tremblay MR , Castro AC (2016) **Discovery of a Selective Phosphoinositide-3-Kinase (PI3K)- $\gamma$  Inhibitor (IPI-549) as an Immuno-Oncology Clinical Candidate** *ACS Med Chem Lett* **7**:862–867  
<https://doi.org/10.1021/acsmedchemlett.6b00238>

Gangadhara G , Dahl G , Bohnacker T , Rae R , Gunnarsson J , Blaho S , Öster L , Lindmark H , Karabelas K , Pemberton N , Tyrchan C , Mogemark M , Wymann MP , Williams RL , Perry MWD , Papavoine T , Petersen J (2019) **A class of highly selective inhibitors bind to an active state of PI3K $\gamma$**  *Nature Chemical Biology* **15**:348–357  
<https://doi.org/10.1038/s41589-018-0215-0>

Hansen SD , Huang WYC , Lee YK , Bieling P , Christensen SM , Groves JT (2019) **Stochastic geometry sensing and polarization in a lipid kinase-phosphatase competitive reaction** *Proc Natl Acad Sci U S A* **116**:15013–15022  
<https://doi.org/10.1073/pnas.1901744116>

Hawkins PT , Stephens LR (2015) **PI3K signalling in inflammation** *Biochim Biophys Acta* **1851**:882–897  
<https://doi.org/10.1016/j.bbailip.2014.12.006>

Jenkins ML , Ranga-Prasad H , Parson MAH , Harris NJ , Rathinaswamy MK , Burke JE (2023) **Oncogenic mutations of PIK3CA lead to increased membrane recruitment driven by reorientation of the ABD, p85 and C-terminus** *Nat Commun* **14**  
<https://doi.org/10.1038/s41467-023-35789-6>

Jin JR , Gogvadze E , Xavier AR , Bohnacker T , Voelzmann J , Wymann MP (2020) **PI3K $\gamma$  Regulatory Protein p84 Determines Mast Cell Sensitivity to Ras Inhibition-Moving Towards Cell Specific PI3K Targeting?** *Front Immunol* **11**  
<https://doi.org/10.3389/fimmu.2020.585070>

Johannessen CM , Boehm JS , Kim SY , Thomas SR , Wardwell L , Johnson LA , Emery CM , Stransky N , Cogdill AP , Barretina J , Caponigro G , Hieronymus H , Murray RR , Salehi-Ashtiani K , Hill DE , Vidal M , Zhao JJ , Yang X , Alkan O , Kim S , Harris JL , Wilson CJ , Myer VE , Finan PM , Root DE , Roberts TM , Golub T , Flaherty KT , Dummer R , Weber BL , Sellers WR , Schlegel R , Wargo JA , Hahn WC , Garraway LA (2010) **COT drives resistance to RAF inhibition through MAP kinase pathway reactivation** *Nature* **468**:968–972  
<https://doi.org/10.1038/nature09627>

Kaneda MM , Cappello P , Nguyen AV , Ralainirina N , Hardamon CR , Foubert P , Schmid MC , Sun P , Mose E , Bouvet M , Lowy AM , Valasek MA , Sasik R , Novelli F , Hirsch E , Varner JA (2016) **Macrophage PI3Ky Drives Pancreatic Ductal Adenocarcinoma Progression** *Cancer Discov* **6**:870–885

<https://doi.org/10.1158/2159-8290.CD-15-1346>

Kaneda MM , Messer KS , Ralainirina N , Li H , Leem CJ , Gorjestani S , Woo G , Nguyen AV , Figueiredo CC , Foubert P , Schmid MC , Pink M , Winkler DG , Rausch M , Palombella VJ , Kutok J , McGovern K , Frazer KA , Wu X , Karin M , Sasik R , Cohen EEW , Varner JA (2016) **PI3Ky is a molecular switch that controls immune suppression** *Nature* **539**:437–442

<https://doi.org/10.1038/nature19834>

Kurig B , Shymanets A , Bohnacker T , Prajwal Brock C , Ahmadian MR , Schaefer M , Gohla A , Harteneck C , Wymann MP , Jeanclos E , Nürnberg B (2009) **Ras is an indispensable coregulator of the class IB phosphoinositide 3-kinase p87/p110gamma** *Proc Natl Acad Sci USA* **106**:20312–20317

<https://doi.org/10.1073/pnas.0905506106>

Laffargue M , Calvez R , Finan P , Trifilieff A , Barbier M , Altruda F , Hirsch E , Wymann MP (2002) **Phosphoinositide 3-kinase gamma is an essential amplifier of mast cell function** *Immunity* **16**:441–451

Lanahan SM , Wymann MP , Lucas CL (2022) **The role of PI3Ky in the immune system: new insights and translational implications** *Nat Rev Immunol*

<https://doi.org/10.1038/s41577-022-00701-8>

Lawrence MS , Stojanov P , Mermel CH , Robinson JT , Garraway LA , Golub TR , Meyerson M , Gabriel SB , Lander ES , Getz G (2014) **Discovery and saturation analysis of cancer genes across 21 tumour types** *Nature* **505**:495–501

<https://doi.org/10.1038/nature12912>

Li H , Prever L , Hirsch E , Gulluni F (2021) **Targeting PI3K/AKT/mTOR Signaling Pathway in Breast Cancer** *Cancers (Basel)* **13**

<https://doi.org/10.3390/cancers13143517>

Li Z , Jiang H , Xie W , Zhang Z , Smrcka AV , Wu D (2000) **Roles of PLC-beta2 and -beta3 and PI3Kgamma in chemoattractant-mediated signal transduction** *Science* **287**:1046–1049

<https://doi.org/10.1126/science.287.5455.1046>

Liu X , Zhou Q , Hart JR , Xu Y , Yang S , Yang D , Vogt PK , Wang M-W (2022) **Cryo-EM structures of cancer-specific helical and kinase domain mutations of PI3Ka** *Proc Natl Acad Sci U S A* **119**

<https://doi.org/10.1073/pnas.2215621119>

Lucas CL , Kuehn HS , Zhao F , Niemela JE , Deenick EK , Palendira U , Avery DT , Moens L , Cannons JL , Biancalana M , Stoddard J , Ouyang W , Frucht DM , Rao VK , Atkinson TP , Agharahimi A , Hussey AA , Folio LR , Olivier KN , Fleisher TA , Pittaluga S , Holland SM , Cohen JI , Oliveira JB , Tangye SG , Schwartzberg PL , Lenardo MJ , Uzel G (2014) **Dominant-activating germline mutations in the gene encoding the PI(3)K catalytic subunit p110δ result in T cell senescence and human immunodeficiency** *Nat Immunol* **15**:88–97

<https://doi.org/10.1038/ni.2771>

Luo L , Wall AA , Tong SJ , Hung Y , Xiao Z , Tarique AA , Sly PD , Fantino E , Marzolo M-P , Stow JL (2018) **TLR Crosstalk Activates LRP1 to Recruit Rab8a and PI3Ky for Suppression of Inflammatory Responses** *Cell Rep* **24**:3033–3044

<https://doi.org/10.1016/j.celrep.2018.08.028>

Mandelker D , Gabelli SB , Schmidt-Kittler O , Zhu J , Cheong I , Huang C-H , Kinzler KW , Vogelstein B , Amzel LM (2009) **A frequent kinase domain mutation that changes the interaction between PI3Kalpha and the membrane** *Proc Natl Acad Sci USA* **106**:16996–17001

<https://doi.org/10.1073/pnas.0908444106>

Masson GR , Burke JE , Ahn NG , Anand GS , Borchers C , Brier S , Bou-Assaf GM , Engen JR , Englander SW , Faber J , Garlish R , Griffin PR , Gross ML , Guttman M , Hamuro Y , Heck AJR , Houde D , Jacob RE , Jørgensen TJD , Kaltashov IA , Klinman JP , Konermann L , Man P , Mayne L , Pascal BD , Reichmann D , Skehel M , Snijder J , Strutzenberg TS , Underbakke ES , Wagner C , Wales TE , Walters BT , Weis DD , Wilson DJ , Wintrobe PL , Zhang Z , Zheng J , Schriemer DC , Rand KD (2019) **Recommendations for performing, interpreting and reporting hydrogen deuterium exchange mass spectrometry (HDX-MS) experiments** *Nat Methods* **16**:595–602

<https://doi.org/10.1038/s41592-019-0459-y>

Miled N , Yan Y , Hon W-C , Perisic O , Zvelebil M , Inbar Y , Schneidman-Duhovny D , Wolfson HJ , Backer JM , Williams RL (2007) **Mechanism of two classes of cancer mutations in the phosphoinositide 3-kinase catalytic subunit** *Science* **317**:239–242

<https://doi.org/10.1126/science.1135394>

Mirdita M , Schütze K , Moriwaki Y , Heo L , Ovchinnikov S , Steinegger M (2022) **ColabFold: making protein folding accessible to all** *Nat Methods* **19**:679–682

<https://doi.org/10.1038/s41592-022-01488-1>

Okkenhaug K (2013) **Signaling by the phosphoinositide 3-kinase family in immune cells** *Annu Rev Immunol* **31**:675–704

<https://doi.org/10.1146/annurev-immunol-032712-095946>

Pacold ME , Suire S , Perisic O , Lara-Gonzalez S , Davis CT , Walker EH , Hawkins PT , Stephens L , Eccleston JF , Williams RL (2000) **Crystal structure and functional analysis of Ras binding to its effector phosphoinositide 3-kinase gamma** *Cell* **103**:931–943

Perez-Riverol Y , Bai J , Bandla C , García-Seisdedos D , Hewapathirana S , Kamatchinathan S , Kundu DJ , Prakash A , Frericks-Zipper A , Eisenacher M , Walzer M , Wang S , Brazma A , Vizcaíno JA (2022) **The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences** *Nucleic Acids Res* **50**

<https://doi.org/10.1093/nar/gkab1038>

Perino A , Ghigo A , Ferrero E , Morello F , Santulli G , Baillie GS , Damilano F , Dunlop AJ , Pawson C , Walser R , Levi R , Altruda F , Silengo L , Langeberg LK , Neubauer G , Heymans S , Lembo G , Wymann MP , Wetzker R , Houslay MD , Iaccarino G , Scott JD , Hirsch E (2011) **Integrating Cardiac PIP(3) and cAMP Signaling through a PKA Anchoring Function of p110gamma** *Mol Cell* **42**:84–95  
<https://doi.org/10.1016/j.molcel.2011.01.030>

Pettersen EF , Goddard TD , Huang CC , Couch GS , Greenblatt DM , Meng EC , Ferrin TE (2004) **UCSF Chimera--a visualization system for exploratory research and analysis** *J Comput Chem* **25**:1605–1612  
<https://doi.org/10.1002/jcc.20084>

Punjani A , Rubinstein JL , Fleet DJ , Brubaker MA (2017) **cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination** *Nat Methods* **14**:290–296  
<https://doi.org/10.1038/nmeth.4169>

Rathinaswamy MK , Burke JE (2019) **Class I phosphoinositide 3-kinase (PI3K) regulatory subunits and their roles in signaling and disease** *Adv Biol Regul* **100**:657  
<https://doi.org/10.1016/j.jbior.2019.100657>

Rathinaswamy MK , Dalwadi U , Fleming KD , Adams C , Stariha JTB , Pardon E , Baek M , Vadas O , DiMaio F , Steyaert J , Hansen SD , Yip CK , Burke JE (2021) **Structure of the phosphoinositide 3-kinase (PI3K) p110γ-p101 complex reveals molecular mechanism of GPCR activation** *Sci Adv* **7**  
<https://doi.org/10.1126/sciadv.abj4282>

Rathinaswamy MK , Fleming KD , Dalwadi U , Pardon E , Harris NJ , Yip CK , Steyaert J , Burke JE (2021) **HDX-MS-optimized approach to characterize nanobodies as tools for biochemical and structural studies of class IB phosphoinositide 3-kinases** *Structure* **29**:1371–1381  
<https://doi.org/10.1016/j.str.2021.07.002>

Rathinaswamy MK , Gaieb Z , Fleming KD , Borsari C , Harris NJ , Moeller BE , Wymann MP , Amaro RE , Burke JE (2021) **Disease-related mutations in PI3Ky disrupt regulatory C-terminal dynamics and reveal a path to selective inhibitors** *Elife* **10**  
<https://doi.org/10.7554/eLife.64691>

Rathinaswamy MK , Jenkins ML , Duwell BR , Zhang X , Harris NJ , Evans JT , Stariha JTB , Dalwadi U , Fleming KD , Ranga-Prasad H , Yip CK , Williams RL , Hansen SD , Burke JE (2023) **Molecular basis for differential activation of p101 and p84 complexes of PI3Ky by Ras and GPCRs** *Cell Rep* **42**  
<https://doi.org/10.1016/j.celrep.2023.112172>

Rynkiewicz NK , Anderson KE , Suire S , Collins DM , Karanasios E , Vadas O , Williams R , Oxley D , Clark J , Stephens LR , Hawkins PT (2020) **Gβγ is a direct regulator of endogenous p101/p110γ and p84/p110γ PI3Ky complexes in mouse neutrophils** *Sci Signal* **13**  
<https://doi.org/10.1126/scisignal.aaz4003>



Samuels Y, Wang Z, Bardelli A, Silliman N, Ptak J, Szabo S, Yan H, Gazdar A, Powell S, Riggins G, Willson J, Markowitz S, Kinzler K, Vogelstein B, Velculescu V (2004) **High frequency of mutations of the PIK3CA gene in human cancers** *Science* **304**  
<https://doi.org/10.1126/science.1096502>

Shymanets A, Prajwal P, Bucher K, Beer-Hammer S, Harteneck C, Nürnberg B (2013) **p87 and p101 subunits are distinct regulators determining class IB PI3K specificity** *J Biol Chem* **288**:31059–68  
<https://doi.org/10.1074/jbc.M113.508234>

Siempelkamp BD, Rathinaswamy MK, Jenkins ML, Burke JE (2017) **Molecular mechanism of activation of class IA phosphoinositide 3-kinases (PI3Ks) by membrane-localized HRas** *J Biol Chem* **292**:12256–12266  
<https://doi.org/10.1074/jbc.M117.789263>

Stephens LR, Eguinoa A, Erdjument-Bromage H, Lui M, Cooke F, Coadwell J, Smrcka AS, Thelen M, Cadwallader K, Tempst P, Hawkins PT (1997) **The Gbg sensitivity of a PI3K is dependent upon a tightly associated adaptor, p101** *Cell* **89**:105–114

Takeda AJ, Maher TJ, Zhang Y, Lanahan SM, Bucklin ML, Compton SR, Tyler PM, Comrie WA, Matsuda M, Olivier KN, Pittaluga S, McElwee JJ, Long Priel DA, Kuhns DB, Williams RL, Mustillo PJ, Wymann MP, Koneti Rao V, Lucas CL (2019) **Human PI3Ky deficiency and its microbiota-dependent mouse model reveal immunodeficiency and tissue immunopathology** *Nat Commun* **10**:4364–12  
<https://doi.org/10.1038/s41467-019-12311-5>

Vadas O, Dbouk HA, Shymanets A, Perisic O, Burke JE, Abi Saab WF, Khalil BD, Harteneck C, Bresnick AR, Nürnberg B, Backer JM, Williams RL (2013) **Molecular determinants of PI3Ky-mediated activation downstream of G-protein-coupled receptors (GPCRs)** *Proc Natl Acad Sci U S A* **110**:18862–18867  
<https://doi.org/10.1073/pnas.1304801110>

Vanhaesebroeck B, Perry MWD, Brown JR, André F, Okkenhaug K (2021) **PI3K inhibitors are finally coming of age** *Nat Rev Drug Discov* **20**:741–769  
<https://doi.org/10.1038/s41573-021-00209-1>

Vasan N, Cantley LC (2022) **At a crossroads: how to translate the roles of PI3K in oncogenic and metabolic signalling into improvements in cancer therapy** *Nat Rev Clin Oncol*  
<https://doi.org/10.1038/s41571-022-00633-1>

Walker EH, Perisic O, Ried C, Stephens L, Williams RL (1999) **Structural insights into phosphoinositide 3-kinase catalysis and signalling** *Nature* **402**:313–320  
<https://doi.org/10.1038/46319>

Walser R, Burke JE, Gogvadze E, Bohnacker T, Zhang X, Hess D, Küenzi P, Leitges M, Hirsch E, Williams RL, Laffargue M, Wymann MP (2013) **PKCβ phosphorylates PI3Ky to activate it and release it from GPCR control** *PLoS Biol* **11**  
<https://doi.org/10.1371/journal.pbio.1001587>

Zhang X , Vadas O , Perisic O , Anderson KE , Clark J , Hawkins PT , Stephens LR , Williams RL (2011) **Structure of lipid kinase p110 $\beta$ /p85 $\beta$  elucidates an unusual SH2-domain-mediated inhibitory mechanism** *Mol Cell* **41**:567–578  
<https://doi.org/10.1016/j.molcel.2011.01.026>

## Author information

### Noah J Harris

Department of Biochemistry and Microbiology, University of Victoria

### Meredith L Jenkins

Department of Biochemistry and Microbiology, University of Victoria

ORCID iD: [0000-0002-0685-5048](https://orcid.org/0000-0002-0685-5048)

### Sung-Eun Nam

Department of Biochemistry and Molecular Biology, The University of British Columbia, Vancouver

### Manoj K Rathinaswamy

Department of Biochemistry and Microbiology, University of Victoria

### Matthew AH Parson

Department of Biochemistry and Microbiology, University of Victoria

ORCID iD: [0000-0001-6270-559X](https://orcid.org/0000-0001-6270-559X)

### Harish Ranga-Prasad

Department of Biochemistry and Microbiology, University of Victoria

### Udit Dalwadi

Department of Biochemistry and Molecular Biology, The University of British Columbia, Vancouver

### Brandon E Moeller

Department of Biochemistry and Microbiology, University of Victoria

### Eleanor Sheekey

Department of Biochemistry and Microbiology, University of Victoria

### Scott D Hansen

Department of Chemistry and Biochemistry, Institute of Molecular Biology, University of Oregon

ORCID iD: [0000-0001-7005-6200](https://orcid.org/0000-0001-7005-6200)

### Calvin K Yip

Department of Biochemistry and Molecular Biology, The University of British Columbia, Vancouver

**For correspondence:** [calvin.yip@ubc.ca](mailto:calvin.yip@ubc.ca)

ORCID iD: [0000-0003-1779-9501](https://orcid.org/0000-0003-1779-9501)

**John E Burke**

Department of Biochemistry and Microbiology, University of Victoria, Department of Biochemistry and Molecular Biology, The University of British Columbia, Vancouver

**For correspondence:** [jeburke@uvic.ca](mailto:jeburke@uvic.ca)

ORCID iD: [0000-0001-7904-9859](https://orcid.org/0000-0001-7904-9859)

**Editors**

Reviewing Editor

**Amy Andreotti**

Iowa State University, United States of America

Senior Editor

**Volker Dötsch**

Goethe University, Germany

**Reviewer #1 (Public Review):**

The authors examine signaling factors that differentiate parallel routes to activating phosphoinositide 3-kinase gamma (PI3K $\gamma$ ). Dissecting the convergent pathways that control PI3K $\gamma$  activity is critical because PI3K $\gamma$  is a therapeutic target for treating inflammatory disease and cancer. Here, the authors employ a multipronged approach to reveal new aspects for how p84 and p101 pair with p110 $\gamma$  to activate the PI3K $\gamma$  heterodimer. The key instigator to this study is a previously reported inhibitory Nanobody, NB7. The hypothesized mechanism for NB7 allosteric inhibition of p84- p110 $\gamma$  was previously proposed to involve blockage of the Ras-binding domain. The authors revise the allosteric inhibition model based on meticulous profiling of various PI3K $\gamma$  complex interactions with NB7. In parallel, a cryo-EM-derived model of NB7 bound to the p110 $\gamma$  subunit convincingly reveals a Nanobody interaction pocket involving the helical domain and regulatory motifs of the kinase domain. This revelation shifts the focus to the helical domain, a known target of PKC phosphorylation. While the connections between NB7 interactions and the effects of PKC phosphorylation are sometimes tenuous, it could be argued that the Nanobody served as a tool to reveal the importance of the helical domain to p110 $\gamma$  regulation.

The sites of PKC-mediated p110 $\gamma$  helical domain phosphorylation were unexpectedly inaccessible in the available structural models. Nevertheless, mass spectrometry (MS)-based phosphorylation profiling indicates that PKC can phosphorylate the helical domain of p110 $\gamma$  and p84/p110 $\gamma$  (but not p101/p110 $\gamma$ ) in vitro. The authors hypothesize that helical domain dynamics dictate susceptibility to PKC phosphorylation. To explore this notion, carefully executed, rigorous H/D exchange MS (HDX-MS) experiments were performed comparing phosphorylated vs. unphosphorylated p110 $\gamma$ . Notably, this design reveals more about the consequences of p110 $\gamma$  phosphorylation, rather than the mechanisms of p84/p101 promoting/resisting phosphorylation. Nevertheless, HDX-MS is very well suited to exploring secondary structure dynamics, and helical domain phosphorylation strikingly increases dynamics consistent with increased regional accessibility. The increased dynamics also nicely map to the pocket enveloped by the inhibitory NB7 Nanobody.

Ultimately, this study reveals an unexpected p110 $\gamma$  pocket that allows an engineered Nanobody to allosterically inhibit PI3K $\gamma$  complexes. The cryo-EM characterization of the interaction inspired an HDX-MS investigation of known sites of phosphorylation in the

region. These insights could be linked to differences/convergences of p84 and p101 complex formation and activation of PI3K $\gamma$ , and future work may clarify these mechanisms further. The data presented herein will also be useful for broadening the target surface for future therapeutic developments. New allosteric connections between effector binding sites and post-translational modifications are always welcome.

## Reviewer #2 (Public Review):

Harris et al. have described the cryo-EM structure of PI3K p110 $\gamma$  in a complex with a nanobody that inhibits the enzyme. This provided the first structure of full-length of PI3K $\gamma$  in the absence of a regulatory subunit. This nanobody is a potent allosteric inhibitor of the enzyme, and might provide a starting point for developing allosteric, isotype-specific inhibitors of the enzyme. One distinct effect of the nanobody is to greatly decrease the dynamics of the enzyme as shown by HDX-MS, which is consistent with a growing body of observations suggesting that for the whole PI3K superfamily, enzyme activators increase enzyme dynamics.

The most remarkable outcome of the study is that upon observing the site of nanobody binding, the authors searched the literature and found that there was a previous report of a PKC $\beta$  phosphorylation of PI3K $\gamma$  in the helical domain that is near the nanobody binding site. This led the authors to re-examine the consequence of the phosphorylation armed with better structural models and the tools to study the effects of this phosphorylation on enzyme dynamics. They found that the site of phosphorylation is buried in the helical domain, suggesting that a large conformational change would have to take place to enable the phosphorylation. HDX-MS showed that phosphorylation at three sites clustered in the helical domain generate a distinctly different conformation with rapid deuterium exchange. This suggests that the phosphorylation locks the enzyme in a more dynamic state. Their enzyme kinetics show that the phosphorylated, dynamic enzyme is activated.

While this phosphorylation was reported before, the authors have provided a mechanism for why this activates the enzyme, and they have shown why binders that stabilise the helical domain (such as binding to the p101 regulatory subunit and the nanobody) prevent the phosphorylation. It is this insight into the dynamics of the PI3K $\gamma$  that will likely be the long-lasting influence of the work.

The paper is well written and the methods are clear.

## Author Response:

The following is the authors' response to the original reviews.

We appreciate the in depth review of our manuscript, and the excellent suggestions from the two reviewers. We have addressed all concerns as described in the point by point response below. We have also added all of these changes to a revised version posted to biorxiv on May 23rd 2023 (BIORXIV/2023/536585).



### **Reviewer #1 (Recommendations For The Authors):**

*It is sometimes difficult to connect the rationalizations behind the transitions between NB7 binding interaction, the compare/contrast of p84 and p101 effectors, and the synergy with phosphorylation. More explanation of the rationalizations behind these transitions in the Results would be helpful.*

We agree that the manuscript would benefit from better transitions between the sections. We have added a new paragraph in the final section describing the nanobody structure before the helical domain phosphorylation that fully describes the rationale for how both inform on the critical role of helical domain dynamics in kinase activity. This paragraph is shown below.

*'The interface of NB7 with p110\_g is distant from both the putative membrane binding surface, as well as the catalytic machinery of the kinase domain. To further understand how this nanobody could so potently inhibit PI3K activity we examined any other potential modulators of PI3K activity localised in this region. There are two regulatory phosphorylation sites in the helical (Walser et al., 2013) and kinase domain (Perino et al., 2011) localised at the NB7 interface. This is intriguing as helical domain phosphorylation is activating, and kinase domain phosphorylation is inhibitory. This suggested a critical role in the regulation of p110\_g is the dynamics of this kinase-helical interface. To fully define the role of NB7 in altering the dynamics of the helical domain we needed to study other modulators of helical domain dynamics.'*

*The Methods section would benefit from careful copy editing for clarity and consistency.*

We have gone through the methods section and edited for clarity and consistency throughout.

*There's a minor ambiguity throughout when referring to the phosphorylation of S594/S595. Although close inspection makes it clear that this refers to the monophosphorylation of either site S594 or S595, there are several references to "S594/S595" that could be interpreted as phosphorylation of both residues.*

We agree that this was ambiguous in the original text. We have added an explicit statement describing this as a single phosphorylation event.

*The modification at this site results in a single phosphorylation event, but due to CID MS/MS fragmentation we cannot determine which site is modified, and will be described as S594/S595 throughout the manuscript.'*

*In Figure 2B, the authors show the cryo-EM density map and the structural model based on this map. It would be helpful to also include an image of the structural model fit in the density map to allow readers to evaluate the quality of the map and model. The 2F panel provides an important view of this fit, but CD3 models are difficult to discern.*

We agree that this would help interpret model quality. We have added a new supplemental figure showing the fit of both p110 and NB7 into this Cryo EM density (see new Fig S2).

*Paragraph starting at line 258: The shift to monitoring ATPase activity is confusing here. ATPase activity indicates production of ADP + phosphate (rather than ADP + PIP3). However, an explanation is provided that states that measuring ADP production serves as a surrogate for measuring PIP3 production. The apparent absence of membrane PIP2 substrate in Figure 4E (left) suggests that there is a true ATPase background activity in this kinase. If so, does the increase in ADP production in Fig. 4E reflect the inclusion of PIP2 substrate, increased background ATPase activity, or both?*

We agree that this was worded confusingly in the original version. We have now clarified exactly what we are observing in these ATPase assays. The new paragraph is appended below

*To further explore the potential role of phosphorylation in mediating p110g activity, we examined the kinase activity of p110g under two conditions: basal ATP turnover, and with PIP2 containing lipid membranes. The experiments in the absence of PIP2 measure turnover of ATP into ADP and phosphate, and is a readout of basal catalytic competency. Experiments with PIP2, measured ATP consumed in the generation of PIP3, as well as in non-productive ATP turnover. The p110g enzyme in the absence of stimulators is very weakly active towards PIP2 substrate with only ~2 fold increased ATP turnover compared to in the absence of membranes. This is consistent with very weak membrane recruitment of p110g complexes in the absence of lipid activators (Rathinaswamy et al., 2023). PKCb-mediated phosphorylation enhanced the ATPase activity of p110g ~2-fold in both the absence and presence of membranes (Fig. 4E). This suggests that the effect of phosphorylation is to change the intrinsic catalytic efficiency of phosphorylated p110g, with limited effect on membrane binding.'*

*In the section "Nanobody blocks p110gamma phosphorylation," it's not entirely convincing that "the presence of NB7 showed even lower phosphorylation than p110gamma-p101." This does not seem to be subject to a significance test in Figure 5A/B. The follow-up point about "complete abrogation of phosphorylation," however, is readily apparent.*

We agree that we could have been more precise with our language, as this is not a complete block of phosphorylation, it is merely a significant decrease in phosphorylation. We have removed the comparison to p110-p101, and also removed the statement about complete abrogation of phosphorylation. This is now reworded to

*The presence of NB7 showed a significant decrease in p110g phosphorylation at both sites (Fig. 5A-B).'*

*Figure 1: Legend needs to include more detail to define the data. (1A) Representations of variance need to be clarified (e.g., replicates, error bar meaning). Consider "Normalized lipid kinase activity" as a y-axis label and expand on the activity measurement and normalization in the legend. (1B) How was error calculated? (1C) Mislabeled as 1B? Also, consider clarifying the first title highlighting the comparison to class IA PI3Ks. (1D) Typo: "Y647-p84/p110gamma." Also, would it not be more accurate to say "effect of nanobody NB7 on PI3K displacement..." for this experiment?*

We apologise for these oversights. See details on what has been changed in Fig 1A, 1B, 1C, and 1D.

For Fig 1A we now show the data where each replicate is indicated in the graph in the absence of error bars, and have also more clearly expanded on this activity measurement in the figure legend and also stated the replicate number.

For Fig 1B we now clearly state how the error was generated.

For Fig 1C we have fixed the typo

For Fig 1D, we have fixed this typo and also changed the sub-heading as suggested.

New figure legend is below as well

**Figure 1. The inhibitory nanobody NB7 binds tightly to all p110 $\gamma$  complexes and inhibits kinase activity, but does not prevent membrane binding**

**A.** Cartoon schematic depicting nanobody inhibition of activation by lipidated G $\beta\gamma$  (1.5  $\mu$ M final concentration). Lipid kinase assays show a potent inhibition of lipid kinase activity with increasing concentrations of NB7 (3–3000 nM) for the different complexes. Experiments are carried out in triplicate ( $n=3$ ) with each replicate shown. The y-axis shows lipid kinase activity normalised for each complex activated by G $\beta\gamma$  in the absence of nanobody. Concentrations of each protein were selected to give a lipid kinase value in the detectable range of the ATPase transcreener assay. The protein concentration of p110 $\gamma$  (300 nM), p110 $\gamma$ -p84 (330 nM) and p110 $\gamma$ -p101 (12 nM) was different due to intrinsic differences of each complex to be activated by lipidated G $\beta\gamma$ , and is likely mainly dependent for the difference seen in NB7 response.

**B.** Association and dissociation curves for the dose response of His-NB7 binding to p110 $\gamma$ , p110 $\gamma$ -p84 and p110 $\gamma$ -p101 (50 – 1.9 nM) is shown. A cartoon schematic of BLI analysis of the binding of immobilized His-NB7 to p110 $\gamma$  is shown on the left. Dissociation constants (KD) were calculated based on a global fit to a 1:1 model for the top three concentrations and averaged with error shown. Error was calculated from the association and dissociation value ( $n=3$ ) with standard deviation shown. Full details are present in the source data.

**C.** Association and dissociation curves for His-NB7 binding to p110 $\gamma$ , p110 $\alpha$ -p85 $\alpha$ , p110 $\beta$ -p85 $\beta$ , and p110 $\delta$ -p85 $\delta$ . Experiments were performed in duplicate with a final concentration of 50 nM of each class I PI3K complex.

**D.** Effect of NB7 on PI3K recruitment to supported lipid bilayers containing H-Ras(GTP) and farnesyl-Gbg as measured by Total Internal Reflection Fluorescence Microscopy (TIRF-M). DY647-p84/p110g displays rapid equilibration kinetics and is insensitive to the addition of 500 nM nanobody (black arrow, 250 sec) on supported lipid bilayers containing H-Ras(GTP) and farnesyl-Gbg.

**E.** Kinetics of 50 nM DY647-p84/p110g membrane recruitment appears indistinguishable in the absence and presence of nanobody. Prior to sample injection, DY647-p84/p110g was incubated for 10 minutes with 500 nM nanobody.

**F.** Representative TIRF-M images showing the localization of 50 nM DY647-p84/p110g visualized in the absence or presence of 500 nM nanobody (+NB7). Membrane composition for panels C-E: 93% DOPC, 5% DOPS, 2% MCC-PE, Ras(GTP) covalently attached to MCC-PE, and 200 nM farnesyl-Gbg.

Figure 2: (1A) For consistency with the rest of the paper, p110g can be updated with the Greek character. (1B) This may have been intentional to attract attention to subdomains interacting with NB7, but "colored according to the schematic" omits the purple RBD. (2F) the figure legend should specify whether p110gamma surfaces depicted are the cryo-EM density or a surface rendition of the structural model.

We agree and have fixed the p110 typo, and have also colored the schematic the same as shown in the cartoon model.

The data shown in Fig 1B is indeed the Cryo EM density and this is now clearly indicated in the legend.

*Figure 3: (3B) Specifying the [M+H] as [M+2H]<sup>2+</sup> and [M+4H]<sup>4+</sup> would help the reader understand the delta mass for monophosphorylation here. Given the broad readership of this journal, it would be useful to define 't' and 'e' as 'theoretical' and 'experimental' in the legend. It may also help to be explicit about the meaning of the red spectra and residues in the legend. (3C-E) autocorrect typo for "(C)" and an opportunity to update "b" for Greek character beta.*

We agree that clearly defining the charge state of each spectra will make it more obvious that we are dealing with a mono-phosphorylation and have made this change as suggested in the figure. We have also clearly define m/z t and m/z e in the figure legend, as well as the black and red lines, and characters. Finally we have added PKCb for all descriptions of PKC treatment in the figure, and fixed the incorrect PKC'b' in the legend.

*Figure 4: (4C) Given the common use of "ND" for other terms, it would be useful to spell out "no deuterium" or "undeuterated." (4E) the parenthetical "(concentration, 12nM to 1000nM)" could be clarified. How are the (presumably p100gamma) concentration ranges reflected in the three plotted data points per treatment? See also Figure 5E.*

We agree and have redefined ND as undeuterated. We apologise for the typo in the figure legend, as the concentrations of p110 gamma were the same for both phosphorylated and non-phosphorylated, with this being a typo (all concentrations of enzyme were 1000 nM). This has been changed here and in Fig 5E.

*Figure 5: (5A/B) Some clarification that we're looking at extracted ion chromatograms would be very useful in this figure legend. On a related note, the experimental details on the LC-MS methodology for this data appear to be split between two sections of the Methods: the "Phosphorylation analysis" paragraph (line 526) and the HDX-MS section. Some more explicit cross-referencing would clarify this experiment. (5E) Clarify inclusion of PIP2 membranes here.*

We have clearly described that we are looking at extracted ion chromatograms in both panel A and B. We also have normalized the experimental methods in the LC-MS as these used exactly the same procedure. Finally, we now clearly describe the assays shown in Fig 5E were performed in the absence of PIP2 membranes.

*Miscellaneous typos:  
Line 205: reference omitted for "Previous study.."*

We have added this reference

*Line 196: "unambiguous"*

Fixed to unambiguously

### **Reviewer #2 (Recommendations For The Authors):**

*The only mistake I spotted was that on line 729 there is a reference to Fig 3C that should actually be Fig. 4C*

We have changed this to the correct Fig 4C.