



# The catalytic mechanism of the RNA methyltransferase METTL3

Ivan Corbeski, Pablo Andrés Vargas-Rosales, Rajiv Kumar Bedi, Jiahua Deng, Dylan Coelho, Emmanuelle Braud, Laura Iannazzo, Yaozong Li, Danzhi Huang, Mélanie Ethève-Quelquejeu, Qiang Cui, Amedeo Caflisch 

Department of Biochemistry, University of Zurich, Zurich CH-8057, Switzerland • Department of Chemistry, Boston University, Boston, Massachusetts 02215, United States • Université Paris Cité, CNRS, Laboratoire de Chimie et Biochimie Pharmacologiques et Toxicologiques, Paris F-75006, France • Department of Physics, Boston University, Boston, Massachusetts 02215, United States • Department of Biomedical Engineering, Boston University, Boston, Massachusetts 02215, United states

 [https://en.wikipedia.org/wiki/Open\\_access](https://en.wikipedia.org/wiki/Open_access)

 Copyright information

## Reviewed Preprint

Revised by authors after peer review.

## About eLife's process

## Reviewed preprint version 2

February 22, 2024 (this version)

## Reviewed preprint version 1

November 30, 2023

## Sent for peer review

September 27, 2023

## Posted to preprint server

September 6, 2023

## Abstract

The complex of methyltransferase-like proteins 3 and 14 (METTL3-14) is the major enzyme that deposits N<sup>6</sup>-methyladenosine (m<sup>6</sup>A) modifications on mRNA in humans. METTL3-14 plays key roles in various biological processes through its methyltransferase (MTase) activity. However, little is known about its substrate recognition and methyl transfer mechanism from its cofactor and methyl donor S-adenosylmethionine (SAM). Here, we study the MTase mechanism of METTL3-14 by a combined experimental and multiscale simulation approach using bisubstrate analogues (BAs), conjugates of a SAM-like moiety connected to the N<sup>6</sup>-atom of adenosine. Molecular dynamics simulations based on crystal structures of METTL3-14 with BAs suggest that the Y406 side chain of METTL3 is involved in the recruitment of adenosine and release of m<sup>6</sup>A. A crystal structure with a bisubstrate analogue representing the transition state of methyl transfer shows a direct involvement of the METTL3 side chains E481 and K513 in adenosine binding which is supported by mutational analysis. Quantum mechanics/molecular mechanics (QM/MM) free energy calculations indicate that methyl transfer occurs without prior deprotonation of adenosine-N<sup>6</sup>. Furthermore, the QM/MM calculations provide further support for the role of electrostatic contributions of E481 and K513 to catalysis. The multidisciplinary approach used here sheds light on the (co)substrate binding mechanism, catalytic step, and (co)product release catalysed by METTL3, and suggests that the latter step is rate-limiting. The atomistic information on the substrate binding and methyl transfer reaction of METTL3 can be useful for understanding the mechanisms of other RNA MTases and for the design of transition state analogues as their inhibitors.

## One sentence summary

A combined structural, biochemical, and computational approach reveals two distinct binding conformations of adenosine, the transition state of methyl transfer, and the catalytic mechanism of the human m<sup>6</sup>A RNA methyltransferase METTL3-14.

## eLife assessment

This **important** study combines experimental and computational data to address crucial aspects of RNA methylation by a vital RNA methyltransferase (MTase). The authors have provided **compelling**, strong evidence, utilizing well-established techniques, to elucidate aspects of the methyl transfer mechanism of methyltransferase-like protein 3 (METTL3), which is a part of the METTL3-14 complex. This work will be of broad interest to biochemists, biophysicists, and cell biologists alike.


## Introduction

### METTL3-14 is the main human mRNA m<sup>6</sup>A MTase

There are more than 170 RNA modifications forming the epitranscriptome.<sup>1</sup> N<sup>6</sup>-methyladenosine (m<sup>6</sup>A) is the most frequent internal modification of messenger RNA (mRNA) within the consensus sequence GGACU that is enriched near stop codons and in 3' untranslated regions.<sup>2-4</sup> m<sup>6</sup>A affects most aspects of RNA regulation, i.e., alternative polyadenylation<sup>5</sup>, splicing<sup>6</sup>, nuclear export<sup>7</sup>, stability<sup>8</sup>, and translation initiation.<sup>3,9</sup> The complex of methyltransferase-like protein 3 (METTL3) and METTL14 (abbreviated as METTL3-14 in the following) is the main m<sup>6</sup>A-RNA methyltransferase (MTase).<sup>10</sup>

The METTL3-14 heterodimer is involved in a wide variety of diseases including type 2 diabetes,<sup>11</sup> viral infections,<sup>12</sup> and several types of cancer.<sup>13</sup> METTL3-mediated m<sup>6</sup>A deposition is directly involved in the development of acute myeloid leukaemia (AML) by promoting the translation of genes involved in cell growth, differentiation, and apoptosis.<sup>14,15</sup> It has been demonstrated that inhibition of the METTL3 catalytic function is sufficient to induce apoptosis and differentiation in AML cells and in a mouse model of the disease but not in normal non-leukaemic haematopoietic cells.<sup>16,17</sup> While it has been well established that METTL3-14 dysregulation is related to cancer development, the role of METTL3-14 varies in different cancer types, i.e., it can act as oncogene or tumour suppressor.<sup>18</sup> Despite growing knowledge of the diverse pathways involving METTL3-14, the mechanism of how m<sup>6</sup>A regulates gene expression remains poorly understood. Little is known about the recognition of specific RNA transcripts, the binding and release of the adenosine substrate, and the methyl transfer mechanism catalysed by METTL3. Furthermore, inhibiting the MTase function of METTL3-14 is a promising therapeutic strategy for several diseases.<sup>17,19</sup> Hence, understanding the mechanism of this complex would be helpful to develop new therapies.

METTL3-14 is the catalytic complex that transfers the methyl group from S-adenosylmethionine (SAM) to the substrate adenosine (**Figure 1**).<sup>20-23</sup> METTL3 comprises a low-complexity region at the N-terminus, a zinc finger responsible for substrate binding, and the catalytic MTase domain at the C-terminus (**Figure 1A**).<sup>24</sup> The METTL3 MTase domain has the catalytically active SAM binding site and adopts a Rossmann fold that is characteristic of Class I SAM-dependent MTases (**Figure 1B**). METTL14 has an MTase domain, too, however, with a redundant active site of hitherto unknown function, and so-called RGG repeats at its C-terminus essential for RNA binding.<sup>23</sup> METTL14 plays a structural role for complex stabilization and RNA-binding. It forms a positively charged groove at the interface with METTL3 which is predicted to be the RNA binding site.



**Figure 1.**

**METTL3-14 domain architecture and structure.**

**(A)** Domain architecture of METTL3 and METTL14. ZnF = zinc finger, NHM = N-terminal  $\alpha$ -helical motif, CTM = C-terminal motif, RGG = arginine-glycine-glycine motif. **(B)** Crystal structure of the MTase domains of METTL3-14. Ribbon representations (left) are coloured as in panel (A). Surface renderings (right) are coloured according to the electrostatic potential. SAM and the putative RNA binding site are indicated.

## MD simulations for mechanistic studies of RNA MTases

Thus far, and to the best of our knowledge, no computational study of the conformational landscape or catalytic mechanism of eukaryotic RNA MTases has been reported. Molecular dynamics (MD) studies have mainly focused on protein MTases, and MTases from bacteria.<sup>25–28</sup> In the latter, dynamic cross correlation analysis, a technique usually applied for the study of allosteric processes,<sup>29</sup> showed that MTase conformational changes can influence the orientation of the substrate.<sup>28</sup> In the former, Chen et al. explored the conformational landscape of SETD8, a histone MTase.<sup>30</sup> The study showed how slow conformational motions and conformational states of the MTase are relevant to catalysis.


Among RNA MTases, research has focused on viral enzymes. One example is the MD simulation analysis of the binding mechanism of SAH and m<sup>7</sup>GTP to the Zika virus NS5 protein.<sup>31</sup> This enabled a detailed analysis of their interaction and understanding of the effects of MTase inhibitors as antiviral drugs. Another study focused on the mechanisms of the SARS-CoV-2 MTase nsp16 and its heterodimeric partner nsp10 which acts as a stimulator of SAM binding.<sup>32</sup> The study provides a comprehensive understanding of the dynamic, thermodynamic, and allosteric processes of MTase complex formation and function.

## Bisubstrate analogues as structural tools to investigate the mechanisms of MTases

Only a few structures of RNA-bound MTases are currently available due to the intrinsic instability of RNA and RNA-enzyme complexes and resulting difficulties in obtaining their structures. Of the known m<sup>6</sup>A RNA MTases, only METTL16, a SAM homeostasis factor, has been crystallized in complex with substrate RNA.<sup>33</sup> The lack of more such structures results in poorly understood RNA recognition and methyl transfer mechanisms. In contrast, structures are available for many MTases in complex with either the cosubstrate SAM or the coproduct S-adenosylhomocysteine (SAH), allowing for a well-understood cofactor binding within this protein family. Cofactor binding guided the initial design of bisubstrate analogues (BAs) as chemical tools to study the catalytic mechanisms of m<sup>6</sup>A-MTases (**Figure 2**).<sup>34</sup>

BAs aim to mimic the transition state in which both the substrate nucleoside and the cosubstrate SAM are bound in the catalytic pocket of the enzyme while the methyl group is transferred from SAM to the adenosine N<sup>6</sup>-atom of the substrate RNA during catalysis (**Figure 2A**).<sup>34–38</sup> They consist of a SAM analogue (5'-N-SAM) covalently linked to the N<sup>6</sup>-position of an adenosine of a ribonucleotide(-like) fragment (**Figure 2B**).<sup>35,39</sup> The only structural information on these molecules is their binding mode in the bacterial m<sup>6</sup>A RNA MTase RlmJ.<sup>34,37</sup> There, in the mononucleoside containing compounds BA2 and BA4, the substrate adenosine is positioned in the presumed substrate binding pocket of RlmJ, and for the cofactor moiety, the methionine chain is bound like SAM. However, the structural studies resulted in a (co)substrate conformation that is not always biologically relevant, as the adenosine of the SAM analogue was rotated out of the canonical SAM binding pocket caused by  $\pi$ -stacking with the substrate adenine ring.<sup>34</sup> Therefore, the binding modes revealed therein are diverse and not always suitable for mechanistic studies. In a subsequent study, using a dinucleotide containing BA (GA\* in **Figure 2B**), the SAM moiety had the correct orientation.<sup>37</sup> Furthermore, the N<sup>6</sup>-atom of adenosine was positioned, through the alkyl chain of the linker, at 3 Å distance from the carbon corresponding to the C $\epsilon$ -atom of the methionine moiety in the cosubstrate SAM. Such a distance, in two non-linked moieties, would allow for an SN2 methyl transfer from SAM to the N<sup>6</sup>-atom of adenosine.<sup>40</sup> We therefore hypothesized that targeting human RNA MTases, in particular METTL3, with BAs might lead to structures that provide a suitable basis for understanding their MTase mechanism.

In the present study, we use a multidisciplinary approach to study the substrate binding, m<sup>6</sup>A methylation reaction, and release of m<sup>6</sup>A product in the human RNA MTase METTL3-14. To shed light on its catalytic mechanism we combine crystal structures of METTL3-14-bisubstrate analogue



**Figure 2.**

### Bisubstrate analogues as transition state mimics for METTL3.

**(A)** METTL3-catalysed transfer of the methyl group of SAM to the N<sup>6</sup>-atom of A in a GGACU-motif containing mRNA and the production of m<sup>6</sup>A and SAH. The inset shows the design principle of bisubstrate analogues (BAs) as transition state analogues. The point of linkage in the BA is indicated with a double arrow (red) between the N<sup>6</sup>-atom of adenosine and 5'-N of the SAM analogue. **(B)** Chemical structures of the BAs used in this study. Substrate adenosine = black; SAM analogue = blue, linker = red. Compound names are as previously published: BA1/2/3/4/6, ref. [34](#); Compound 12, ref. [36](#); GA\*, ref. [37](#).

complexes with in vitro experiments, multiscale atomistic simulations, namely classical molecular dynamics (MD) and quantum mechanics/molecular mechanics (QM/MM) free energy calculations. Crystal structures show the binding mode of the substrate adenosine in two different conformations, representing an encounter complex of RNA binding and the transition state of catalysis, respectively. These structures are validated through mutational analysis. Classical MD simulations are used to investigate the binding of the substrates SAM and adenosine and dissociation of the products SAH and  $m^6A$ . QM/MM free energy calculations reveal the details of the methylation reaction. Taken together, we elucidate the reaction catalysed by METTL3-14 at atomic level of detail. This knowledge will help in the further investigation of other MTases and the optimization of chemical probes that target their function.

## Results and Discussion

### Bisubstrate analogues bind in the METTL3 active site

We evaluated a series of bisubstrate analogues (BAs) as catalytic inhibitors and investigated the structural similarity between their binding mode in METTL3-14 and the putative RNA substrate and SAM cosubstrate during methyl transfer (see **Figure 2B**). First, we measured the inhibitory activity of the BAs on METTL3-14 enzymatic activity (**Table 1** and **Supplementary Figure S1**). For this, we used our in-house developed reader-based enzymatic assay that quantifies the  $N^6$ -adenosine methyl transfer based on homogeneous time-resolved fluorescence (HTRF).<sup>41</sup> The assay detects  $m^6A$  using the natural  $m^6A$ -reader YTHDC1. The  $m^6A$ -RNA and -reader are fluorescently labeled such that their proximity during binding causes Förster resonance energy transfer (FRET). The BAs that inhibit METTL3 reduce the  $m^6A$  level, and thus decrease the FRET signal, in a dose-dependent manner. The low micromolar inhibitory activity of the most potent BAs motivated us to conduct a structural investigation on their binding mode to METTL3-14. We conducted crystallization trials and obtained crystal structures of four of the BAs (BA1, BA2, BA4 and BA6) by soaking them into METTL3-14 crystals (see **Table 1** and **Supplementary Table S1**).

All the crystallized BAs bind in the METTL3 active site (**Figure 3A**). BA1 and BA6, that are missing the methionine part of the SAM analogue or have a polar urea group in the linker, respectively, have the highest  $IC_{50}$  values of the crystallized compounds in the enzymatic assay (see **Table 1**). Furthermore, in their crystal structures with METTL3, they reveal divergent interactions of their SAM-like moiety compared to SAM (**Supplementary Figure S2**). For BA2 and BA4, however, the interaction of their SAM-like moiety is the same as for SAM (**Figure 3B-F** and **Supplementary Figure S3**). Both ligands bind with their SAM moiety in the active site lined with METTL3 hydrophobic residues and form polar contacts to conserved residues, namely D395, R536, H538, and N539 via the methionine part and D377, I378, F534, N549, and Q550 via the adenosine moiety of 5'-N-SAM. These structures therefore validate their further analysis for the interpretation of the catalytic mechanism of METTL3.

### Adenosine has two distinct binding conformations in the METTL3 active site

In the METTL3-BA2 and -BA4 structures, the SAM moiety is superimposable with the METTL3-bound conformation of SAM, and adenosine is bound in the presumed catalytic site (**Figure 4A**). Adenosine, based on BA2 and BA4, is involved in an intricate network of interactions with side chains of METTL3 (**Figure 4B**).

The classical  $m^6A$  MTase catalytic <sup>395</sup>DPPW<sup>398</sup> motif (D/N-PP-Y/F/W) of METTL3 in its flexible active site loop 1 (ASL1, METTL3 residues 395-410) participates to the binding of adenosine. D395 forms a salt bridge to SAM and its mutation to alanine was previously shown to abolish METTL3

Bisubstrate analogue name <sup>a</sup>	IC <sub>50</sub> [μM] <sup>b</sup> ± SD	PDB ID <sup>c</sup>	Resolution <sup>c</sup> [Å]
<b>BA1</b>	346 ± 66	8PW9	2.3
<b>BA2</b>	9 ± 1	8PW8	2.3
<b>BA3</b>	21 ± 2	NA	NA
<b>BA4</b>	18 ± 3	8PWA	2.1
<b>BA6</b>	≈ 500	8PWB	2.5
<b>Compound 12</b>	≈ 500	NA	NA
<b>GA*</b>	32 ± 3	NA	NA

NA = Not available

a. Compound names as previously published: BA1/2/3/4/6, ref. <sup>34</sup>; Compound 12, ref. <sup>36</sup>; GA\*, ref. <sup>37</sup>.


b. IC<sub>50</sub> = Half maximal inhibitory concentration from the enzymatic assay; for comparison, IC<sub>50</sub> of SAH is 0.51 μM.<sup>41</sup>

c. Data for the METTL3-14-BA complex structures deposited to the Protein Data Bank (PDB).

**Table 1.**

**BAs for METTL3-14 characterized in this study.**






**Figure 3.**

### Crystal structures of METTL3-14 show that BAs bind in the METTL3 active site.

(A) Superposition of the crystal structures of METTL3-14 bound to SAM and the four BAs. METTL3 backbone is shown as ribbon, side chains involved in polar interactions with SAM or intramolecularly are shown as sticks, waters as red spheres. SAM (cyan) is shown as sticks and indicated, and BAs are shown as transparent sticks (BA1 = yellow, BA2 = magenta, BA4 = salmon, BA6 = palegreen). Black dashes indicate polar contacts between METTL3 and SAM in the crystal structure. (B) Outline of METTL3/SAM interactions from a LigPlot+ analysis.<sup>42,43</sup> Black dashed lines indicate polar contacts between METTL3 and SAM in the crystal structure, residues forming the binding pocket environment are shown in grey. (C) Structure of METTL3-BA2. Figure composition as in (A). BA2 is coloured magenta, its SAM and adenosine moieties are indicated. The green dashes indicate a hydrogen bond that does not form with BA2, but is likely to have favourable geometry to form between D395 and adenosine-N<sup>6</sup> (i.e., the NH<sub>2</sub> group) of the natural RNA substrate. (D) Outline of METTL3/BA2 interactions from a LigPlot+ analysis, as in (B). The SAM analogue and adenosine parts of the BA are indicated. Black lightnings highlight residues in METTL3 involved in hydrophobic contacts with the adenosine moiety of the BA. (E) Structure of METTL3-BA4. Figure composition as in (D). BA4 is coloured salmon, its SAM and adenosine moieties are indicated. Note that BA4 is missing the ribose of the substrate adenosine moiety due to lack of electron density in the crystal structure probably due to flexibility of this group. (F) Outline of METTL3/BA4 interactions from a LigPlot+ analysis, as in (D). The missing ribose of the substrate adenosine moiety in the crystal structure is indicated with a lighter colour.





**Figure 4.**

### Crystal structures of METTL3-14 with BA2 and BA4 reveal two distinct adenosine binding modes.

**(A)** Superposition of the structures of SAM (cyan), BA2 (magenta) and BA4 (salmon) bound to METTL3. The ligands and their moieties are indicated. METTL3 in light/dark grey for BA2/BA4, backbone is shown as ribbon with side chains involved in the interactions with the adenosine moiety of the BAs shown as sticks. Waters are shown as red spheres. Black dashes indicate polar contacts common to BA2/BA4. Magenta/salmon dashes indicate hydrogen bonds unique to BA2/BA4. The green dashes indicate a hydrogen bond that does not form with BA2, but is likely to have favourable geometry to form between D395 and adenosine-N<sup>6</sup> (i.e., the NH<sub>2</sub> group) of the natural RNA substrate. Note that there is no electron density for the side chain of Y406 in the complex with BA2 and for the ribose of the adenosine moiety in the complex with BA4 which is most likely due to flexibility of these groups. **(B)** Ligplot+ analysis showing key interactions between METTL3, adenosine, and SAM based on the BA2 and BA4 structures. Dashed lines indicate polar contacts as in (A). Small magenta/salmon lightnings highlight residues in METTL3 involved in hydrophobic contacts with the adenosine moiety in the BA2/BA4 conformation. **(C)** Mutational analysis of the enzymatic activity of METTL3 active site residues involved in adenosine binding. **(D)** The melting temperature ( $T_m$ ) and its shift ( $\Delta T_m$ , in red) for METTL3 WT and mutants with DMSO as control (light grey bars) or in the presence of SAH (dark grey bars) measured using differential scanning fluorimetry. The error bars represent standard deviation from triplicate measurements.

activity and SAM binding, confirming its involvement in cosubstrate binding.<sup>20,22</sup> The METTL3-BA2 structure reveals that there can also be a hydrogen bond formed between D395 and the N<sup>6</sup>-atom of adenosine in the case of the non-alkylated RNA substrate (**Figure 4B**). P396 in the <sup>395</sup>DPPW<sup>398</sup> motif, through its carbonyl group, forms a hydrogen bond to the N<sup>6</sup>-atom of adenosine. In addition, P397 from the <sup>395</sup>DPPW<sup>398</sup> motif makes hydrophobic contacts with SAM. The BA2 structure further shows that adenosine is stabilized by additional hydrogen bonds from its N<sup>7</sup>- and O2'-atoms to the METTL3 side chains of E481 and K513, respectively. While W398 from the <sup>395</sup>DPPW<sup>398</sup> motif is not directly involved in adenosine binding, its backbone carbonyl group forms a water-bridged hydrogen bond with the Y406 side chain in the BA4 structure. The latter is involved in hydrophobic interactions with adenosine in the BA4 structure whereas it remains flexible in the BA2-bound METTL3 structure (see **Figure 4A**).

Taken together, adenosine swaps conformation from solvent exposed in the METTL3-BA4 structure to buried in the METTL3-BA2 structure where it forms hydrogen bonds with residues in the active site loop 2 (ASL2, METTL3 residues 507-515). Each moiety of the BAs is involved in hydrophobic interactions with METTL3 residues, and both adenines (of SAM and adenosine) form hydrogen bonds to conserved residues in METTL3. Alanine mutation of D395, Y406, E481, and K513, which are involved in adenosine binding as seen in the BA2 and BA4 structures, almost completely abolishes the METTL3 catalytic activity (**Figure 4C**). Importantly, these residues are highly conserved in METTL3 (**Supplementary Figure S4**). The loss of activity of the Y406, E481, and K513 mutants originates mainly from abolished binding capability to adenosine and not SAM, because they can still bind SAH, as seen from a thermal shift assay (**Figure 4D**). Upon binding SAH, the thermal shift for the Y406, E481, and K513 mutants is similar as for the WT which suggests that their side chains are not involved in binding SAM/SAH but rather the RNA substrate. In contrast, mutation of D395 impairs SAH binding, which is consistent with the involvement of the D395 side chain in SAM binding.<sup>20</sup>

## Molecular dynamics simulations reveal BA2 as the stable adenosine binding pose

Because of the different conformations of METTL3 and adenosine in the BA2 and BA4 structures, we went on to characterize the enzyme dynamics in the presence of adenosine. We carried out multiple MD simulations of the METTL3-14 heterodimer in complex with (co)substrates and (co)products. Apo trajectories were generated in a previous study.<sup>43</sup> The MD simulations were started from the crystal structures of the complex with bisubstrate analogues BA2 and BA4 described above. Based on the position of each BA, the (co)substrates SAM and adenosine monophosphate (AMP) or the (co)products SAH and m<sup>6</sup>AMP were positioned in the protein, aligning them to their respective moiety of the BA. The interaction of each ligand to the protein, and of intramolecular polar contacts, was monitored throughout the trajectories. The dissociation time for each of the substrates and products was analysed and modelled by fitting a single exponential to the fraction of bound ligands (**Table 2** using block averaging and **Supplementary Figure S5** considering all trajectories). The SAM and SAH cofactors remained bound in all but one of the sampled trajectories and thus fitting was not possible. The experimentally determined  $k_{\text{off}}$  rates of SAM and SAH are  $8 \times 10^{-4} \text{ s}^{-1}$  and  $2 \times 10^{-4} \text{ s}^{-1}$ , respectively, which corresponds to mean lifetimes of more than 20 min and 83 minutes, respectively.<sup>44</sup> Hence, we would not expect to observe dissociations of SAM or SAH in our 500 ns MD simulations.

Simulations that were started with AMP or m<sup>6</sup>AMP in the conformation of BA4 showed immediate dissociation of these ligands (see **Table 2** and **Supplementary Figures S6** and **S7**). This could mean that this conformation represents a short-lived intermediate during substrate binding or product release. Possibly, it represents an encounter complex between the METTL3-SAM holo complex and RNA substrate. This could be promoted through electrostatic steering of the negatively charged RNA backbone by the positively charged sulfonium ion of SAM.

<i>Initial structure</i>	<i>Fit</i>	<i>Substrates</i>		<i>Products</i>	
		SAM	AMP	SAH	m <sup>6</sup> AMP
BA2	A	>> 500	426 ± 196	>> 500	12.87 ± 0.02
	B		579 ± 386 (0.9 ± 0.1)		12.70 ± 4.95 (1.1 ± 0.1)
BA4	A	>> 500	0	>> 500	0

**Table 2.**

**Kinetic parameters of ligand dissociation. Mean lifetime ( $\tau$ ) in ns of the analysed (co)substrates or (co)products as calculated from fitting of a single exponential (A) or an exponential with a multiplicative factor (in parentheses) (B).**

The simulations started from the BA2 conformation show that the AMP and m<sup>6</sup>AMP ligands dissociated in several of the analysed MD simulation runs (**Supplementary Figures S8 and S9**). Binding of AMP is more stable, while the methylated product dissociates more quickly (see **Table 2**). This difference originates, at least in part, from the long-range monopole-monopole electrostatic interaction between the positively charged SAM and the negatively charged AMP. After the methyl transfer reaction, SAH is neutral which does not favour interaction with m<sup>6</sup>AMP resulting in its quicker dissociation.

One caveat is that we simulate only a mononucleotide, resulting in much quicker dissociation of both the substrate AMP and product m<sup>6</sup>AMP than would be expected for the longer, canonical RNA substrate/product. To the best of our knowledge, there are no published affinities of AMP and m<sup>6</sup>AMP for METTL3-14 binding. Based on the mean lifetimes determined from the MD simulations, and assuming diffusion-limited association rates of 10<sup>9</sup> M<sup>-1</sup>s<sup>-1</sup>, the dissociation constants of AMP and m<sup>6</sup>AMP are ~2 and ~78 mM, respectively. In contrast to these low affinities, the long GGACU motif-containing mRNA can interact with the binding groove at the METTL3-14 interface with dissociation constants in the nanomolar range, resulting in a more stable complex.<sup>45</sup> Nevertheless, our model is useful as it emphasizes the differences in binding affinity due to the different electrostatic interactions between the SAM-AMP and SAH-m<sup>6</sup>AMP pairs. Besides the structural stability of the BA2-like pose of AMP, the MD simulations also suggest that AMP can transiently populate a binding mode similar to BA4 (**Supplementary Figure S10**). This further supports that BA4 could represent an intermediate binding conformation of adenosine.

## A polar interaction network stabilizes the BA2 conformation of adenosine in METTL3

The crystal structure of the complex with BA2 reveals a string of ionic interactions which involves charged side chains of METTL3 and the amino and carboxyl groups of the SAM analogue. The string consists of seven charged groups (four of which are positive): R536<sup>+</sup> – SAM COO<sup>-</sup> – SAM NH<sub>3</sub><sup>+</sup> – D395<sup>-</sup> – K513<sup>+</sup> – E481<sup>-</sup> – K459<sup>+</sup> (see **Figure 4A,B**). We decided to monitor these salt bridges and the following monopole-dipole interactions in the MD simulations: D395<sup>-</sup> – AMP-N<sup>6</sup>, K513<sup>+</sup> – AMP-N<sup>7</sup>, and E481<sup>-</sup> – AMP-2'OH (see **Supplementary Figures S6 to S9**). In the BA2 conformation, the polar interactions between AMP and the charged side chains D395 and K513 are stronger than the interaction between E481 and the 2' hydroxyl group of the ribose (see **Supplementary Figure S8**, black traces). For SAM, the interaction between its positively charged amine and D395 is more stable than the one between its negatively charged carboxy with R536 which can adopt different conformations (see **Supplementary Figure S8**, blue traces). The intramolecular salt bridges are also stable throughout the sampling (see **Supplementary Figure S8**, grey traces). The salt bridge D395<sup>-</sup> – K513<sup>+</sup> seems to be the most stable, while E481<sup>-</sup> – K459<sup>+</sup> seems to change frequently between direct and water-separated contacts. In METTL3 apo trajectories, the D395<sup>-</sup> – K513<sup>+</sup> salt bridge is not present in the initial structure but is formed during the course of the simulation (**Supplementary Figure S11**). The bond between E481 and K513 is stable but separated by water when SAM and AMP are bound. This contrasts with the METTL3 apo simulations, where this bond forms transiently. The simulations with the products reveal similar interactions except for the faster dissociation of m<sup>6</sup>AMP (see **Table 2** and **Supplementary Figure S9**). Another difference is the weaker interaction between SAH and D395 which fluctuates more than with SAM. The intramolecular interactions are also observed for the simulations initiated from the BA4 conformation of adenosine (see **Supplementary Figures S6 and S7**). These MD results validate the mechanistic interpretation of the BA-bound METTL3-14 crystal structures, and give a dynamic view of the behaviour of the complex before (substrate bound) and after (product bound) the methyl transfer reaction.

## The flexibility of METTL3 Y406 supports the recruitment of adenosine


In structures of METTL3-14 in the apo state or bound to SAH or SAM, METTL3 residue Y406 is found in different conformations.<sup>20–22</sup> One study suggested that Y406 makes a hydrogen bond with S511 in ASL2 and thereby caps the SAH product.<sup>22</sup> However, another study suggested that Y406 might be important for the interaction with nucleotide bases.<sup>21</sup> The different conformations of Y406 seen in our crystal structures with BA2 and BA4 support the latter and suggest an involvement of the Y406 side chain in RNA nucleotide binding, probably as a first step of RNA recognition (see **Figure 4A**). Indeed, mutation of Y406 to alanine (in this and a previous study) or a cysteine (in a previous study) abolishes MTase activity (see **Figure 4C**).

The flexibility of Y406 in the MD simulations was analysed by monitoring the distance between its hydroxyl oxygen atom and the backbone carbonyl oxygen of METTL3 residue W398. There is a water-bridged polar interaction between these two oxygen atoms in the BA4 crystal structure (see **Figure 3E**). In the MD simulations, Y406 transitions multiple times (on the  $\mu$ s time scale) from orientations far away from the W398 carbonyl oxygen (distance of  $\sim 15$  Å) to shorter distances of  $\sim 6$  Å (**Figure 5A**). This corresponds to a transition from the extended ASL1, with Y406 pointing outside the pocket, to a conformation compatible with the water-bridged hydrogen bond observed in the crystal structure with BA4. **Figure 5B** shows a superposition of frames of a BA2 substrate MD trajectory illustrating this behaviour. We observe that the adenine ring system of AMP can be involved in a  $\pi$ - $\pi$  interaction with Y406 for several ns before exiting the pocket, and is then captured again later by it, but is not brought back into the pocket (**Supplementary Movie S1**). The full binding mechanism thus probably requires the rest of the substrate RNA, though the role of Y406 emerges already from the present simulations with the mononucleotide. The different conformations of Y406 reflect different steps in the binding and catalysis reaction. In the METTL3 SAM-bound state, the Y406 side chain is flexible. To bind RNA effectively, Y406 needs to stabilize through the water-bridged polar interaction with the backbone of W398. The substrate adenosine can then bind through selection of this conformation. Additionally, Y406 plays a role in positioning the adenosine molecule at the catalytic site.

## SAM binding primes the METTL3 active site for adenosine recognition

Given the high concentration of SAM in the cell (60 to 160  $\mu$ M in the rat liver), the cosubstrate SAM is expected to bind before the RNA substrate.<sup>46</sup> SAM binding results in large conformational changes of several side chains in the METTL3 active site (**Figure 6**). In the apo state, the K513 side chain points away from the putative RNA binding site and is involved in intramolecular polar contacts with the side chains of Y518, E532, and, via a water molecule, D395, that probably help to stabilize the apo protein (**Figure 6A**). SAM binding disrupts the water-mediated hydrogen bond between K513 and D395. This leads to a conformational change in the K513 side chain which then points in the direction of the RNA binding site where it can form a direct salt bridge with D395. In that conformation, the K513 side chain can also readily form a hydrogen bond to the adenosine-N<sup>7</sup> of an RNA substrate as seen in the BA2 structure (**Figure 6B**). Furthermore, the side chain of H512, which is part of the SAM binding pocket environment, also undergoes a conformational change upon SAM binding (see **Figure 6A**). In the apo state, the H512 side chain points inwards and thus blocks the adenosine binding site. However, once SAM is bound, the H512 side chain is attracted to form a  $\pi$ - $\pi$  interaction with H538 whose side chain rotates from the apo state to interact with SAM in the holo state. This conformational change of the H512 side chain makes space for adenosine to bind as seen in the BA2 structure (see **Figure 6B**). MD simulations show that the side chains of both H512 and K513 are flexible in the apo state, but undergo stabilization upon SAM and adenosine binding (**Supplementary Figures S8, S11, and S12**). Together, the conformational switches of the H512 and K513 side chains upon SAM binding can be seen as priming METTL3 for adenosine binding.






**Figure 5.**

### Flexibility of Y406 is evidenced by MD simulations.

**(A)** Distance time series of five 500-ns MD trajectories started from the BA2 conformation with substrates (top), products (middle), or apo METTL3-14 (bottom). The distance between the Y406 side chain and the backbone O of W398 (grey trace) reports on the orientation of Y406 and the flexibility of the loop. The data points are coloured black if AMP/ $m^6$ AMP is bound, and grey if not. Bound AMP is defined by a distance of less than 6 Å between N<sup>6</sup> of adenosine and Cy of D395. **(B)** The conformations of the flexible METTL3 backbone (ribbon) and Y406 side chain (sticks) are shown at different timepoints between 0-300 ns of the simulation with SAM and AMP, the Y406 side chain is coloured from red/start to blue/end. SAM and AMP are shown as sticks at the start of the simulation.



**Figure 6.**

### SAM binding primes the METTL3 active site for adenosine binding.

**(A)** Structural overlay of the METTL3 apo (grey) and SAM bound holo state (cyan). Black/cyan dashes indicate intramolecular polar contacts in apo/olo METTL3. Residues are shown as sticks and labelled. SAM and METTL3 residues in the holo state are shown as transparent sticks. Waters are shown as red spheres. **(B)** Structural overlay of the METTL3 apo (grey) and BA2 bound state (magenta). Black dashes indicate intramolecular polar contacts in apo METTL3, magenta dashes indicate polar contacts in BA2 bound METTL3. Residues are shown as sticks and labelled. BA2 is shown in magenta as sticks, its SAM and adenosine moieties are indicated. METTL3 residues in the apo state are shown as transparent sticks. Waters are shown as red spheres.



## BA2 represents a transition state analogue of the METTL3 catalysed reaction

We compared the structure of METTL3-BA2 with the structures of RNA MTases METTL4 and METTL16 bound to their substrates (**Figure 7**). When the SAM moiety of BA2 is superimposed with SAM bound to METTL4 and METTL16, the adenosine moiety of BA2 is situated in a very similar position as the substrates of the other MTases (**Figure 7A**). In METTL4 and METTL16, the N<sup>6</sup>-atom of their substrate adenosine is positioned at a distance of 2.8 and 2.1 Å from the methyl group of SAM, respectively (**Figure 7B,C**). The angle formed between SAM-5'S – Cε – adenosine-N<sup>6</sup> in the substrate-cosubstrate pairs of METTL4 and METTL16 is 162.5° and 175.4°, respectively. In BA2, the adenosine-N<sup>6</sup> is situated, through the alkyl chain of the linker, at 1.9 Å away from the carbon corresponding to the Cε-atom of the methionine moiety in the SAM cofactor (**Figure 7D**). The 1.9 Å distance in the crystal structure with BA2 is similar to the corresponding distance in METTL4 and METTL16, which, in two non-linked moieties, would allow for an SN<sub>2</sub> transfer of the methyl group of a METTL3-bound SAM to the N<sup>6</sup>-atom of an adenosine substrate. Strikingly, the angle formed by the SAM-5'S – Cε – adenosine-N<sup>6</sup> in METTL3 is 160.9° which is similar to the other MTases, and very close to the optimal 180° for an SN<sub>2</sub> reaction in which the adenosine-N<sup>6</sup> attacks the SAM-Cε and SAH becomes the leaving group. Hence, the adenosine of BA2 is in a suitable orientation for methyl transfer and BA2 represents a transition state mimic for METTL3. This is useful atomistic information for setting up QM/MM free energy calculations to study the catalytic reaction (see below).


## The METTL3 catalytic pocket supports direct methyl transfer without prior deprotonation

We carried out hybrid quantum mechanical/molecular mechanical (QM/MM<sup>47–54</sup>) free energy simulations to establish the catalytic mechanism of RNA methylation by the METTL3-14 complex.<sup>22,55–56</sup> The crystal structure with the bisubstrate analogue BA2 (see **Figure 3C**) was used as the starting point for the QM/MM simulations (**Figure 8**).

In the simplest mechanism (**Figure 8A**), the methyl cation in the SAM cofactor is transferred directly to the N<sup>6</sup> position of the adenosine substrate, prior to the deprotonation of N<sup>6</sup>H<sub>2</sub>, which has a very high pK<sub>a</sub> of ~17.<sup>57</sup> Indeed, DFTB3/MM free energy simulations show that this mechanism is energetically favourable (by about ~4 kcal/mol) with a barrier of 15-16 kcal/mol (**Figure 8B** and **Supplementary Figure S13 and S14**). The catalytic turnover of METTL3-14 as measured by an enzymatic RNA methylation assay is 0.2-0.6 min<sup>-1</sup> at ambient temperature which implies a barrier of ~20 kcal/mol.<sup>58–60</sup> Hence, the methyl transfer is not the rate-limiting step. Taken together, the QM/MM and MD simulations suggest that the dissociation of the coproduct SAH and product RNA is likely the rate-limiting step.

Compared to the model reaction in solution computed using a continuum solvation model (**Table 3**), the reaction in the enzyme is substantially more exoergic, suggesting that the enzyme environment stabilizes the product of the methyl transfer reaction. Inspection of the active site structure based on DFTB3/MM simulations suggests that such stabilization primarily comes from the hydrogen-bonding interactions between the adenosine N<sup>6</sup> group and nearby polar groups, in particular the side chain of D395 and the backbone carbonyl of P396 (**Figure 8C**).


This catalytic mechanism is similar to N<sup>6</sup>-adenine DNA methyl transferase M-TaqI, in which the adenosine N<sup>6</sup> group is hydrogen-bonded to an Asn side chain and backbone carbonyl of a Pro in the protein.<sup>61</sup> In the absence of any catalytic base, it was proposed that the methyl transfer occurs first, leading to an m<sup>6</sup>NH<sub>2</sub><sup>+</sup> group well stabilized by hydrogen bonding interactions with the Asn side chain and Pro backbone carbonyl. The mechanism was supported by QM/MM free energy simulations with a barrier height of ~20 kcal/mol.<sup>62</sup> On the other hand, the same QM/MM study suggested that when the active site Asn was replaced by an Asp, a mechanism in which



**Figure 7.**

**The crystal structure of the complex of METTL3-14 with the bisubstrate analogue BA2 represents the transition state of catalysis.**

(A) Overlap of the crystal structure of the complex of METTL3-14 (grey) with BA2 (magenta) to the complex with SAM (cyan) and the substrate-cosubstrate pairs of METTL4 (green) and METTL16 (orange). The overlap was generated by aligning SAM from each (co)substrate pair to the SAM of METTL3. (B-D) Measurements of distances and angles between the adenosine-N<sup>6</sup> and SAM-CH<sub>3</sub> groups in the respective (co)substrate pairs shown in (A). (B) The METTL4 (co)substrate pair was generated by aligning the structure of METTL4-AM to METTL4-SAM. (C) The METTL16 (co)substrate pair was generated by aligning the structure of METTL16-MAT2A 3'UTR hairpin 1 to METTL16-SAH. SAM was then generated from SAH using the Chem3D software. (D) The METTL3 overlay was generated by aligning the structure of METTL3-BA2 to METTL3-SAM. The distance was measured between the N<sup>6</sup> of BA2 and C $\epsilon$  of SAM, the angle was measured between the N<sup>6</sup> of BA2, C $\epsilon$  of SAM, and 5'S of SAM.



**Figure 8.**

### Methyl transfer catalysed by METTL3 without prior deprotonation of adenosine is energetically favourable based on DFTB3/MM simulations.

**(A)** Proposed mechanism of the methyl transfer reaction catalysed by METTL3. **(B)** Potential of mean force (PMF) along the antisymmetric stretch coordinate that describes the methyl transfer between the N<sup>6</sup> in adenosine and the SAM sulphur atom computed using multiple walker metadynamics simulations. The solid curve represents the PMF averaged over three independent runs, and the shaded area represents the standard error of the mean of the three replicas. **(C)** Snapshots of the active site for the reactant (left panel), transition state (middle panel), and product windows (right panel). Key distances (in Å) involving the reactive groups and the nearby ion-pair (D395-K513) and P396 backbone carbonyl are shown. METTL3 backbone is shown in grey ribbon representation with side chains shown as sticks and labelled. SAM/adenosine and SAH/m<sup>6</sup>(NH<sub>2</sub><sup>+</sup>)-adenosine are shown as sticks and labelled in the reactant and product window, respectively, together with the transferred methyl group. The methylium group (CH<sub>3</sub><sup>+</sup>) is indicated in the transition state window.

Reaction <sup>b</sup>	Different computational models						
	B3LYP-D3	ωB97X-D	B3LYP-D3	ωB97X-D	DFTB3/3O-B	B3LYP-D3	ωB97XD
	aug-cc-pVDZ		aug-cc-pVTZ			CPCM <sup>c</sup> , aug-cc-pVTZ	
SAM + Ade → SAH + Ade-CH <sub>3</sub>	5.3	5.0	7.3	6.9	2.5	10.2	9.6
SAM + dp-Ade → SAH + dp-Ade-CH <sub>3</sub>	-129.1	-130.6	-128.1	-129.8	-140.5	-37.7	-38.9
Difference	-134.5	-135.6	-135.4	-136.8	-143.0	-47.9	-48.5

- a. Different density functional theory methods (B3LYP-D3, ωB97XD) and basis set combinations (aug-cc-pVDZ and aug-cc-pVTZ), along with a semi-empirical quantum mechanical method, DFTB3/3OB, are used to evaluate the intrinsic energetics for model methyl transfer reactions. The comparison also helps validate the more approximate DFTB3/3OB method, which is used in QM/MM free energy simulations.
- b. dp-Ade indicates a deprotonated adenosine at the N<sup>6</sup> position. SAM and SAH are modelled by replacing the adenosine and amino moieties by ethyl groups.
- c. A dielectric constant of 78.4 is used for the conductor-like polarizable continuum model (CPCM) to estimate the reaction energetics in solution. All other results are based on gas phase calculations.

**Table 3.**

**Reaction energetics (in kcal/mol) computed for model methyl transfer reactions that involve SAM and adenosine using different levels of theory.<sup>a</sup>**

deprotonation of the adenine N<sup>6</sup> group by the Asp preceded the methyl transfer from SAM also had a comparable free energy barrier. Due to the involvement of the proton transfer, the corresponding transition state exhibited rather different charge distributions from that in the WT M-TaqI. DNA N<sup>6</sup>-methyltransferases in the  $\alpha/\beta$  groups feature an Asp in the active site, while those in the  $\gamma$  group have an Asn at the equivalent position.<sup>63</sup> Hence, the QM/MM computational results suggest that transition states with distinct charge distributions are involved in different groups of enzymes (i.e.,  $\alpha/\beta$  vs.  $\gamma$ ), giving rise to the opportunity of designing transition state analogues as inhibitors unique to specific classes of methyl transferases.<sup>62</sup>


This raises the question whether methyl transfer in METTL3-14 may also occur following deprotonation of the adenosine N<sup>6</sup>H<sub>2</sub>. The structural features of the METTL3-14 active site do not support this mechanism. The N<sup>6</sup> position has a very high pK<sub>a</sub> of ~17, and thus its deprotonation requires a particularly strong base, which is absent in the active site of METTL3-14. For example, while there is a carboxylate nearby (D395), it forms a salt-bridge with K513, and therefore is expected to feature a too low pK<sub>a</sub> value to deprotonate the adenosine N<sup>6</sup>H<sub>2</sub>. These considerations are congruent with the observation that the computed DFTB3/MM free energy profile without N<sup>6</sup> deprotonation is possible within the reported experimental kinetics.<sup>58,64,65</sup>

The difference in the methylation energetics of adenosine in different protonation states shown in **Table 3** is consistent with the pK<sub>a</sub> difference of adenosine before and after methylation. Thus, the large difference suggests that N<sup>6</sup> becomes much more acidic following methylation, which is consistent with literature estimates of the pK<sub>a</sub> values of N<sup>6</sup>-protonated adenosine derivatives in the range of -3 to -10.<sup>66</sup> Close inspection of the active site structure in the product state of DFTB3/MM simulations reveals that the proton release may occur readily through water wires that connect the N<sup>6</sup> position to the protein-solvent interface (**Figure 9**). Therefore, favourable salt-bridges in the active site (e.g., D395-K513) do not have to break to allow the proton release following methylation of the substrate.

Finally, we note that the recently solved crystal structures of METTL4, which belongs to a subclade of MT-A70 family members of MTases, showed an active site very similar to that of METTL3.<sup>67</sup> A nearby ion-pair (D233-K364) is engaged in a hydrogen-bonding network involving both the substrate and SAM. Therefore, we expect that the catalytic mechanism discussed here applies also to METTL4 and potentially other MT-A70 family members of MTases.

## Complete atomistic model of METTL3 binding site plasticity and methyl transfer mechanism

The complementarity of the methodologies and the congruence of the experimental data and simulation results allow us to construct a model of the METTL3 catalytic reaction (**Figure 10**). This model shows that in the apo state (**Figure 10**, **State 0**), the side chain of K513 is involved in intramolecular interactions that stabilize the protein. SAM binding displaces the K513 side chain and brings it in the right conformation where it can form a hydrogen bond to the N<sup>7</sup> of the adenosine substrate (**Figure 10**, **State 1**). Space for adenosine to bind is further conditioned by the conformational change of the H512 side chain which blocks the catalytic site in the apo state, but is drawn to interact with H538 in the SAM-bound state. The proper recognition of adenosine is conditioned by the interaction with the aromatic side chain of Y406 to which the adenosine substrate can bind and stabilize the ASL1 loop through interaction in the BA4 conformation (**Figure 10**, **State 2**). The Y406 side chain acts as gatekeeper and swaps out to allow the adenosine to flip and slip into the catalytic site where it is stabilized through hydrogen bonds to E481 and K513 in the BA2 conformation (**Figure 10**, **State 3**). The bond between K513 and adenosine is especially stable in the MD simulations started from the binding mode of BA2. Hydrogen bonds with the D395 side chain and P396 backbone enhance the nucleophilicity of the adenosine-N<sup>6</sup> and trigger the SN2 reaction with the electrophilic methyl group (**Figure 10**, **State 4**). The deprotonation of m<sup>6</sup>NH<sub>2</sub><sup>+</sup> following the methyl transfer may proceed along multiple water-



**Figure 9.**

**Deprotonation of  $m^6NH_2^+$  may occur readily through water wires that connect the  $N^6$  position to the protein-solvent interface.**

Shown is a snapshot of the product state from DFTB3/MM simulations illustrating that the deprotonation of  $N^6$  following the methyl transfer may proceed along multiple water-mediated pathways that lead to the protein/solvent interface. METTL3 backbone is shown in grey ribbon representation with side chains shown as sticks and labelled, water molecules are shown as spheres. SAH and  $m^6(NH_2^+)$ -adenosine are shown as sticks and labelled together with the transferred methyl group. Hydrogen bonds are indicated with dotted lines. The movement of protons through water channels is indicated with arrows.



mediated pathways that lead to the protein-solvent-interface (**Figure 10**, [State 5](#)). The m<sup>6</sup>A product can then flip and slip back into the BA4 conformation (**Figure 10**, [State 6](#)) before the flexibility of the ASL1 loop then facilitates its release (**Figure 10**, [State 7](#)).

## Conclusions

We have conducted a combined experimental and computational study of METTL3-14 to characterize the methyl transfer reaction from SAM to the N<sup>6</sup> of adenosine in RNA. Crystal structures of METTL3-14 complexed with bisubstrate analogues were solved and allowed us to carry out classical MD simulations and QM/MM free-energy calculations. The encounter complex between the METTL3-SAM holo complex and RNA is promoted by electrostatic steering of the negatively charged RNA backbone by a surface groove of positively charged electrostatic potential centred around the sulfonium ion of SAM. The crystal structures together with mutational analysis and MD simulations have revealed a key role of Y406 in recruiting the adenosine of the RNA substrate into a catalytically competent position and orientation. The proper recognition of adenosine is conditioned by interaction with the aromatic side chain of Y406 positioned in the flexible active site loop of METTL3. The side chain of Y406 recruits the RNA-adenosine, accommodates it into the catalytic site, and facilitates release after methyl transfer. The adenosine ring forms hydrogen bonds to the side chains of E481 and K513 in the catalytic site. Alanine mutants of these newly identified adenosine binding residues show abolished MTase activity compared to the wild-type METTL3-14. Importantly, these mutants are still folded and able to bind the SAH cofactor. This confirms their contribution to RNA-substrate binding, as shown also by the stability of their hydrogen bonds to adenosine in the MD simulations. Hydrogen bonds with the D395 side chain and P396 backbone enhance the nucleophilicity of the adenosine-N<sup>6</sup> and trigger the SN<sub>2</sub> reaction with the electrophilic methyl group of SAM. The QM/MM calculations provide evidence that the transfer of the methyl group from SAM to adenosine proceeds without previous deprotonation of the adenosine-N<sup>6</sup>. Furthermore, the height of the QM/MM free-energy barrier indicates that the methyl transfer step is not rate-determining. MD simulations suggest that the release of the coproduct SAH is the rate-limiting step.

In conclusion, the present study provides evidence that bisubstrate analogues and multiscale atomistic simulations can be used to decipher RNA recognition by human RNA MTases. The multidisciplinary strategy described here can be used for other m<sup>6</sup>A MTases to probe their active site by adapting the RNA sequence of the BA to the RNA substrate of the MTase of interest.


## Materials and Methods

### Chemical synthesis of bisubstrate analogues

The synthesis of the bisubstrate analogues was as previously described: BA1/2/3/4/6, ref. [35](#); Compound 12, ref. [36](#); GA\*, ref. [37](#).

### METTL3-14 expression, purification, and site-directed mutagenesis

For determining the half maximal inhibitory concentration (IC<sub>50</sub>) with the full-length complex and for crystallization studies with the truncated complex METTL3<sup>MTD</sup>:METTL14<sup>MTD</sup> containing just the methyltransferase domains (MTD) of METTL3 (residues 354–580) METTL14 (residues 107–395), the recombinant complex constructs were expressed using the baculovirus/Sf9 insect cell expression system and purified as described previously.[22](#)



**Figure 10.**

**The experimental and computational data elucidate the individual steps of substrate binding, product release, and methyl transfer catalysed by METTL3.**

Schematic illustration of the individual steps making up the (co)substrate binding, methyl transfer, and (co)product release mechanism of METTL3; dashed grey and black lines indicate polar contacts that are intramolecular in METTL3 (including water mediated (waters shown as red spheres)) and intermolecular to the substrate adenosine, respectively. The ASL1 loop containing Y406 is either flexible (grey colour) or stabilized by the interaction with the substrate adenosine (black colour). Step 1: binding of SAM and flexibility of Y406 (supporting evidence from crystal structures and MD simulations of apo and SAM bound states); Step 2: substrate recognition (crystal structure of the complex with BA4); Step 3: flip and slip of the substrate adenosine into the catalytic site (crystal structure of the complex with BA2); Steps 4 and 5: methyl transfer and deprotonation of m<sup>6</sup>NH<sub>2</sub><sup>+</sup> through water channels (movement of protons indicated by arrows) (QM/MM free energy calculations); Steps 6 and 7: slip and flip of the product m<sup>6</sup>A out of the catalytic site and subsequent dissociation from METTL3; Step 8: SAH release to re-establish the apo state (MD simulations).

For mutational analysis, alanine mutants were generated in the baculovirus vector pFastBacDual-StrepII-GFP-TEV-METTL3-His-TEV-METTL14 using the QuikChange site-directed mutagenesis protocol and confirmed by sequencing. Recombinant baculoviruses were generated using the Bac-to-Bac system. For protein expression, suspension cultures of Sf9 cells in Sf-90 II SFM medium (Thermo Fisher) were infected at a density of  $2 \times 10^6 \text{ ml}^{-1}$ . Cells were harvested 72 hours post infection, resuspended in Buffer A (50 mM Tris-HCl pH 8.0, 500 mM NaCl) supplemented with Protease Inhibitor Cocktail (Roche Diagnostics GmbH, Germany), phenylmethylsulfonyl fluoride (PMSF), Salt Active Nuclease (Merck), and lysed by sonication. The protein complex was purified by Ni-affinity chromatography on a 5 mL HisTrap HP column (Cytiva) equilibrated and washed with Buffer A and eluted with 250 mM imidazole. Proteins were further purified by Strep-tag purification using a 5 mL StrepTrap XT column (Cytiva) equilibrated and washed with buffer A and eluted with 50 mM biotin. The affinity tags were removed by digestion with TEV protease overnight at 4°C, followed by further purification by size exclusion chromatography using a Superdex 200 Increase 10/300 GL column (Cytiva) in 20 mM Tris-Cl, pH 8.0, and 200 mM KCl. The proteins were concentrated, flash-frozen in liquid nitrogen, and stored at  $-80^\circ\text{C}$  until further use.

## Protein crystallization

The SAH (S-adenosyl-L-homocysteine)-bound holo protein crystals of METTL3<sup>MTD</sup>:METTL14<sup>MTD</sup> were obtained as previously described.<sup>22</sup> The bisubstrate analogues (BAs) were dissolved in DMSO at concentrations of 100 mM. Complex structures were solved by soaking BAs into holo protein crystals and replacing the bound SAH in the METTL3 catalytic pocket. First, 1  $\mu\text{L}$  of the BA dissolved in DMSO was left overnight to evaporate the solvent at room temperature. The next day, 1  $\mu\text{L}$  of mother liquor containing 30% PEG-3350 and 200 mM Mg-acetate was added on top of the dried compound stamp. One holo crystal was then transferred into the mother liquor over the target compound stamp. After 16 h of incubation at 22°C, the crystals were harvested and flash-frozen in liquid nitrogen.

## Data collection and structure solution

Diffraction data were collected at the PXIII beamline at the Swiss Light Source (SLS) of the Paul Scherrer Institute (PSI, Villigen, Switzerland). Data were processed using XDS.<sup>68</sup> The crystal structures were solved by molecular replacement by employing the 5L6D structure as the search model in the Phaser program (Phenix package).<sup>69</sup> Crystallographic models were constructed through iterative cycles of manual model building with COOT and refinement with Phenix.refine.<sup>70–73</sup>

## Reader-based TR-FRET assay

The inhibitory potencies of the bisubstrate analogues (BAs) for METTL3 were quantified by a homogeneous time resolved fluorescence (HTRF)-based enzyme assay as previously described.<sup>41</sup> Briefly, the level of  $\text{m}^6\text{A}$  in an RNA substrate after the reaction catalysed by METTL3-14 was quantified by measuring specific binding to the  $\text{m}^6\text{A}$  reader domain of YTHDC1 (residues 345-509) by HTRF. BAs that inhibit METTL3 decrease the  $\text{m}^6\text{A}$  level and thus reduce the HTRF signal. Dose response curves of titrations with the BAs were plotted in OriginLab 2018 and fitted with nonlinear regression “log(inhibitor) vs. normalized response with variable slope” from which  $\text{IC}_{50}$  values were determined.

For the mutational analysis, the HTRF assay was used with some modifications. In the reaction step, METTL3-14 (wildtype or mutant) (40 nM final concentration) methylates the 5'-biotinylated ssRNA (5'-AAGAACCGGACUAAGCU-3' (Microsynth)) (200 nM final concentration). The co-substrate SAM (Cisbio, 62SAHZLD) (450 nM final concentration) was added as the last component and thus initiated the methylation reaction. The final reaction volume was 15  $\mu\text{L}$  in 20 mM Tris-HCl, pH 7.5, 0.01% (w/v) bovine serum albumin (BSA). The reaction was let to incubate for 1 hour at room temperature (RT) and then stopped by addition of 5  $\mu\text{L}$  detection buffer (50 mM HEPES, pH 7.5, 150

mM NaCl, 200 mM KF, 0.05% (w/v) BSA, 25 nM GST-tagged m<sup>6</sup>A reader YTHDC1(345-509), 3 nM XL665-conjugated streptavidin (Cisbio, 610SAXLB), 1x anti-GST Eu<sup>3+</sup>-labelled antibody (from 400x stock (Cisbio, 61GSTKLB))). Capture of the m<sup>6</sup>A-modified RNA by the m<sup>6</sup>A reader and the biotinylated RNA by Streptavidin was allowed to proceed for 3 hours at RT and in the dark before the HTRF signal was measured using a Tecan Spark plate reader (Tecan). The plate reader recorded with a delay of 100 μs the emission at 620 and 665 nm after the excitation of the HTRF donor with UV light at 320 nm. The emission signal was read over an integration time of 400 μs. The  $\Delta F$  ( $((\text{ratio}_{\text{sample}} - \text{ratio}_{\text{background}}) / \text{ratio}_{\text{background}}) * 100$ ) used as background the reaction without SAM as an internal control.

## Differential scanning fluorimetry (thermal shift assay, TSA)

Experiments were conducted as previously described with some modifications.<sup>16</sup> Briefly, METTL3-14 (wildtype or mutants) at a final concentration of 0.5 μM was mixed with SAH at a final concentrations of 500 μM in a final volume of 20 μL in a buffer consisting of 20 mM Tris-Cl, pH 8.0, and 200 mM KCl. DMSO concentration was kept at 1 % (v/v). SYPRO Orange was added at a final dilution of 1 : 1000 (v/v) as a fluorescence probe (ex/em 465/590 nm). Differential scanning fluorimetry was performed on a LightCycler® 480 Instrument II (Roche Diagnostics, Indianapolis, IN). The temperature was raised in steps of 3.6°C per minute from 20°C to 85°C and fluorescence readings were taken at each interval. The T<sub>m</sub> values were determined as the transition midpoints of the individual samples. The  $\Delta T_m$  values were calculated as the difference between the transition midpoints of the individual samples and the reference wells containing the protein and DMSO only from the same plate. Samples were measured in triplicates.

## Molecular Dynamics simulations

The crystal structures of the METTL3-14 heterodimer in complex with the BA2 and BA4 ligands were used as starting conformation for simulating the heterodimer with its substrates SAM and AMP and products SAH and m<sup>6</sup>AMP. The addition of the phosphate to adenosine and m<sup>6</sup>-adenosine aims to mimic one element of the substrate RNA chain. The ligands were aligned to their respective moiety of the BA, keeping the original coordinates for present atoms, and reconstructing the missing parts. The missing segments of the METTL3-14 crystal structures were reconstructed using the SWISSMODEL web server with the structures as templates. The simulated construct spans residues L354 to L580 of METTL3 and S104 to L289 of METTL14. All MD simulations were performed with GROMACS 2021.5 using the CHARMM36m July 2021 force field. The models were solvated in a 9 nm box and equilibrated with Na<sup>+</sup> and Cl<sup>-</sup> ions to a concentration of 150 mM. Energy minimization was applied and a canonical equilibration under all-atom positional restraints was performed for 5 ns to reach 300 K. A further canonical equilibration was performed for 10 ns with the Cα atoms of the proteins, the adenine moiety of SAM/SAH and of AMP/m<sup>6</sup>AMP under positional restraints. These partial restraints were set to allow a relaxation of the side chains around the ligands. Sixteen independent runs of canonical MD simulations were then started for each of the systems, sampling 500 ns per run.

The existence and behaviour of salt bridges observed in the crystals were monitored throughout the runs. Furthermore, the stability of the binding was calculated by defining a binding pocket for the BA2 and the BA4 conformations. The distance between residues was calculated and contacts were defined as a distance smaller than 5 Å. For each of the simulation groups, the binding pocket was determined as the residues with a mean contact presence higher than the 90 percentiles of mean contacts. A dissociation event was defined as the mean of the ligand to pocket distances surpassing a threshold of 10 Å. The dissociation rates were predicted using exponential fitting, modelled as a single exponential. A single exponential with a multiplicative factor was used to check the quality of the fitting, with a preexponential factor close to 1 indicative of a good fit. No fitting was done for SAM/SAH as they remained bound in almost every single trajectory. The rates were calculated by fitting the dissociation time of two blocks of the trajectories at a time. The 16 choose 8 (12870) possible combinations were considered, and the mean of all the values was

reported as the dissociation rate in **Table 2**. The standard deviation was used as error. The unbinding rates using all the trajectories without block averaging are shown in **Supplementary Figure S5**.

## QM model calculations

To understand the intrinsic energetics of the methyl transfer reaction, we conducted QM calculations of an infinitely separated model substrate (adenosine) and a truncated model for the cofactor SAM in which the adenosine and amino moieties were replaced by ethyl groups. The N<sup>6</sup> position of the model substrate was taken to be either protonated (-NH<sub>2</sub>) or deprotonated. Calculations were performed in the gas phase using two different density functional theory (DFT) methods (B3LYP with the D3 dispersion correction<sup>74–77</sup> and  $\omega$ B97XD<sup>78</sup>) and two different basis sets (aug-cc-pVDZ and aug-cc-pVTZ).<sup>79–81</sup> Calculations were also conducted at the DFTB3/3OB level for calibration as the same QM method is used in subsequent QM/MM free energy simulations. To probe the effect of solvation on the methyl transfer energetics, single point calculations were carried out with the conductor-like polarizable continuum model (CPCM<sup>82,83</sup>) using the gas-phase optimized structures at both B3LYP-D3 and  $\omega$ B97XD levels. DFT calculations using B3LYP-D3 and  $\omega$ B97XD were conducted using the Gaussian16 software<sup>84</sup>, and DFTB3<sup>85</sup> calculations were carried out using the CHARMM program.<sup>86</sup>

## QM/MM free energy simulations

We employed QM/MM free energy simulations to probe the mechanism of adenosine N<sup>6</sup>-methylation catalysed by the METTL3-14 complex. As illustrated in **Figure 3C**, the crystal structure (at 2.3 Å resolution) of the complex with a transition state analogue (the bisubstrate analogue BA2) suggests a rather straightforward mechanism through which the methyl group is transferred from SAM to the adenosine N<sup>6</sup> position. The deprotonation of N<sup>6</sup> by a nearby base in principle may occur either before or after the methyl transfer, but QM/MM calculations strongly suggest that deprotonation occurs after the methyl transfer.

In the QM/MM simulations, the bisubstrate analogue was first converted to a SAM non-covalently bonded to the adenosine: the nitrogen N<sub>BS</sub> atom in the crystal structure was replaced by a sulphur atom and the extra carbon atom was deleted. The O5'-PA bond was cut off and a hydrogen atom was patched to the location. The QM region included the cofactor SAM, the model substrate adenosine (Ade), the carbonyl groups of A394, D395, P396, and P397 without the backbone carbonyl group, and the side chain of K513. Link atoms were added between C and C<sub>α</sub> of A394 and P397, and between C<sub>β</sub> and C<sub>α</sub> of K513 to saturate the valence of the QM boundary atoms using the divided frontier charge (DIV) scheme.<sup>87</sup> The QM atoms were treated with the third-order density functional tight binding (DFTB3) method with the 3OB parameter set<sup>88,89</sup>; benchmark calculations using model compounds (see **Table 3**) indicate that the DFTB3/3OB method describes the energetics of the methyl transfer reaction rather well in comparison to DFT calculations with a large basis set. The MM region was described with the CHARMM36 force field for proteins.<sup>90</sup>

In the generalized solvent boundary potential (GSBP<sup>91,92</sup>) framework, the inner region contained atoms within a 27 Å-radius sphere centred at the N<sup>6</sup> in Ade. Newtonian equations of motion were solved for atoms within 25 Å. Protein atoms in the buffer region (25–27 Å) were harmonically restrained with force constants determined from the crystallographic B factors and Langevin equations of motion were solved with a bath temperature of 300 K.<sup>93</sup> The remaining portion of the system in the outer region was frozen. All water molecules were subject to a weak geometrical (GEO) type of restraining potential to keep them inside the inner sphere.<sup>86</sup> Weak GEO restraints were added on adenosine to make sure it was well-bounded during the simulations. Electrostatic interactions among inner region atoms were treated with extended electrostatics and a group-based cut-off scheme.<sup>94</sup> The static field due to the outer region atoms was evaluated with the linearized Poisson-Boltzmann (PB) equations using a focusing scheme,



which employed a coarse grid of 1.2 Å and a fine grid of 0.4 Å.<sup>95</sup> The reaction field matrix was evaluated using spherical harmonics up to the 20th order. In the PB calculations, dielectric constants of the protein and water were set to 1 and 80, respectively, and the salt concentration was set to zero.

To probe the energetics of the methyl transfer reaction catalysed by the METTL3-14 complex, well-tempered multiple-walker metadynamics simulations<sup>96,97</sup> were carried out using the PLUMED-CHARMM interface.<sup>98,99</sup> The antisymmetric stretch that describes the methyl transfer process between the N<sup>6</sup> in adenosine and the SAM sulphur-methyl group was chosen as the collective variable (CV), i.e.,  $\xi = r(\text{CE-SD}) - r(\text{CE-N6})$ . The corresponding C-S distance and C-N distance are also monitored but no bias potential was added. SHAKE was applied to all bonds involving hydrogen and used to avoid undesired proton transfer reactions.<sup>100</sup>

The first two metadynamic runs were not well-tempered for the efficiency of sampling. In the subsequent well-tempered runs, the bias factor was set to be 35. A new Gaussian biasing potential was added every 0.2 ps with an initial height of 0.3 kJ/mol and a width of 0.05. Twenty-four walkers with different initial velocities were used per simulation in parallel while sharing hill history among all walkers every 1 ps. Each walker was run for 250 ps for a total of 6 ns of sampling, and convergence was evaluated by comparing the PMF as a function of the number of Gaussians added.

## Acknowledgements

This work was supported by a grant of the Swiss National Science Foundation to A.C. (grant number 310030-212195). The MD simulations were carried out at Eiger@Alps at the Swiss National Supercomputing Center (in Lugano, Switzerland). This work was further supported by a grant of the French Agence Nationale de la Recherche (ANR), Project ARNTools, to M.E.Q. (grant ARNtools-19-CE07-0028-01). The QM/MM study was supported by the NIH Grant R35-GM141930 to Q.C. Computational resources from the project BIO230101 allocated through ACCESS are greatly appreciated; part of the computational work was performed on the Shared Computing Cluster which is administered by Boston University's Research Computing Services (URL: [www.bu.edu/tech/support/research/](http://www.bu.edu/tech/support/research/)).

## Author contributions

Conceptualization: MEQ, QC, AC

Investigation: IC, PAVR, RKB, JD, DC, EB, LI, YL, DH

Visualization: IC, PAVR, JD

Funding acquisition: MEQ, QC, AC

Project administration: AC

Supervision: MEQ, QC, AC

Writing – original draft: IC, PAVR, JD, QC, AC


Writing – review & editing: IC, PAVR, RKB, JD, MEQ, QC, AC



## Competing interests

None of the authors declare a competing interest.

## Data availability

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. The coordinates of the METTL3-14-bisubstrate analogue complexes have been deposited in the Protein Data Bank under accession numbers 8PW9 (BA1 complex), 8PW8 (BA2 complex), 8PWA (BA4 complex), 8PWB (BA6 complex). The MD simulation trajectories and the PLUMED file that was used for the DFTB3/MM metadynamics simulations (plumed.dat) are available via Zenodo under the link: <https://doi.org/10.5281/zenodo.10277884> .

## Supplementary Materials

Supplementary Figures S1 to S14

Supplementary Table S1

Supplementary Movie S1

Supplementary references

## References

- 1 Boccaletto P. *et al.* (2022) **MODOMICS: a database of RNA modification pathways. 2021 update** *Nucleic Acids Res* **50**:D231–D235
- 2 Roundtree I. A., Evans M. E., Pan T., He C (2017) **Dynamic RNA Modifications in Gene Expression Regulation** *Cell* **169**:1187–1200
- 3 Fu Y., Dominissini D., Rechavi G., He C (2014) **Gene expression regulation mediated through reversible m(6)A RNA methylation** *Nat Rev Genet* **15**:293–306
- 4 Linder B., Grozhik A. V., Olarerin-George A. O., Meydan C., Mason C. E., Jaffrey S. R (2015) **Single-nucleotide-resolution mapping of m6A and m6Am throughout the transcriptome** *Nat Methods* **12**:767–772
- 5 Ke S. *et al.* (2015) **A majority of m6A residues are in the last exons, allowing the potential for 3' UTR regulation** *Genes Dev* **29**:2037–2053
- 6 Ke S., Pandya-Jones A., Saito Y., Fak J., Vågbo C. B., Geula S., Hanna J. H., Black D. L., Darnell J. E., Darnell R. B. (2017) **m6A mRNA modifications are deposited in nascent pre-mRNA and are not required for splicing but do specify cytoplasmic turnover** *Genes & Development* **31**:990–1006
- 7 Lesbirel S., Wilson S. A (2019) **The m(6)A-methylase complex and mRNA export** *Biochim Biophys Acta Gene Regul Mech* **1862**:319–328
- 8 Lee Y., Choe J., Park O. H., Kim Y. K (2020) **Molecular Mechanisms Driving mRNA Degradation by m(6)A Modification** *Trends Genet* **36**:177–188
- 9 Kadumuri R. V., Janga S. C (2018) **Epitranscriptomic Code and Its Alterations in Human Disease** *Trends Mol Med* **24**:886–903
- 10 Liu J. *et al.* (2014) **A METTL3-METTL14 complex mediates mammalian nuclear RNA N6-adenosine methylation** *Nat Chem Biol* **10**:93–95
- 11 De Jesus D. F., Zhang Z., Kahraman S., Brown N. K., Chen M., Hu J., Gupta M. K., He C., Kulkarni R (2019) **N. m(6)A mRNA Methylation Regulates Human beta-Cell Biology in Physiological States and in Type 2 Diabetes** *Nat Metab* **1**:765–774
- 12 Dang W., Xie Y., Cao P., Xin S., Wang J., Li S., Li Y., Lu J (2019) **N(6)-Methyladenosine and Viral Infection** *Front Microbiol* **10**
- 13 Chen X. Y., Zhang J., Zhu J. S (2019) **The role of m(6)A RNA methylation in human cancer** *Mol Cancer* **18**
- 14 Barbieri I. *et al.* (2017) **Promoter-bound METTL3 maintains myeloid leukaemia by m(6)A-dependent translation control** *Nature* **552**:126–131
- 15 Vu L. P. *et al.* (2017) **The N(6)-methyladenosine (m(6)A)-forming enzyme METTL3 controls myeloid differentiation of normal hematopoietic and leukemia cells** *Nat Med* **23**:1369–1376

- 16 Moroz-Omori E. V., Huang D., Kumar Bedi R., Cheriyaunkunel S. J., Bochenkova E., Dolbois A., Rzeczkowski M. D., Li Y., Wiedmer L., Caflisch A (2021) **METTL3 Inhibitors for Epitranscriptomic Modulation of Cellular Processes** *ChemMedChem* **16**:3035–3043
- 17 Yankova E. *et al.* (2021) **Small-molecule inhibition of METTL3 as a strategy against myeloid leukaemia** *Nature* **593**:597–601
- 18 Zeng C., Huang W., Li Y., Weng H (2020) **Roles of METTL3 in cancer: mechanisms and therapeutic targeting** *J Hematol Oncol* **13**
- 19 Dolbois A., Bedi R. K., Bochenkova E., Muller A., Moroz-Omori E. V., Huang D. Z., Caflisch A. (2021) **1,4,9-Triazaspiro[5.5]undecan-2-one Derivatives as Potent and Selective METTL3 Inhibitors** *J Med Chem* **64**:12738–12760
- 20 Wang X. *et al.* (2016) **Structural basis of N(6)-adenosine methylation by the METTL3-METTL14 complex** *Nature* **534**:575–578
- 21 Wang P., Doxtader K. A., Nam Y (2016) **Structural Basis for Cooperative Function of Mettl3 and Mettl14 Methyltransferases** *Mol Cell* **63**:306–317
- 22 Sledz P., Jinek M (2016) **Structural insights into the molecular mechanism of the m6A writer complex** *eLife* **5**
- 23 Yoshida A., Oyoshi T., Suda A., Futaki S., Imanishi M (2022) **Recognition of G-quadruplex RNA by a crucial RNA methyltransferase component, METTL14** *Nucleic Acids Res* **50**:449–457
- 24 Huang J. *et al.* (2019) **Solution structure of the RNA recognition domain of METTL3-METTL14 N(6)-methyladenosine methyltransferase** *Protein Cell* **10**:272–284
- 25 Sun J., Shi F., Yang N (2019) **Exploration of the Substrate Preference of Lysine Methyltransferase SMYD3 by Molecular Dynamics Simulations** *ACS Omega* **4**:19573–19581
- 26 Sun J., Li Z., Yang N (2021) **Mechanism of the Conformational Change of the Protein Methyltransferase SMYD3: A Molecular Dynamics Simulation Study** *Int J Mol Sci* **22**
- 27 Singh A. P., Kumar R., Gupta D (2021) **Structural insights into the mechanism of human methyltransferase hPRMT4** *J Biomol Struct Dyn* :1–14
- 28 Singh W., Karabancheva-Christova T. G., Black G. W., Ainsley J., Dover L., Christov C. Z. (2016) **Conformational Dynamics, Ligand Binding and Effects of Mutations in NirE an S-Adenosyl-L-Methionine Dependent Methyltransferase** *Sci Rep* **6**
- 29 Ichiye T., Karplus M (1991) **Collective motions in proteins: A covariance analysis of atomic fluctuations in molecular dynamics and normal mode simulations. Proteins: Structure Function, and Bioinformatics** **11**:205–217
- 30 Chen S. *et al.* (2019) **The dynamic conformational landscape of the protein methyltransferase SETD8** *Elife* **8**
- 31 Chuang C. H., Chiou S. J., Cheng T. L., Wang Y. T (2018) **A molecular dynamics simulation study decodes the Zika virus NS5 methyltransferase bound to SAH and RNA analogue** *Sci Rep* **8**

- 32 Sk M. F., Jonniya N. A., Roy R., Poddar S., Kar P (2020) **Computational Investigation of Structural Dynamics of SARS-CoV-2 Methyltransferase-Stimulatory Factor Heterodimer nsp16/nsp10 Bound to the Cofactor SAM** *Front Mol Biosci* **7**
- 33 Doxtader K. A., Wang P., Scarborough A. M., Seo D., Conrad N. K., Nam Y (2018) **Structural Basis for Regulation of METTL16, an S-Adenosylmethionine Homeostasis Factor** *Mol Cell* **71**:1001–1011
- 34 Oerum S. *et al.* (2019) **Bisubstrate analogues as structural tools to investigate m(6)A methyltransferase active sites** *RNA Biol* **16**:798–808
- 35 Atdjian C., Iannazzo L., Braud E., Ethève-Quelquejeu M (2018) **Synthesis of SAM-Adenosine Conjugates for the Study of m6A-RNA Methyltransferases** *European Journal of Organic Chemistry* **2018**:4411–4425
- 36 Atdjian C., Coelho D., Iannazzo L., Ethève-Quelquejeu M., Braud E (2020) **Synthesis of Triazole-Linked SAM-Adenosine Conjugates: Functionalization of Adenosine at N-1 or N-6 Position without Protecting Groups** *Molecules* **25**
- 37 Meynier V., Iannazzo L., Catala M., Oerum S., Braud E., Atdjian C., Barraud P., Fonvielle M., Tisne C., Etheve-Quelquejeu M (2022) **Synthesis of RNA-cofactor conjugates and structural exploration of RNA recognition by an m6A RNA methyltransferase** *Nucleic Acids Res* **50**:5793–5806
- 38 Coelho D., Le Corre L., Bartosik K., Iannazzo L., Braud E., Etheve-Quelquejeu M (2023) **Synthesis of Bisubstrate Analogues for RNA Methylation Studies using two Transition-Metal-Catalyzed Reactions** *Chemistry*
- 39 Schapira M (2016) **Structural Chemistry of Human RNA Methyltransferases** *ACS Chem Biol* **11**:575–582
- 40 O'Hagan D., Schmidberger J. W (2010) **Enzymes that catalyse SN2 reaction mechanisms** *Nat Prod Rep* **27**:900–918
- 41 Wiedmer L., Eberle S. A., Bedi R. K., Sledz P., Caflisch A (2019) **A Reader-Based Assay for m(6)A Writers and Erasers** *Anal Chem* **91**:3078–3084
- 42 Laskowski R. A., Swindells M. B (2011) **LigPlot+: multiple ligand-protein interaction diagrams for drug discovery** *J Chem Inf Model* **51**:2778–2786
- 43 Bedi R. K., Huang D., Li Y., Caflisch A (2023) **Structure-Based Design of Inhibitors of the m6A-RNA Writer Enzyme METTL3** *ACS Bio & Med Chem Au*
- 44 Selberg S., Blokhina D., Aatonen M., Koivisto P., Siltanen A., Mervaala E., Kankuri E., Karelson M (2019) **Discovery of Small Molecules that Activate RNA Methylation through Cooperative Binding to the METTL3-14-WTAP Complex Active Site** *Cell Rep* **26**:3762–3771
- 45 Qi S., Mota J., Chan S. H., Villarreal J., Dai N., Arya S., Hromas R. A., Rao M. K., Correa I. R., Gupta Y. K (2022) **RNA binding to human METTL3-METTL14 restricts N(6)-deoxyadenosine methylation of DNA in vitro** *Elife* **11**
- 46 Finkelstein J. D., Martin J. J (1984) **Methionine metabolism in mammals. Distribution of homocysteine between competing pathways** *Journal of Biological Chemistry* **259**:9508–9513

- 47 Warshel A., Levitt M (1976) **Theoretical Studies of Enzymic Reactions - Dielectric, Electrostatic and Steric Stabilization of Carbonium-Ion in Reaction of Lysozyme** *J. Mol. Biol* **103**:227–249
- 48 Field M. J., Bash P. A., Karplus M (1990) **A Combined Quantum-Mechanical and Molecular Mechanical Potential for Molecular-Dynamics Simulations** *J. Comput. Chem* **11**:700–733
- 49 Senn H. M., Thiel W. (2009) **QM/MM methods for biomolecular systems** *Angew. Chem. Int. Ed.* **48**:1198–1229
- 50 Lipkowitz K. B., Boyd D. B. (1995) **Reviews in Computational Chemistry** :119–185
- 51 Brunk E., Rothlisberger U (2015) **Mixed Quantum Mechanical/Molecular Mechanical Molecular Dynamics Simulations of Biological Systems in Ground and Electronically Excited States** *Chem. Rev* **115**:6217–6263
- 52 Chung L. W. *et al.* (2015) **The ONIOM Method and Its Applications** *Chem. Rev.* **115**:5678–5769
- 53 Lu X., Fang D., Ito S., Okamoto Y., Ovchinnikov V., Cui Q (2016) **QM/MM Free Energy Simulations: Recent Progress and Challenges** *Mol. Simul* **42**:1056–1078
- 54 Hu H., Yang W. T (2008) **Free Energies of Chemical Reactions in Solution and in Enzymes with Ab Initio Quantum Mechanics/Molecular Mechanics Methods** *Annu. Rev. Phys. Chem* **59**:573–601
- 55 Wu B., Li L., Huang Y., Ma J., Min J (2017) **Readers, writers and erasers of N6-methylated adenosine modification** *Curr. Opin. Struc. Biol* **47**:67–76
- 56 Oerum S., Meynier V., Catala M., Tisne C (2021) **A comprehensive review of m6A/m6Am RNA methyltransferase structures** *Nuc. Acids Res* **49**:7239–7255
- 57 Lippert B (2005) **Alternations of nucleobase pKa values upon metal coordination: origins and consequences** *Prog. Inorg. Chem* **54**:385–447
- 58 Buker S. M., Gurard-Levin Z. A., Wheeler B. D., Scholle M. D., Case A. W., Hirsch J. L., Ribich S., Copeland R. A., Boriack-Sjodin P. A (2020) **A Mass Spectrometric Assay of METTL3/METTL14 Methyltransferase Activity** *SLAS Discov* **25**:361–371
- 59 Garcia-Viloca M., Gao J., Karplus M., Truhlar D. G (2004) **How Enzymes Work: Analysis by Modern Rate Theory and Computer Simulations** *Science* **303**:186–195
- 60 Glowacki D. R., Harvey J. N., Mulholland A. J (2012) **Taking Ockham's razor to enzyme dynamics and catalysis** *Nat Chem* **4**:169–176
- 61 Goedecke K., Pignot M., Goody R. S., Scheidig A. J., Weinhold E (2001) **Structure of the N6-adenine DNA methyltransferase M•TaqI in complex with DNA and a cofactor analog** *Nature Structural Biology* **8**:121–125
- 62 Aranda J., Zinovjev K., Roca M., Tunon I (2014) **Dynamics and reactivity in Thermus aquaticus N6-adenine methyltransferase** *J Am Chem Soc* **136**:16227–16239
- 63 Malone T., Blumenthal R. M., Cheng X (1995) **Structure-guided Analysis Reveals Nine Sequence Motifs Conserved among DNA Amino-methyl-transferases, and Suggests a Catalytic Mechanism for these Enzymes** *Journal of Molecular Biology* **253**:618–632

- 64 Woodcock C. B. *et al.* (2019) **Human MettL3-MettL14 complex is a sequence-specific DNA adenine methyltransferase active on single-strand and unpaired DNA in vitro** *Cell Discov* **5**
- 65 Xiao S. *et al.* (2022) **High-Throughput-Methyl-Reading (HTMR) assay: a solution based on nucleotide methyl-binding proteins enables large-scale screening for DNA/RNA methyltransferases and demethylases** *Nucleic Acids Res* **50**
- 66 Kettani A., Guéron M., Leroy J. L (1997) **Amino Proton Exchange Processes in Mononucleosides** *Journal of the American Chemical Society* **119**:1108–1115
- 67 Luo Q. *et al.* (2022) **Structural insights into molecular mechanism for N(6)-adenosine methylation by MT-A70 family methyltransferase METTL4** *Nat Commun* **13**
- 68 Kabsch W. Xds. (2010) **Kabsch, W. Xds. Acta Crystallogr D Biol Crystallogr** **66**, 125–132 (2010). *Acta Crystallogr D Biol Crystallogr* **66**:125–132
- 69 McCoy A. J., Grosse-Kunstleve R. W., Adams P. D., Winn M. D., Storoni L. C., Read R. J. (2007) **Phaser crystallographic software** *J Appl Crystallogr* **40**:658–674
- 70 Emsley P., Lohkamp B., Scott W. G., Cowtan K. (2010) **Features and development of Coot** *Acta Crystallogr D Biol Crystallogr* **66**:486–501
- 71 Afonine P. V., Grosse-Kunstleve R. W., Echols N., Headd J. J., Moriarty N. W., Mustyakimov M., Terwilliger T. C., Urzhumtsev A., Zwart P. H., Adams P. D (2012) **Towards automated crystallographic structure refinement with phenix.refine** *Acta Crystallogr D Biol Crystallogr* **68**:352–367
- 72 Liebschner D. *et al.* (2019) **Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix** *Acta Crystallogr D Struct Biol* **75**:861–877
- 73 Emsley P., Cowtan K (2004) **Coot: model-building tools for molecular graphics** *Acta Crystallogr D Biol Crystallogr* **60**:2126–2132
- 74 Becke A. D (1988) **Density-functional exchange-energy approximation with correct asymptotic behavior** *Physical Review A* **38**:3098–3100
- 75 Becke A. D. (1993) **Density-functional thermochemistry. III. The role of exact exchange** *The Journal of Chemical Physics* **98**:5648–5652
- 76 Lee C., Yang W., Parr R. G (1988) **Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density** *Physical Review B* **37**:785–789
- 77 Grimme S., Antony J., Ehrlich S., Krieg H (2010) **A consistent and accurate ab initio parametrization of density functional dispersion correction (DFT-D) for the 94 elements H-Pu** *The Journal of Chemical Physics* **132**
- 78 Chai J.-D., Head-Gordon M (2008) **Systematic optimization of long-range corrected hybrid density functionals** *The Journal of Chemical Physics* **128**
- 79 Dunning T. H. (1989) **Gaussian basis sets for use in correlated molecular calculations I. The atoms boron through neon and hydrogen. The Journal of Chemical Physics** **90**:1007–1023



- 80 Kendall R. A., Dunning T. H., Harrison R. J (1992) **Electron affinities of the first-row atoms revisited. Systematic basis sets and wave functions** *The Journal of Chemical Physics* **96**:6796–6806
- 81 Woon D. E., Dunning T. H. (1993) **Gaussian basis sets for use in correlated molecular calculations III. The atoms aluminum through argon.** *The Journal of Chemical Physics* **98**:1358–1371
- 82 Barone V., Cossi M (1998) **Quantum Calculation of Molecular Energies and Energy Gradients in Solution by a Conductor Solvent Model** *The Journal of Physical Chemistry A* **102**:1995–2001
- 83 Cossi M., Rega N., Scalmani G., Barone V (2003) **Energies, structures, and electronic properties of molecules in solution with the C-PCM solvation model** *Journal of Computational Chemistry* **24**:669–681
- 84 Gaussian 16 Rev. C.01 (2016) **Gaussian 16 Rev. C.01 (Wallingford, CT, 2016).**
- 85 Gaus M., Cui Q., Elstner M (2011) **DFTB3: Extension of the Self-Consistent-Charge Density-Functional Tight-Binding Method (SCC-DFTB)** *Journal of Chemical Theory and Computation* **7**:931–948
- 86 Brooks B. R. *et al.* (2009) **CHARMM: The biomolecular simulation program** *Journal of Computational Chemistry* **30**:1545–1614
- 87 König P. H., Hoffmann M., Frauenheim T., Cui Q (2005) **A Critical Evaluation of Different QM/MM Frontier Treatments with SCC-DFTB as the QM Method** *The Journal of Physical Chemistry B* **109**:9082–9095
- 88 Gaus M., Goez A., Elstner M (2013) **Parametrization and Benchmark of DFTB3 for Organic Molecules** *Journal of Chemical Theory and Computation* **9**:338–354
- 89 Gaus M., Lu X., Elstner M., Cui Q (2014) **Parameterization of DFTB3/3OB for Sulfur and Phosphorus for chemical and biological applications** *J. Chem. Theory Comput* **10**:1518–1537
- 90 Huang J., MacKerell Jr A. D. (2013) **CHARMM36 all-atom additive protein force field: Validation based on comparison to NMR data** *J. Comput. Chem* **34**:2135–2145
- 91 Im W., Berneche S., Roux B (2001) **Generalized solvent boundary potential for computer simulations** *J. Chem. Phys* **114**:2924–2937
- 92 Schaefer P., Riccardi D., Cui Q (2005) **Reliable treatment of electrostatics in combined QM/MM simulation of macromolecules** *J. Chem. Phys* **123**
- 93 Brooks C. L., Karplus M. (1983) **Deformable stochastic boundaries in molecular dynamics** *The Journal of Chemical Physics* **79**:6312–6325
- 94 Stote R. H., States D. J., Karplus M (1991) **On the Treatment of Electrostatic Interactions in Biomolecular Simulations** *AIP Conference Proceedings* **239**:117–117
- 95 Im W., Beglov D., Roux B (1998) **Continuum solvation model: Computation of electrostatic forces from numerical solutions to the Poisson-Boltzmann equation** *Computer Physics Communications* **111**:59–75

- 96 Valsson O., Tiwary P., Parrinello M (2016) **Enhancing Important Fluctuations: Rare Events and Metadynamics from a Conceptual Viewpoint** *Annual Review of Physical Chemistry* **67**:159–184
- 97 Barducci A., Bussi G., Parrinello M (2008) **Well-Tempered Metadynamics: A Smoothly Converging and Tunable Free-Energy Method** *Physical Review Letters* **100**
- 98 Bonomi M. *et al.* (2009) **PLUMED: A portable plugin for free-energy calculations with molecular dynamics** *Computer Physics Communications* **180**:1961–1972
- 99 Bonomi M. *et al.* (2019) **The, P. c. Promoting transparency and reproducibility in enhanced molecular simulations** *Nature Methods* **16**:670–673
- 100 Ryckaert J.-P., Ciccotti G., Berendsen H. J. C (1977) **Numerical integration of the cartesian equations of motion of a system with constraints: molecular dynamics of n-alkanes** *Journal of Computational Physics* **23**:327–341

## Article and author information

### Ivan Corbeski

Department of Biochemistry, University of Zurich, Zurich CH-8057, Switzerland  
ORCID iD: [0000-0002-5881-8425](https://orcid.org/0000-0002-5881-8425)

### Pablo Andrés Vargas-Rosales

Department of Biochemistry, University of Zurich, Zurich CH-8057, Switzerland  
ORCID iD: [0000-0001-5198-620X](https://orcid.org/0000-0001-5198-620X)

### Rajiv Kumar Bedi

Department of Biochemistry, University of Zurich, Zurich CH-8057, Switzerland  
ORCID iD: [0000-0002-8193-9006](https://orcid.org/0000-0002-8193-9006)

### Jiahua Deng

Department of Chemistry, Boston University, Boston, Massachusetts 02215, United States  
ORCID iD: [0000-0001-8865-4786](https://orcid.org/0000-0001-8865-4786)

### Dylan Coelho

Université Paris Cité, CNRS, Laboratoire de Chimie et Biochimie Pharmacologiques et Toxicologiques, Paris F-75006, France

### Emmanuelle Braud

Université Paris Cité, CNRS, Laboratoire de Chimie et Biochimie Pharmacologiques et Toxicologiques, Paris F-75006, France

### Laura Iannazzo

Université Paris Cité, CNRS, Laboratoire de Chimie et Biochimie Pharmacologiques et Toxicologiques, Paris F-75006, France

### Yaozong Li

Department of Biochemistry, University of Zurich, Zurich CH-8057, Switzerland  
ORCID iD: [0000-0002-5796-2644](https://orcid.org/0000-0002-5796-2644)

**Danzhi Huang**

Department of Biochemistry, University of Zurich, Zurich CH-8057, Switzerland

**Mélanie Ethève-Quelquejeu**

Université Paris Cité, CNRS, Laboratoire de Chimie et Biochimie Pharmacologiques et Toxicologiques, Paris F-75006, France

ORCID iD: [0000-0002-4105-3243](https://orcid.org/0000-0002-4105-3243)

**Qiang Cui**

Department of Chemistry, Boston University, Boston, Massachusetts 02215, United States,  
Department of Physics, Boston University, Boston, Massachusetts 02215, United States,  
Department of Biomedical Engineering, Boston University, Boston, Massachusetts 02215,  
United states

ORCID iD: [0000-0001-6214-5211](https://orcid.org/0000-0001-6214-5211)

**Amedeo Caflisch**

Department of Biochemistry, University of Zurich, Zurich CH-8057, Switzerland

**For correspondence:** [caflisch@bioc.uzh.ch](mailto:caflisch@bioc.uzh.ch)

ORCID iD: [0000-0002-2317-6792](https://orcid.org/0000-0002-2317-6792)

**Copyright**

© 2023, Corbeski et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

**Editors**

Reviewing Editor

**Aaron Frank**

Arrakis Therapeutics, Waltham, United States of America

Senior Editor

**Amy Andreotti**

Iowa State University, Ames, United States of America

**Reviewer #1 (Public Review):**

Summary:

This important study nicely integrates a breadth of experimental and computational data to address fundamental aspects of RNA methylation by an important for biology and health RNA methyltransferases (MTases).

**Strengths:** The authors offer compelling and strong evidence, based on carefully performed with appropriate and well-established techniques to shed light on aspects of the methyl transfer mechanism of the methyltransferase-like protein 3 (METTL3), which is part of the methyltransferase-like proteins 3 & 14 (METTL3-14) complex.

There are no weaknesses that we identified in the revised version.

<https://doi.org/10.7554/eLife.92537.2.sa2>

**Reviewer #2 (Public Review):****Summary:**

Caffisch and coworkers investigate the methyltransferase activity of the complex of methyltransferase-like proteins 3 and 14 (METTL3-14). To obtain an high resolution description of the complete catalytic cycle they have carefully designed a combination of experiments and simulations. Starting from the identification of bisubstrate analogues (BAs) as binder to stabilise a putative transition state of the reaction they have determined multiple crystal structures and validated relevant interactions by mutagenesis and enzymatic assays.

Using the resolved structure and classical MD simulations they obtained a kinetic picture of the binding and release of the substrates. Of note, they accumulate very good statistics on these processes using 16 simulation replicates over a time scale of 500 ns. To compare the time scale of the release of the products with that of the catalytic step they performed state-of-the-art QM/MM free energy calculations (testing multiple levels of theory) and obtain a free energy barrier that indicates how the release of the product is slower than the catalytic step.

**Strengths:**

All the work proceeds through clear hypothesis testing based on a combination of literature and new results. Eventually, this allows them to present in Figure 10 a detailed step-by-step description of the catalytic cycle. The work is very well crafted and executed.

<https://doi.org/10.7554/eLife.92537.2.sa1>

**Reviewer #3 (Public Review):****Summary:**

The manuscript by Coberski et al describes a combined experimental and computational study aimed to shed light on the catalytic mechanism in a methyltransferase that transfers a methyl group from S-adenosylmethionine (SAM) to a substrate adenosine to form N6-methyladenosine (m6A).

**Strengths:**

The authors determine crystal structures in complex with so-called bi-substrate analogs that can bridge across the SAM and adenosine binding sites and mimic a transition state or intermediate of the methyl-transfer reaction. The crystal structures suggest dynamical motions of the substrate(s) that are examined further using classical MD simulations. The authors then use QM/MM calculations to study the methyl-transfer process. Together with biochemical assays of ligand/substrate binding and enzyme turnover, the authors use this information to suggest what the key steps are in the catalytic cycle. The manuscript is in most places easy to read.

**Weaknesses:**

After revising the manuscript, there are few weaknesses beyond those listed in the paper.

<https://doi.org/10.7554/eLife.92537.2.sa0>

**Author Response**

The following is the authors' response to the original reviews.

## Public Reviews:

### Reviewer #1 (Public Review):

#### Summary:

*This important study nicely integrates a breadth of experimental and computational data to address fundamental aspects of RNA methylation by an important for biology and health RNA methyltransferases (MTases).*

#### Strengths:

*The authors offer compelling and strong evidence, based on carefully performed work with appropriate and well-established techniques to shed light on aspects of the methyl transfer mechanism of the methyltransferase-like protein 3 (METTL3), which is part of the methyltransferase-like proteins 3 & 14 (METTL3-14) complex.*

#### Weaknesses:

*The significance of this foundational work is somewhat diminished mostly due to mostly efficient communication of certain aspects of this work. Parts of the manuscript are somewhat uneven and don't quite mesh well with one another. The manuscript could be enhanced by careful revision and significant textual and figure edits. Examples of recommended edits that would improve clarity and allow accessibility to a broader audience are highlighted in some detail below.*

We thank the reviewer for the positive evaluation of our work. We have followed the suggestions and modified the text and figures as detailed further in our answers to the specific recommendations.

### Reviewer #2 (Public Review):

#### Summary:

*Caflich and coworkers investigate the methyltransferase activity of the complex of methyltransferaselike proteins 3 and 14 (METTL3-14). To obtain a high-resolution description of the complete catalytic cycle they have carefully designed a combination of experiments and simulations. Starting from the identification of bisubstrate analogues (BAs) as binders to stabilise a putative transition state of the reaction, they have determined multiple crystal structures and validated relevant interactions by mutagenesis and enzymatic assays.*

*Using the resolved structure and classical MD simulations they obtained a kinetic picture of the binding and release of the substrates. Of note, they accumulated very good statistics on these processes using 16 simulation replicates over a time scale of 500 ns. To compare the time scale of the release of the products with that of the catalytic step they performed state-of-the-art QM/MM free energy calculations (testing multiple levels of theory) and obtained a free energy barrier that indicates how the release of the product is slower than the catalytic step.*

#### Strengths:

*All the work proceeds through clear hypothesis testing based on a combination of literature and new results. Eventually, this allows them to present in Figure 10 a detailed step-by-step description of the catalytic cycle. The work is very well crafted and executed.*

We thank the reviewer for the positive evaluation of our work.

*Weaknesses:*

*To fulfill its potential of guiding similar studies for other systems as well as to allow researchers to dig into their vast work, the authors should share the results of their simulations (trajectories, key structures, input files, protocols, and analysis) using repositories like Zenodo, the plumed-nest, figshare or alike.*

The reviewer is right. We have uploaded the simulation materials to Zenodo: the MD simulation data (trajectories, pdb files, parameter files), and the PLUMED file that was used for the DFTB3/MM metadynamics simulations. We provide the link in the “Data availability” section.

**Reviewer #3 (Public Review):**

*Summary:*

*The manuscript by Coberski et al describes a combined experimental and computational study aimed to shed light on the catalytic mechanism in a methyltransferase that transfers a methyl group from Sadenosylmethionine (SAM) to a substrate adenosine to form N6-methyladenosine (m6A).*

*Strengths:*

*The authors determine crystal structures in complex with so-called bi-substrate analogs that can bridge across the SAM and adenosine binding sites and mimic a transition state or intermediate of the methyltransfer reaction. The crystal structures suggest dynamical motions of the substrate(s) that are examined further using classical MD simulations. The authors then use QM/MM calculations to study the methyl-transfer process. Together with biochemical assays of ligand/substrate binding and enzyme turnover, the authors use this information to suggest what the key steps are in the catalytic cycle. The manuscript is in most places easy to read.*

We thank the reviewer for the positive evaluation of our work.

*Weaknesses:*

*My main suggestion for the authors is that they show better how their conclusions are supported by the data. This includes how the electron density maps for example support the key interactions and water molecules in the active site and a better error analysis of the computational analyses.*

We thank the reviewer for the comments and suggestions. We have followed the suggestions and added error analysis of the computational results as well as additional figures (in the supplementary information) that illustrate key interactions and water molecules in the active site supported by the electron density.

**Reviewer #1 (Recommendations For The Authors):**

- *The phrasing of the second sentence in the introduction is difficult to read. I am not sure it is necessary to define the DRACH motif if you are also giving the exact consensus sequence unless providing more context for other instances of the DRACH motif. Referring to this motif instead as "consensus sequence GGACU" may be more effective.*

The reviewer is right. We corrected the sentence accordingly.



- *In the second paragraph of the introduction, a further short description of how METTL3-14 is "involved" in diseases would be appreciated.*

We thank the reviewer for the comment. We made that clearer by including “by promoting the translation of genes involved in cell growth, differentiation, and apoptosis” together with a reference.

- *Is there any evidence that inhibiting METTL3-14 doesn't negatively impact healthy cells?*

We thank the reviewer for the question. Yes, there is such evidence and we added to the sentence “but not in normal non-leukaemic haemopoietic cells” together with a reference to make this point clearer.

- *Bringing up the MACOM complex in the third paragraph of the introduction is perhaps not necessary unless further discussing the MACOM complex later.*

The reviewer is right. We removed the mention of the MACOM complex.

- *Figure 1B: Color coding is difficult to distinguish on a screen and print out. More contrasting colors would be helpful.*

We thank the reviewer for the suggestion. We removed the transparency from the protein cartoon representation that was the reason for the low contrast.

- *The level of detail in the "MD simulations for mechanistic studies of RNA MTases" is not advised. Would strongly encourage condensing this section to improve clarity and accessibility to a larger audience.*

The reviewer is right. We removed non-essential parts of this paragraph.

- *Confirming the role of the hydroxyl in Y406 would be better supported by a Y406 -> F406 mutant because the A406 mutant could bind differently due to a loss of pi-stacking interactions.*

The purpose of the Y406A mutant was to eliminate the interaction of the aromatic sidechain with adenosine as seen from the structure with BA4. Since there is no involvement of the Y406-OH group with adenosine, mutating to F did not seem sufficient. Furthermore, by mutating Y406 to alanine, we also eliminate the possibility for a water-mediated hydrogen bond to the W398 backbone. Hence, with the alanine mutant we achieve the strongest possible effect on the enzymatic activity while the integrity of the active site is maintained as seen from the thermal shift assay.

- *For Figure 4D, can the authors justify why SAH was used as a metric for SAM binding instead of using SAM directly? Additionally, referring to the RNA as "ligand" instead of "RNA" in the Figure caption is more confusing than simply calling it RNA.*

We thank the reviewer for the comment. With the TSA, we wanted to show that with the adenosine binding mutants, the integrity of the METTL3 active site is still intact. It was shown that SAH is bound with higher affinity than SAM by METTL3 (DOI:

10.1016/j.celrep.2019.02.100). Since the magnitude of the thermal shift depends also on the affinity, we chose the higher-affinity binder SAH. There is no RNA per se shown in this figure. “Ligands” in the figure caption (A) refers to the three bound molecules that are shown and mentioned in the previous sentence: SAM, BA2, and BA4. “Ligand” in the figure caption (D) refers to “SAH” that was used in the experiment described and mentioned just after, but is now removed.

- *Figure 5D is very difficult to interpret. Removing the ribbons representing Y406 movement may make it easier to see. Color coding the Supplementary Movie 1 to match would be also helpful.*

The reviewer is right. We have changed the figure to make the different conformations of METTL3 and its Y406 sidechain clearer. However, we left the coloring of the different conformations as the colors are connected to different time points of the simulation. Following the suggestion of the reviewer we changed the coloring of SAM and AMP to match that of the supplementary movie.

- *Figure 10 is overwhelming as is. Removing the grey area around the binding sites and toning down the color of the substrate binding sites would help with visibility. The size of the chemical structures and illustrations is currently too small to easily be made out. A full page-sized figure may be beneficial for this figure.*

We agree with the reviewer and have changed the figure to make each reaction step clearer and better recognizable.

Minor >edits

- *Change "Despite the growing knowledge on the diverse pathways" to "Despite growing knowledge of the diverse pathways involving METTL3-14".*

We corrected the sentence.

- *Perhaps use "redundant active site" instead of "degenerate active site".*

We changed the word as suggested.

- *Consider moving "The METTL3 MTase domain has the catalytically active SAM binding site and adopts a Rossmann fold that is characteristic of Class I SAM-dependent MTases" to before "METTL14 also has an MTase domain, however, with a degenerate active site of hitherto unknown function, and so-called RGG repeats at its C-terminus essential for RNA binding" to keep information about METTL3 together.*

We shifted the part of the text as suggested.

- *"Molecular dynamics studies have mainly focused on protein and bacterial MTases"? Does this mean bacterial MTases that methylate proteins?*

We thank the reviewer for the comment. This means bacterial MTases in general. The example that we mention is of a bacterial MTase that methylates a chemical precursor. We changed the sentence slightly to make that clearer.

- In "Bisubstrate analogues bind in the METTL3 active site", please consider the following:
- Change "and to investigate" to "and investigated".
- Briefly describe the enzymatic assay in the main text.
- Either more clearly defining "least potent" or change to "have the highest IC50 values".

We made all the suggested changes to improve the description of the assay and its outcomes.

- In Figure 3, remove some of the amino acid labels from panels A, C, and E for clarity, especially since panels B, D, and F more clearly demonstrate the interactions.

We removed amino acids that were not involved in polar contacts and adapted the figure caption accordingly.

- In panels 3D, 3F, and 4B, the lightning bolts are too small to make out as lightning bolts. An asterisk or other symbol may be easier to distinguish.

We made the lightnings more than double the size to make them better recognizable.

- In Figure 4C, no units are provided on the y-axis. Additionally, I do not believe the arrows indicating "Loss of activity" are necessary.

These are arbitrary units as it is a ratio which is explained in the materials and methods section. We removed the arrows following the suggestion of the reviewer.

- While demonstrating mutants with no activity still retain SAM binding is suggestive of the mutant impacting RNA binding, this would still be better supported with RNA binding studies. Electrophoretic mobility shift assays would be sufficient if Tm studies are time-consuming. While these experiments could be informative, we also acknowledge that they may be outside the scope of this current report.

We thank the reviewer for suggesting these experiments and acknowledging that they would be outside of the scope of the current study. Such RNA binding experiments can turn out to be very time consuming, both in TSA and EMSA. The reason is mainly this: The RNA substrate must be chosen such that it binds sufficiently strong to the WT to cause an effect (thermal shift or electrophoretic mobility shift), but also to observe a clear difference in binding between WT and mutant proteins. Since many more residues of METTL3 and METTL14 contribute to RNA binding, the effects of individual mutants on affinity might be too small to be confidently detected in TSA or EMSA. In particular, we only identified the substrate adenosine binding residues, and mutating them and hence preventing adenosine binding alone, might not have a big effect on overall RNA binding affinity. The enzymatic assay that we used, on the other hand, is more sensitive since the detection is fluorescence based and quantifies the conversion of A to m6A in an RNA substrate, and more factors than just affinity play a role for enzymatic activity, such as correct orientation and stability of the adenosine in the active site and stabilization of the transition state.

- *A written narrative to accompany Supplementary Movie 1 would make it much more accessible to those unfamiliar with modeling and simulations.*

We thank the reviewer for the comment. We expanded the caption to the movie with a narrative describing different events at different time points in the movie.

- *Table 3 could be made clearer to those without MD experience by defining/indicating the top row as different computational models.*

The reviewer is right. We have added a footnote to Table 3 to clearly indicate the different density functional theory and semi-empirical density functional tight binding method used in this study. We also added another line in the table.

- *In the conclusion, the authors state "the height of the QM/MM free-energy barrier indicates that the methyl transfer step is not rate-determining." How does this compare to experimental data? Additional kinetic assays to demonstrate this experimentally would go a long way in convincing the reader of this conclusion.*

We thank the reviewer for the question. Kinetic assays have been performed for METTL3-14 and we mention and reference them in the text. We believe that further kinetic experiments would be outside of the scope of this study. Furthermore, the METTL3 mutants that we made show no activity in our enzymatic assay and hence kinetic studies would be probably impossible to do with them.

As we show from QM/MM and describe in the text, the methyl cation in the SAM cofactor is transferred directly to the N6 position of the adenosine substrate. DFTB3/MM free energy simulations show that this mechanism has an energetic barrier of 15-16 kcal/mol. The turnover as published based on an enzymatic assay is 0.2-0.6 min<sup>-1</sup> at ambient temperature which implies a barrier of ~20 kcal/mol. This value is higher than that determined for the methyl transfer alone as determined by QM/MM. Hence, in the overall mechanism, there must be a step that is slower than the methyl transfer and hence we conclude that the methyl transfer is not the rate-limiting step.

### **Reviewer #3 (Recommendations For The Authors):**

*I only have a few comments about the work.*

*(1) It would be good if the authors could show more of the data that is used as the basis for their conclusions. For example, IC<sub>50</sub> values are presented (Table 1) without error estimates or an indication of the quality of the data that is used to estimate the data.*

We thank the reviewer for the suggestion. We included errors of the IC<sub>50</sub> values and show the dose response curved from the enzymatic assay with the BAs as inhibitors in a new Supplementary Figure S1.

*(2) More substantially, it would be good to have a more detailed analysis of the crystal structures in terms of the properties that are mentioned/analysed. While the structures are relatively good (2.1 Å-2.5Å), it is not clear to the reader how this data supports the interactions that are proposed. For example, the authors pinpoint a number of hydrogen bonding interactions and water molecules in the complexes. They might consider showing support for some of these in the electron density maps. Similarly, it would be good to show the densities that support the substantial differences of the Ade in the BA2 and BA4 complexes. These might be supplementary files. I note also that the structures*

*are not yet released or available for analysis [which of course is a valid choice but also means that I cannot inspect the maps myself].*

We have added supplementary figures supporting the conformations of the BAs and their interactions with METTL3 with electron density, for BA1 and BA6 in Supplementary Figure S2, and for BA2 and BA4 in a new Supplementary Figure S3.

*(3) It would be useful with an error analysis of the off-rates estimated from the MD simulations and a discussion of the accuracy of these estimates. Even the slower dissociation events seem quite fast. What are the rough affinities of these molecules and how fast would the binding need to be to be compatible with the affinity and estimated off-rates?*

We expanded upon this in the results paragraph concerning the MD simulations. The affinities of METTL3-14 binding to AMP or m6AMP can be expected to be very low, with Kd values in the millimolar range. We have not measured these Kd values, nor have we found any published data, but we have conducted thermal shift assays with A and m6A and did not observe any significant thermal shifts in the melting temperature of METTL3-14 at high micromolar concentrations of these compounds, indicative of a very low binding affinity. This is to be expected because METTL3-14 should not methylate adenosines unspecifically but rather in the GGACU motif of substrate mRNA.

*(4) The authors use QM/MM simulations with metadynamics to estimate the energy profile of the methyl transfer reaction. They find a barrier of ca. 15 kcal/mol and suggest this to be compatible with the enzymatic turnover rate of ca. 0.3/min. Here it would be good with a clearer description of the possible sources of error and assumptions in making these statements. First, what is the error on the estimated energy profile from the metadynamics? The authors mention the analysis of progression of the PMF as a function of time, but that is in itself not a strong test for convergence (the PMF may stay constant if there is little sampling). What does the time series of the CV look like? Second, it seems as if the authors are assuming a large pre-exponential factor ( $10^9/s$ ?). Is that correct, and how sure are they of this value? Finally, when linking the barrier of the methyl-transfer reaction to the overall turnover rate it sounds like they assume that other parts of the reaction do not affect the turnover rate. Is that correctly understood, and what is the evidence for that? It sounds like the authors are saying that step 5 in the cycle (Figure 10) is limiting.*

We thank the reviewer for the questions. Accordingly, we have carried out additional simulations and statistical error analyses.

(i) We have carried out two additional sets of multi-walker metadynamics simulations with the same setup as the original calculation, except for using different initial random seeds. Using the three independent sets of metadynamics simulations, we can better estimate the statistical uncertainty for the computed potential of mean force (PMF). We have updated the PMF in Fig. 8b, in which the solid curve represents the result averaged over three independent runs, and the shaded area represents the standard error of the mean of the three replicas. The figure caption of Fig. 8b is revised accordingly.

(ii) To further illustrate the convergence behavior of the metadynamics simulations, we have included the following supplementary files: (1). Potentials of mean force computed with different numbers of deposited Gaussians are compared. (2). As suggested by the reviewer, we show the time series of the collective variable (CV) sampled by the 24 independent walkers during one set of metadynamics simulations. These results clearly indicate that the CV exhibits diffusive behaviors between the reactant and product regions, further supporting the adequate sampling and convergence of our metadynamics simulations.

(iii) Regarding the issue of pre-factor used in the rate estimate, we have indeed used the common approximation of  $kT/h$  as in the regular transition state theory. Many studies in the literature support the use of this expression for very localized chemical reactions in enzymes. We have included several representative references along this line: (1) M. Garcia-Viloca, J. Gao, M. Karplus, D. G. Truhlar, How enzymes work: Analysis by modern rate theory and computer simulations, *Science*, 303, 186-195 (2004) (2) D. R. Glowacki, J. N. Harvey, A. J. Mulholland, Taking Ockham's razor to enzyme dynamics and catalysis, *Nat. Chem.* 4, 169-176 (2012)

(iv) Regarding the nature of the rate-limiting event, please see our response to reviewer 1.

*(5) The authors should ideally make the input files for their simulations available and deposit the plumed files in for example plumed-nest (as indicated in their reference 100).*

We agree with the reviewer. Accordingly, we have uploaded the PLUMED file that we have used for the DFTB3/MM metadynamics simulations (plumed.dat) together with the MD simulation trajectories to Zenodo.

*Minor*

*(1) Many of the details in Figure 10 are very small and difficult to read without zooming in. Consider whether some parts could be made larger.*

The reviewer is right. We have changed the figure to make each reaction step clearer and better recognizable.

*In conclusion, we would like to thank you for the very efficient handling of our manuscript, and the reviewers for the useful comments and suggestions. We think that the manuscript has improved and it should be accepted in its revised form at eLife.*