



# Early evolution of the ecdysozoan body plan

Deng Wang, Yaqin Qiang, Junfeng Guo , Jean Vannier, Zuchen Song, Jiaxin Peng, Boyao Zhang, Jie Sun, Yilun Yu, Yiheng Zhang, Tao Zhang, Xiaoguang Yang, Jian Han 

State Key Laboratory of Continental Dynamics, Shaanxi Key Laboratory of Early Life & Environments and Department of Geology, Northwest University; Xi'an, China • School of Earth Science and Resources, Key Laboratory of Western China's Mineral Resources and Geological Engineering, Ministry of Education, Chang'an University; Xi'an, China • Université de Lyon, Université Claude Bernard Lyon 1, ENS de Lyon, CNRS, Laboratoire de Géologie de Lyon: Terre, Planètes, Environnement (CNRS-UMR 5276), Villeurbanne 69622, France • University of Chinese Academy of Sciences; Beijing, China • Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences; Beijing, China • School of Information Science and Technology, Northwest University; Xi'an, China

## Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.


## About eLife's process

**Reviewed preprint version 1**  
February 19, 2024 (this version)

**Posted to preprint server**  
January 18, 2024

**Sent for peer review**  
December 3, 2023

 [https://en.wikipedia.org/wiki/Open\\_access](https://en.wikipedia.org/wiki/Open_access)

 Copyright information

## Abstract

Extant ecdysozoans (moulting animals) are represented by a great variety of vermiform or articulated organisms. However, controversies remain about the nature of their ancestral body plan although the vermiform hypothesis seems to prevail. We describe here *Beretella spinosa* gen et sp. nov. a tiny ecdysozoan from the early Cambrian, Yanjiahe Formation, South China, with an unusual sack-like appearance, single opening, and spiny ornament. *Beretella* has no equivalent among animals, except *Saccorhytus* from the basal Cambrian. Phylogenetic analyses resolve both forms as a sister group (Saccorhytida) to all known Ecdysozoa, thus suggesting that ancestral ecdysozoans may have been non-vermiform animals. Saccorhytids are likely to represent an early dead-end off-shot along the stem-line Ecdysozoa that possibly evolved through anatomical simplification (e.g. lack of anus). Although extinct during the Cambrian, this animal lineage provides precious insight into the early evolution of Ecdysozoa and the nature (possibly non-vermiform) of the earliest representatives of the group.

### eLife assessment

This study provides a **fundamental** advance in palaeontology by reporting the fossils of a new invertebrate, *Beretella spinosa*, and inferring its relationship with already described species. The analysis placed the newly described species in the earliest branch of moulting invertebrates. The study, supported by **convincing** fossil observation, hypothesizes that early moulting invertebrate animals were not vermiform.

## Introduction

The Ediacaran–Cambrian transition is marked by the appearance in the fossil record of a variety of new body plans that prefigure the majority of present-day animal lineages, including the ecdysozoans a huge clade that encompasses all invertebrate animals growing through successive moulting stages, such as panarthropods (Arthropoda, Onychophora, Tardigrada), scalidophoran (incl. Priapulida) and nematoid worms (Erwin, 2020 [↗](#)). Altogether ecdysozoans represent a very high percentage of animal biodiversity and disparity, inhabiting almost all possible ecological niches on Earth (Brusca et al., 2016 [↗](#)). The nature of the last common ancestor of Ecdysozoa (LCAE) remains largely unresolved, even though worms are prevalent before the rise of panarthropods as trace and body fossils in basal Cambrian and late Ediacaran rocks (Buatois et al., 2014 [↗](#); Liu et al., 2014 [↗](#); Vannier et al., 2010 [↗](#)). Some molecular phylogenies also predict that the most basal ecdysozoans were vermiform organisms (Howard et al., 2022 [↗](#); Laumer et al., 2019 [↗](#)) that possibly diverged in the Ediacaran (Howard et al., 2022 [↗](#); Rota-Stabelli et al., 2013 [↗](#)). Current reconstruction based on fossil and developmental evidence features the ancestral ecdysozoan as a millimeter-sized worm (Budd, 2001 [↗](#); Valentine and Collins, 2000 [↗](#)) with a terminal (Ortega-Hernandez et al., 2019 [↗](#)) or ventral mouth (Martín-Durán and Hejnol, 2015 [↗](#); Nielsen, 2019 [↗](#)). Clearly, the discovery of *Saccorhytus* (Han et al., 2017 [↗](#); Liu et al., 2022 [↗](#); Shu and Han, 2020b [↗](#)) in the basal Cambrian of China (Kuanchuanpu Formation; ca. 535 Ma (Sawaki et al., 2008 [↗](#))) that is anything but a worm sowed doubt among scientists. *Saccorhytus* is a sac-like secondarily phosphatized microscopic animal spiked with conical sclerites and a single opening that was first seen as the earliest known deuterostome (Han et al., 2017 [↗](#)) but is now considered as an ecdysozoan on more solid grounds (Liu et al., 2022 [↗](#); Shu and Han, 2020b [↗](#)), thus broadening the anatomical spectrum of the group and its disparity in the Cambrian and reopening the debate on the nature of LCAE.

We describe here *Beretella spinosa* gen. et sp. nov. from Member 5 of the Yanjiahe Formation (basal Cambrian Stage 2, ca. 529 Ma, Hubei Province, China) that shares morphological traits with *Saccorhytus* such as an ellipsoidal body, a pronounced bilaterality, a spiny ornament made of broad-based sclerites, and a single opening. Cladistic analyses are made to resolve the position of both *Beretella* and *Saccorhytus* that provide key information on the early evolution of the group.

## Systematic palaeontology

Superphylum Ecdysozoa Aguinaldo et al. (Aguinaldo et al., 1997 [↗](#)) Phylum Saccorhytida Han, Shu, Ou and Conway Morris, 2017 stat. nov.

### Remarks

Saccorhytida first appeared in the literature as a new stem-group Deuterostomia that accommodated a single species, *Saccorhytus coronarius* (Han et al., 2017 [↗](#)). Since *Saccorhytus* is no longer considered a primitive deuterostome and, instead, more likely belongs to ecdysozoans, Saccorhytida became an extinct Order of Ecdysozoa (Liu et al., 2022 [↗](#); Shu and Han, 2020b [↗](#)). Because both *Saccorhytus* and *Beretella* display major morphological differences with all other known ecdysozoan phyla (Nematoida, Scalidophora, and Panarthropoda), Saccorhytida is tentatively elevated here to the rank of phylum within Ecdysozoa.

### Emended diagnosis

Microscopic, ellipsoidal body shape with pronounced bilateral symmetry expressed by paired spiny sclerites. Single, presumably oral opening on assumed ventral side (no anus).

## Remarks

Only two forms, *Saccorhytus* and *Beretella* are currently placed within Saccorhytida, making it premature to formally define intermediate taxonomic categories such as an order and a family.

*Beretella spinosa* Han, Guo, Wang and Qiang, gen. et sp. nov.

## Etymology

From ‘*béret*’, French, that designates a soft, visorless cap referring to the overall shape of this species, and ‘*spinosa*’, an adjective (Latin), alluding to its spiny ornament.

## Holotype

CUBar138-12 (**Fig. 1a–c** [↗](#)).

## Paratype

CUBar171-5 **Fig. 1h, i** [↗](#)) and CURBar121-8 (**Fig. 1j, k** [↗](#)).

## Diagnosis

Body with a beret-like lateral profile. Convex side (presumably dorsal) with an elevated (presumably posterior) and lower (presumably anterior one) end. The opposite side (presumably ventral) flattened. Bilateral symmetry well expressed in the overall body shape (sagittal plane) and sclerite distribution. Antero-posterior polarity. Convex side with a slightly elevated sagittal stripe topped with a single row of four aligned spines (S1) and five additional spines (S2) on each side. Six broad-based conical sclerites (S3) distributed in two symmetrical longitudinal rows plus two sagittal ones. Double rows of six marginal spines (S4 and S5). Flattened side often pushed in and partly missing, bearing a possible mouth opening. Possible oral spine.

## Stratigraphy and locality


*Watsonella crosbyi* Assemblage Zone (Guo et al., 2021 [↗](#)), Member 5 of the Yanjiahe Formation (Cambrian Terreneuvian, Stage 2) in the Yanjiahe section near Yichang City, Hubei Province, China (Supplementary Figs. 1, 2).

## Description and comparisons

The body of *Beretella spinosa* is secondarily phosphatized and has a consistent beret-like three-dimensional shape in the lateral view. Its maximum length, width, and height range from 1.0–2.9 mm, to 975–2450  $\mu\text{m}$ , and 500–1000  $\mu\text{m}$ , respectively (**Fig. 1** [↗](#), Supplementary Tables 1–3). The ratio of the maximal length to width is 1.6:1 (Supplementary Fig. 3). As seen in top view, *B. spinosa* shows a small lateral constriction at approximately mid-length (**Fig. 1a, c** [↗](#)).

The body has a convex, assumedly dorsal side with one, presumably posterior end more elevated than the other (**Fig. 1b, e, i, k** [↗](#)). This elevation is gradual along the sagittal plane and then becomes more abrupt near the low elevated, presumably anterior end. The opposite, assumedly ventral side is less well preserved and seems to have been originally flattened.

The convex side bears a complex ornamented pattern made of five sets (S1–S5) of spiny sclerites directed towards the more elevated end (**Figs. 1a, b, d, e, h–k** [↗](#), **2a, b, d** [↗](#)). These sclerites were originally pointed (**Figs. 1a, d, e** [↗](#), **2b, k, l** [↗](#), Supplementary Fig. 4a, b, g), but most of them were



**Fig. 1.**

***Beretella spinosa* gen. et sp. nov. from Member 5 of the Yanjiahe Formation (Cambrian Stage 2), Yichang, Hubei Province, China.**

**a–c,** Holotype, CUBar138-12. **a,** Dorsal view showing the external ornament: (five sclerites at the midline in yellow (S1); flanked by two rows of sclerites in blue (S2); large broad-based conical sclerites in two dorsolateral pairs and one antero-posterior pairs in green (S3)); white arrows indicate lateral constriction. **b,** Right lateral view showing two additional rows of six sclerites (S4 and S5, in light blue and pink, respectively). **c,** Ventral view showing a large opening that may have accommodated the mouth (see the text) and an empty body cavity. **d–g,** CUBar75-45. **d,** Dorsal view showing a broken S3. **e,** Micro-CT image, right lateral view displaying S4. **f,** Ventral view depicting a tiny projection in purple. **g,** An enlargement of the projection of **f**. **h–i,** Paratype, CUBar171-5. **h,** Right dorsal view showing S1–S4. **i,** Right-lateral view showing S4 and S5. **j–k,** Paratype CUBar121-8. **j,** Dorsal view showing poorly preserved S1 and S2. **k,** Right-lateral view showing S3–S5. A, assumed anterior end (see text); ef, exotic fragment; D, assumed dorsal side; L, left; P, posterior end; R, right; tp, tiny spine; V, ventral side. The same abbreviations are used throughout the manuscript including Supplementary materials.

broken thus revealing an internal cavity and an ellipsoidal transverse section (Figs. 1a, b, h–k, 2a–e, g). The broken sclerites show an inner and outer phosphatic layer (thickness ca. 20 to 50  $\mu\text{m}$ ) often separated by a thin empty space (Fig. 2g–i).

The convex side bears six prominent conical sclerites (S3) all with a rounded to elliptical well-delimited broad base, distributed in two longitudinal symmetrical pairs with two additional sclerites at both ends of the sagittal plane (Figs. 1, 2d, Supplementary Figs. 3e–h, 4e–i). A low-relief stripe runs in a sagittal position and vanishes towards the elevated end. It is topped by a row of aligned spines (S1, Fig. 1a); the one closer to the more elevated end being more tubular and longer. This row is flanked on both sides by smaller aligned spines (S2, Figs. 1a, d, h, 2a–c). Two relatively sinuous rows of six tiny spines are present parallel to the lateral margins (S4 and S5, Figs. 1b, e, h–j, 2d, e). The convex side bears a polygonal micro-ornament (mesh size ca. 5  $\mu\text{m}$  wide, Fig. 2f, Supplementary Tables 1–3). However, its exact extension is uncertain due to coarse secondary phosphatization. Clusters of spherical phosphatized grains (diameter ca. 20  $\mu\text{m}$ ) occur near the sclerite base (Supplementary Fig. 4b).

In most specimens, the flattened side is occupied by a relatively large opening (1200 and 600  $\mu\text{m}$  in maximal length and width, respectively) with irregularly defined margins (Fig. 1c, f, see also supplementary information movies 1, 2). The flattened side is often largely missing and opens into a spacious internal cavity with no signs of internal organs (e.g. gut and pharynx) (Fig. 1c, f). One specimen shows a tiny spine on the margin of the flattened side (Fig. 1f, g), which differs from other spiny sclerites (S1–S5).

The length of studied specimens ranges from 1.0 to 2.9 mm (Supplementary Fig. 3e–h). Whether growth was continuous or instead took place via successive moulting stages and cuticular renewal (ecdysis) could not be tested due to the small number of specimens (N=17) available for measurements. No major morphological variations (e.g. a sclerite pattern) can be seen between the smallest and largest specimens of *B. spinosa* (Supplementary Fig. 3e–h).

## Remarks


### Body polarities in *Beretella*

The anterior-posterior (AP) and dorsal-ventral (DV) polarities of *Beretella* are uneasy to define because of the lack of modern equivalent among extant animals. In the vast majority of extinct and extant invertebrates for which antero-posterior polarity is defined on the basis of independent criteria (e.g. position of mouth), sclerites point backwards (e.g. Cambrian scalidophoran worms (Han et al., 2007; Huang et al., 2004) and *Wiwaxia* (Zhang et al., 2015b)). This is most probably also the case with *Beretella* (Fig. 1a, d, j). The dorsoventral polarity of *Beretella* is supported by the fact that protective sclerites such as spines most commonly occur on the dorsal side of bilaterians (Fig. 1a, d, j).

### Comparison with *Saccorhytus* and other ecdysozoans

*Beretella spinosa* has no exact equivalent in any Cambrian animals except *Saccorhytus coronarius*, an enigmatic, sac-like ecdysozoan (Han et al., 2017; Liu et al., 2022; Shu and Han, 2020b). Both forms share a tiny, poorly differentiated ellipsoidal body, and a set of prominent bilaterally arranged spiny sclerites. Indeed, the broad-based conical sclerites (S3) of *Beretella* are almost identical to those of *Saccorhytus* (Supplementary Fig. 4c) and have counterparts among scalidophoran worms (Supplementary Fig. 4d). However, they differ in number, ornamented structures, shape, and spatial arrangement (see details in Supplementary Tables 1–3) which makes the hypothesis of *Saccorhytus* being the larval stage of *Beretella* unlikely. Both *Beretella* and *Saccorhytus* differ from other known ecdysozoans in the lack of a vermiform body, introvert, annulations, and through gut (Supplementary Tables 1–3).





**Fig. 2.**

***Beretella spinosa* gen. et sp. nov.**

**a**, CUBar99-19, dorsal view showing an ornament S1-S4. **b-c**, CUBar136-9, general dorsal view and details. **d**, CUBar136-11, dorsal view showing S1-S5. **e-f**, CUBar73-15 general view and details of the cuticular polygonal reticulation in black. **g-j**, CUBar128-27. **g-i**, general view and details of the bi-layered structure of the cuticular wall as seen in broken conical sclerites. **j**, Micro-CT section showing possibly sclerite infilling. **k-l**, CUBar99-18, cuticular fragment, general view and details of large sclerite (central feature represents possible phosphatic infilling). is, infilling sclerite; pr, polygonal reticulation; sb, sclerite base.

## Discussion

### Ventral mouth

All bilaterian animals have a digestive system with at least one opening that corresponds to the mouth (Brusca et al., 2016). Although the presumed oral area of *Beretella* is poorly preserved (ventral side often pushed in and largely destroyed), its mouth is likely to be found ventrally (see description), since no other opening occurs on its dorsal side, except those created by broken sclerites. The well-defined dorsoventral polarity of *Beretella* would suggest that the animal was resting on its ventral (flattened) side, the spiny dorsal side playing a protective role. Maintaining ventral contact with a substrate seems to be very unlikely unless these microscopic ellipsoidal animals were interstitial.


### Phylogenetic position of *Beretella*

*Beretella*'s phylogenetic affinities remain elusive due to the lack of information concerning its internal anatomy and ventral side. Its scleritome consists of isolated conical sclerites that were the cuticular outgrowths of a seemingly rigid integument that covered both sides of the animal. Such conical sclerites have close counterparts in Cambrian ecdysozoans such as scalidophoran worms (e.g. *Eokinorhynchus* (Zhang et al., 2015a)), lobopodians (e.g. *Onychodictyon ferox* (Hou et al., 1991)) and even more clearly *Saccorhytus* that recent cladistic analyses resolved as a branch of the total-group Ecdysozoa (Liu et al., 2022). These sclerites unknown in other animal groups, suggest that both *Saccorhytus* and *Beretella* belongs to Ecdysozoa in the absence of more direct fossil evidence such as exuviae or features suggesting cuticular moulting (Daley and Drage, 2016; Wang et al., 2019).

Cladistic analyses were performed to test the relation of *Beretella* and *Saccorhytus* to other ecdysozoan groups and, more generally, their phylogenetic relationships with other bilaterian groups (see details in Supplementary Table 4). Both taxa join in a clade (Saccorhytida, Fig. 3a–c) that is resolved as stem species within total-group Ecdysozoa and as the sister group of Cycloneuralia plus Panarthropoda, i.e. crown-group Ecdysozoa (Figs. 3d, 4, Supplementary Figs. 6–9). These results are consistent with the body plan of Saccorhytida being markedly different from that of crown-group ecdysozoans that all have a vermiform body and differentiated structures such as the introvert and pharyngeal complex (Fig. 4).

### The ancestral ecdysozoan body plan

Molecular clock analyses often place the divergence of Ecdysozoa relatively deep into the Ediacaran (Howard et al., 2022; Rota-Stabelli et al., 2013), thus highlighting major discrepancy with the known fossil record of the group. Potential ecdysozoans occur in the late Precambrian as suggested by 1) sclerites resembling those of extant priapulids, found in Ediacaran Small Carbonaceous Fossils assemblages (Moczyłowska et al., 2015) and 2) locomotion traces presumably made by scalidophoran worms (Buatois et al., 2014; Vannier et al., 2010). In the absence of fossil data for other vermiform groups such as nematoids, scalidophorans are potentially the oldest known representatives of Ecdysozoa. Recent Bayesian analyses based on a large molecular data set obtained from the 8 extant ecdysozoan phyla recover Scalidophora as the sister-group to Nematoda + Panarthropoda and suggest that ecdysozoans probably diverged in the Ediacaran possibly some 23 million years before the oldest fossil occurrence (trace fossils) of the group (Howard et al., 2022). Although this study does not speculate on the nature of the last common ancestor of Ecdysozoa, it is consistent with the view that the earliest representatives of the group were probably vermiform. Howard et al. (Howard et al., 2022) drew comparable conclusions based on *Acosmia*, an assumed stem-ecdysozoan worm from early Cambrian Chengjiang Lagerstätte. However, the re-evaluation of the morphological characteristics of this worm rather suggests a less basal position either within the total-group Cycloneuralia (Fig. 3d),




**Fig. 3.**

### Position of *Beretella spinosa* in the animal tree based on cladistic analysis.

**a-c**, artistic three-dimensional reconstructions of *Beretella spinosa* in the anterolateral, dorsal, and posterolateral views. **d**, Animal tree obtained from cladistic analyses using maximum likelihood tree obtained from cladistic analyses using maximal likelihood (IQTREE). *Saccorhytus* and *Beretella* join in a clade (new phylum Saccorhytida) resolved as the sister-group of all other ecdysozoans; numbers at key nodes denote probability. Fossil and extant taxa are in italics and bold, respectively. Known fossil record indicated by thicker bars (after (Shu and Han, 2020a)).





**Fig. 4.**

### Possible evolutionary scenario to explain the origin and early evolution of ecdysozoans.

**a**, Summary tree (see Supplementary Figs. 6-9) showing saccorhytids as a sister-group of Cycloneuralia (Nematoida plus Scalidophora) + Panarthropoda; main morphological features of each group listed along each branch. **b**, Potential evolutionary pathway to evolve Saccorhytida and crown-group Ecdysozoa. Numbers in green, red and blue circles designate pre-ecdysozoan (Spiralia), Saccorhytida and Cycloneuralia, respectively. Light brown gradient (circle) to emphasize ecdysis and sclerite secretion seen as key evolutionary steps. 1, Hypothetical pre-ecdysozoan animal with a ciliated epidermis and glycocalyx. 2, Saccorhytid exemplified by *Beretella* with a cuticle bearing sclerites and a simplified internal organization (e.g. loss of anus). 3, Crown-group ecdysozoan exemplified by a scalidophoran worm with an elongated shape, a differentiated head (introvert) and trunk, sclerites, a through gut, a terminal mouth and abilities to burrow into bottom sediment. Animals not to scale. Abbreviations: a, anus; ci, cilia; cu, cuticle; ec, epidermal cell; gl, glycocalyx (mucous layer); m, mouth; in, introvert; sc, sclerite; se, sediment; TGE, total-group Ecdysozoa. Silhouettes from [phylopic.org](https://www.phylopic.org).

Supplementary Figs. 6, 7) or among crown-group Ecdysozoa (Supplementary Figs. 8, 9). The non-vermiform nature of saccorhytids and their position as the sister group of the crown-group Ecdysozoa clearly reopens the debate on the nature of the ancestral ecdysozoan (Fig. 4) and suggests exploring various evolutionary hypotheses, in particular: 1) does the enigmatic saccorhytid body plan results from anatomical simplification? 2) to what extent may these animals shed light on the nature of the earliest ecdysozoans?

## Do saccorhytids result from simplification?

A relatively simple body plan and tiny size is often seen as resulting from anatomical simplification (e.g. digestive system) and miniaturization (micrometric size) in possible relation with the adaptation to specialized ecological niches or parasitism (Hanken and Wake, 1993). For example, some extant scalidophoran worms living in interstitial (meiobenthic) habitats such as loriciferans have a miniaturized body (Kirstensen, 1983) compared with their macroscopic counterparts (e.g. *Priapul* (Schmidt-Rhaesa, 2013b)). However, they retain a through gut and a functional introvert and show no sign of drastic internal simplification (Schmidt-Rhaesa, 2013a). Anatomical reduction is a typical feature of parasitism (Hanken and Wake, 1993) that is well-represented among extant ecdysozoans such as nematodes (Schmidt-Rhaesa, 2014). Although relatively small (ca. 0.1-2.5 mm long), nematodes underwent no simplification of their digestive system. Saccorhytids have no specialized features (e.g. anchoring or piercing structures) that would point to any adaptation to ecto- or endo-parasitic lifestyles (Cong et al., 2017). *Saccorhytus* has been interpreted<sup>16</sup> as a possible interstitial animal based on its micrometric size which corresponds to that of the extant meiofauna. If we accept the hypothesis of saccorhytids resulting from simplification, then we need to determine its origin. Simplification of saccorhytids from a vermiform animal (e.g. cycloneuralian worm with a through gut and terminal mouth) is difficult to conceive because it would involve considerable anatomical transformations such as the loss of vermiform organization, introvert and pharynx in addition to that of the digestive system (Fig. 4). Alternative options to consider are ancestral and not necessarily vermiform ecdysozoans.

## Early evolution of ecdysozoans: a new scenario

We propose here an alternative evolutionary hypothesis (Fig. 4) in which saccorhytids are replaced within the broader framework of the origin and early diversification of moulting animals. Saccorhytids are seen as an early off-shot from the stem-line Ecdysozoa (see cladistic analysis above) that possibly retained important features of the body plan of ancestral non-vermiform ecdysozoans (see ancestral character state reconstruction in Supplementary Table 4). This scenario must be considered as a working hypothesis whose aim is to stimulate research in this key area of animal evolution.

The cuticular secretion and the loss of cilia (Valentine and Collins, 2000) are seen as the first of a series of evolutionary events (Fig. 4) that led to the rise of Ecdysozoa. Moulting (shedding of the old cuticle via apolysis and its renewal) further reconciled body growth and cuticular protection (Schmidt-Rhaesa, 2007). Cuticle secretion and moulting may have been quasi-simultaneous innovations that took place over a relatively short time interval. The nature of the very first ecdysozoans is hypothetical and lacks fossil evidence. However, they are tentatively represented here as small epibenthic or interstitial slow-moving non-vermiform animals from which saccorhytids may have evolved via an assumed anatomical simplification (i.e. loss of anus seen details in Supplementary Table 4, Fig. 4).

In our scenario, this ancestral ecdysozoan stock would have also given rise to vermiform ecdysozoans through stepwise anatomical transformations such as the body elongation, the differentiation of key morpho-functional structures such as the pharynx and the introvert and the shift of the ventral mouth to a terminal position (Martín-Durán and Hejnol, 2015) (Fig. 4). This mouth shift from ventral to terminal arising in crown ecdysozoans is consistent with the

chronology of divergence of animal lineages and the fact that the mouth of most spiralian is ventral (Martín-Durán and Hejnol, 2015 [DOI](#); Nielsen, 2019 [DOI](#); Ortega-Hernandez et al., 2019 [DOI](#)). Developmental studies show that embryos of extant cycloneuralians have a ventral mouth that moves to a terminal position towards the adult stage (Martín-Durán and Hejnol, 2015 [DOI](#); Nielsen, 2019 [DOI](#)). These assumed major anatomical changes (e.g. functional introvert) must be placed in the ecological context of Cambrian animal radiation. Important changes in the functioning of marine ecosystems occurred in the early Cambrian such as interactive relationships between animal species, exemplified by predation (Vannier and Chen, 2005 [DOI](#); Vermeij, 1977 [DOI](#)) may have acted as drivers in the evolution of early ecdysozoans, in promoting burrowing into sediment and the colonization of endobenthic habitats for the first time (Vannier et al., 2010 [DOI](#)). Burrowing into the sediment could be seen as the evolutionary response of epibenthic animals such as ancestral ecdysozoans to escape visual predation (Daley et al., 2013 [DOI](#); Vannier and Chen, 2005 [DOI](#)). This migration to endobenthic shelters was made possible by the development of a resistant cuticular layer (Fig. 4 [DOI](#)) that strongly reduced physical damage caused by friction with the sediment and provided anchoring points (e.g. scalids and sclerites). Whereas saccorhytids became rapidly extinct during the Cambrian, worms massively colonized endobenthic habitats, resulting in bioturbation and ecological turnover.

## Methods

### Material

Fourteen specimens of *Beretella spinosa* were recovered from samples (siliceous-phosphatic, intraclastic limestone) collected from Member 5 of the Yanjiahe Formation, Yanjiahe section near Yichang City, Hubei Province, China (Guo et al., 2021 [DOI](#)) (Supplementary Tables 1–3). These were obtained by digesting the rocks in 10% acetic acid. Faunal elements associated with *Beretella spinosa* (Supplementary Tables 1–3) in residues are mainly tiny molluscs (CUBar21-4 and CUBar206-6). Comparisons were made with 10 specimens of *Saccorhytus coronarius* (ELIXX25-62, ELIXX34-298, ELIXX45-20, ELIXX48-64, ELIXX58-336, ELIXX61-27, ELIXX65-116, ELIXX65-296, ELIXX99-420) and one coeval scalidophoran specimen (ELIXX57-320) all from Bed 2 of the Kuanchuanpu Formation, Zhangjiagou section near Xixiang County, south Shaanxi Province, China. All specimens of *Beretella* are deposited in the paleontological collections of Chang'an University, Xi'an (CU), those of scalidophoran, and *Saccorhytus* at Northwest University, Xi'an (ELI), China.

### Scanning electron microscopy (SEM)

All specimens were coated with gold and then imaged using a FEI Quanta 400 FEG SEM at Northwest University and a FEI Quanta 650 at Chang'an University.

### X-ray computed microtomography and 3D reconstruction

Micro-CT-images (tiff format, with pixel size 1.1  $\mu\text{m}$ ) of *Beretella* (CUBar75-45, CUBar128-27, CUBar138-12) and *Saccorhytus* (ELIXX65-116, ELIXX99-420) were acquired using the Zeiss Xradia 520 at Northwest University (NWU), Xi'an, China, at an accelerating voltage of 50 kV and a beam current of 80  $\mu\text{A}$ . Micro-CT data were processed using VGStudio Max 3.2 for 3D volume rendering.

### Measurements

Measurements of the length, width, and height of *Beretella* and *Saccorhytus* were obtained from Micro-CT and SEM images by using tipDig2 v.2.16.

## Phylogenetic analysis

We built our matrix with 55 taxa coded using 191 morphological characteristics (Supplementary Texts 1, 2). It is largely based on the data published by Howard et al. (Howard et al., 2020), Vinther and Parry (Vinther and Parry, 2019) and Ou et al. (Ou et al., 2017), although emended and supplemented by recent updates and new observations (Supplementary Text 1). Three characters (37. Through gut, 38. U-shaped gut, and 40. Ventral mouth) in matrix were coded as “?” (uncertain), “?” and “?”, respectively. Because although we can infer a ventral mouth and no anus of *Beretella*, these anatomic structures are invisible in fossils. We analyzed the data matrix using parsimony (TNT), likelihood (IQ-TREE) and Bayesian inference (MrBayes). Parsimony analysis was implemented in TNT under equal and implied (k=3) weight. Parameters are default (Goloboff et al., 2008; Goloboff and Catalano, 2016). The maximum-likelihood tree search was conducted in IQ-TREE (Nguyen et al., 2015), and support was assessed using the ultrafast phylogenetic bootstrap replication method (Hoang et al., 2018; Minh et al., 2013) to run 50,000 replicates. Bayesian inference was conducted in with MrBayes v3.2.6a with default priors and Markov chain Monte Carlo settings (Ronquist et al., 2012). Two independent runs of 7,000,000 Markov chain Monte Carlo generations were performed, each containing four Markov chains under the Mk +  $\Gamma$  model for the discrete morphological character data (Lewis, 2001). In each run (N=2), trees were collected at a sampling frequency of every 5,000 generations and with the first 25% samples discarded as burn-in. The convergence of chains was checked by effective sample size (ESS) values over 1,000 in Tracer v1.7 (Rambaut et al., 2018), 1.0 for the potential scale reduction factor (PSRF) (Gelman and Rubin, 1992), and by an average standard deviation of split frequencies below 0.007.

## Ancestral character state reconstructions

Ancestral character state reconstructions for six morphological characters were performed on the ecdysozoan total group node, the ecdysozoan crown group node and saccorhytid node. Characters selected for ancestral state reconstruction represent traits inferred as ecdysozoan plesiomorphies (ancestral characters) from studies of crown group taxa. These characters included the presence or absence of: (1) through gut; (2) ventral mouth; (3) introvert (see Supplementary Table 4).

This was carried out individually for the selected character in MrBayes. This was employed to calculate the posterior probability of the presence (1) and absence (0) of the selected characters at the selected nodes. Analyses used the MK + gamma model, and always converged after 2 million generations. Average deviation of split frequencies (< 0.01), ESS scores (> 200), and PSRF values (= approx. 1.00) assessed convergence of the MCMC chains (Howard et al., 2020).

## Data availability

The data that support the findings of this study are available in the recent paper and its Supplementary Information.

## Acknowledgements

We thank H. G. for technical assistance. Funding: We thank the National Natural Science Foundation of China (grants 42172016, 41890844 to J.G., 41621003 to J.H., 42202009 to D.W.), the Strategic Priority Research Program of the Chinese Academy of Sciences (grant XDB26000000 grant to J.H. and J.G.), the China Post-doctoral Science Foundation (grant 2022M722568 to D.W.), the Key Scientific and Technological Innovation Team Project in Shaanxi Province (grant to J.G.), and the Région Auvergne Rhône Alpes and Université Claude Bernard Lyon 1 (grant to J.V.) for financial support.



## Author contributions

J.H. and J.G. conceived the research. J.G., Y.Q., Z.S., J.P., and B. Z. collected the material from Yanjiahe Formation. Y.Q. and D.W. prepared all the specimens, photographs, figures except **Figure 4** [\(J.V.\)](#). J.S. performed the analysis by Micro-CT and visualization with Micro-CT data. D.W. and J.H. performed phylogenetic analyses. Y.Y. performed morphospace analyses and Y.Z. and T.Z. performed computational fluid dynamic analyses in initial draft (not used in this version). D.W., J.V., J.H., J.G., and Y.Q. wrote the paper with input from all other authors. All authors approved the final manuscript.

## Competing interests

The authors declare no competing interests.

## Additional information

### Supplementary information

The online version contains supplementary material (tomographic data of *Beretella* and *Saccorhytus*, and movies of 3D-animation of the holotype of *Beretella*) available at <https://figshare.com/s/054f31fc22567a590d7f> [\(J.V.\)](#).

### Correspondence and requests for materials

should be addressed to J. G. or J. H.

## References

- Aguinaldo A.M.A., Turbeville J.M., Linford L.S., Rivera M.C., Garey J.R., Raff R.A., Lake J.A. (1997) **Evidence for a clade of nematodes, arthropods and other moulting animals** *Nature* **387**:489–93
- Brusca R.C., Moore W., Shuster S.M. (2016) **Invertebrates** *Sinauer Associates, Inc* :639–910
- Buatois L.A., Narbonne G.M., Mangano M.G., Carmona N.B., Myrow P. (2014) **Ediacaran matground ecology persisted into the earliest Cambrian** *Nat Commun* **5**
- Budd G.E. (2001) **Why are arthropods segmented?** *Evolution & Development* **3**:332–342
- Cong P., Ma X., Williams M., Siveter D.J., Siveter D.J., Gabbott S.E., Zhai D., Goral T., Edgecombe G.D., Hou X. (2017) **Host-specific infestation in early Cambrian worms** *Nature Ecology & Evolution* **1**:1465–1469
- Daley A.C., Drage H.B. (2016) **The fossil record of ecdysis, and trends in the moulting behaviour of trilobites** *Arthropod Structure & Development* **45**:71–96
- Daley A.C., Paterson J.R., Edgecombe G.D., García-Bellido D.C., Jago J.B., Donoghue P. (2013) **New anatomical information on Anomalocaris from the Cambrian Emu Bay Shale of South Australia and a reassessment of its inferred predatory habits** *Palaeontology* **56**:971–990
- Erwin D.H. (2020) **The origin of animal body plans: a view from fossil evidence and the regulatory genome** *Development* **147**
- Gelman A., Rubin D.B. (1992) **Inference from iterative simulation using multiple sequences (with discussion)** *Statistical Science* **7**:457–472
- Goloboff P.A., Carpenter J.M., Arias J.S., Esquivel D.R.M. (2008) **Weighting against homoplasy improves phylogenetic analysis of morphological data sets** *Cladistics* **24**:758–773
- Goloboff P.A., Catalano S.A. (2016) **TNT version 1.5, including a full implementation of phylogenetic morphometrics** *Cladistics* **32**:221–238
- Guo J.-F., Li G.-X., Qiang Y.-Q., Song Z.-C., Zhang Z.-F., Han J., Wang W.-Z. (2021) **Watsonella crosbyi from the lower Cambrian (Terreneuvian, Stage 2) Yanjiahe Formation in Three Gorges Area, South China** *Palaeoworld* **30**:1–19
- Han J., Conway Morris S., Ou Q., Shu D.G., Huang H. (2017) **Meiofaunal deuterostomes from the basal Cambrian of Shaanxi (China)** *Nature* **542**:228–231
- Han J., Liu J.N., Zhang Z.F., Zhang X.L., Shu D.G. (2007) **Trunk ornament on the palaeoscolecid worms Cricocosmia and Tabelliscolex from the Early Cambrian Chengjiang deposits of China** *Acta Palaeontologica Polonica* **52**:423–431
- Hanken J., Wake D.B. (1993) **Miniaturization of body size: Organismal consequences and evolutionary significance** *Annual Review of Ecology and Systematics* **24**:501–519

- Hoang D.T., Chernomor O., von Haeseler A., Minh B.Q., Vinh L.S. (2018) **UFBoot2: Improving the Ultrafast Bootstrap Approximation** *Mol Biol Evol* **35**:518–522
- Hou X.G., Ramskold L., Bergstrom J. (1991) **Composition and preservation of the Chengjiang fauna -a Lower Cambrian soft-bodied biota** *Zoologica Scripta* **20**:395–411
- Howard R.J., Edgecombe G.D., Shi X., Hou X., Ma X. (2020) **Ancestral morphology of Ecdysozoa constrained by an early Cambrian stem group ecdysozoan** *BMC Evolutionary Biology* **20**
- Howard R.J. *et al.* (2022) **The Ediacaran origin of Ecdysozoa: integrating fossil and phylogenomic data** *Journal of the Geological Society* **179**:jgs2021–107
- Huang D., Vannier J., Chen J. (2004) **Anatomy and lifestyles of Early Cambrian priapulid worms exemplified by *Corynetis* and *Anningvermis* from the Maotianshan Shale (SW China)** *Lethaia* **37**:21–33
- Kirstensen R.M. (1983) **Loricifera, a new phylum with Aschelminthes characters from the meiobenthos** *Zeitschrift für zoologische Systematik und Evolutionsforschung* **21**:163–180
- Laumer C.E., Fernandez R., Lemer S., Combosch D., Kocot K.M., Riesgo A., Andrade S.C.S., Sterrer W., Sorensen M.V., Giribet G. (2019) **Revisiting metazoan phylogeny with genomic sampling of all phyla** *Proceedings of the Royal Society B* **286**
- Lewis P.O. (2001) **A likelihood approach to estimating phylogeny from discrete morphological character data** *Systematic biology* **50**:913–925
- Liu A.G., Matthews J.J., Menon L.R., McIlroy D., Brasier M.D. (2014) **Haootia quadriformis n. gen., n. sp., interpreted as a muscular cnidarian impression from the Late Ediacaran period (approx. 560 Ma)** *Proc Biol Sci* **281**
- Liu Y., Carlisle E., Zhang H., Yang B., Steiner M., Shao T., Duan B., Marone F., Xiao S., Donoghue P.C.J. (2022) **Saccorhytus is an early ecdysozoan and not the earliest deuterostome** *Nature* **609**:541–546
- Martín-Durán J.M., Hejnol A. (2015) **The study of *Priapulius caudatus* reveals conserved molecular patterning underlying different gut morphogenesis in the Ecdysozoa** *BMC Biology* **13**
- Minh B.Q., Nguyen M.A., von Haeseler A. (2013) **Ultrafast approximation for phylogenetic bootstrap** *Mol Biol Evol* **30**:1188–95
- Moczyłowska M., Budd G.E., Agić H. (2015) **Ecdysozoan-like sclerites among Ediacaran microfossils** *Geological Magazine* **152**:1145–1148
- Nguyen L.T., Schmidt H.A., Von Haeseler A., Minh B.Q. (2015) **IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies** *Molecular biology and evolution* **32**:268–274
- Nielsen C. (2019) **Was the ancestral panarthropod mouth ventral or terminal?** *Arthropod Structure & Development* **49**:152–154
- Ortega-Hernandez J., Janssen R., Budd G.E. (2019) **The last common ancestor of Ecdysozoa had an adult terminal mouth** *Arthropod Structure & Development* **49**:155–158

- Ou Q., Han J., Zhang Z., Shu D., Sun G., Mayer G. (2017) **Three Cambrian fossils assembled into an extinct body plan of cnidarian affinity** *Proceedings of the National Academy of Science* **114**:8835–8840
- Rambaut A., Drummond A., Xie D., Baele G., Suchard M. (2018) **Posterior summarisation in Bayesian phylogenetics using Tracer 1.7** *Systematic Biology* **67**:901–904
- Ronquist F., Teslenko M., van der Mark P., Ayres D.L., Darling A., Hohna S., Larget B., Liu L., Suchard M.A., Huelsenbeck J.P. (2012) **MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space** *Systematic Biology* **61**:539–42
- Rota-Stabelli O., Daley A.C., Pisani D. (2013) **Molecular timetrees reveal a Cambrian colonization of land and a new scenario for ecdysozoan evolution** *Current Biology* **23**:392–8
- Sawaki Y., Nishizawa M., Suo T., Komiya T., Hirata T., Takahata N., Sano Y., Han J., Kon Y., Maruyama S. (2008) **Internal structures and U–Pb ages of zircons from a tuff layer in the Meishucunian formation, Yunnan Province, South China** *Gondwana Research* **14**:148–158
- Schmidt-Rhaesa A. (2007) **The evolution of organ systems** :54–73
- Schmidt-Rhaesa A. (2013) **Nematomorpha. In: A. Schmidt-Rhaesa (Editor), Gastrotricha, Cycloneuralia and Gnathifera** *De Gruyter, Germany* :29–146
- Schmidt-Rhaesa A. (2013) **Priapulida. In: A. Schmidt-Rhaesa (Editor), Gastrotricha, Cycloneuralia and Gnathifera** *De Gruyter, Germany* :147–180
- Schmidt-Rhaesa A. (2014) **Handbook of Zoology. Gastrotricha, Cycloneuralia and Gnathifera, 2** *De Gruyter, Germany* :5–12
- Shu D.G., Han J. (2020) **The core value of Chengjiang fauna: the formation of the animal kingdom and the birth of basic human organs** *Earth Science Frontiers* **27**:382–412
- Shu D.G., Han J. (2020) **The core value of Chengjiang fauna: the information of animal kingdom and the birth basic human organs** *Earth Science Frontiers* **27**:1–32
- Valentine J.W., Collins A.G. (2000) **The significance of moulting in ecdysozoan evolution** *Evolution & Development* **2**:152–156
- Vannier J., Calandra I., Gaillard C., Żylińska A. (2010) **Priapulid worms: Pioneer horizontal burrowers at the Precambrian-Cambrian boundary** *Geology* **38**:711–714
- Vannier J., Chen J. (2005) **Early Cambrian Food Chain: New Evidence from Fossil Aggregates in the Maotianshan Shale Biota, SW China** *Palaio* **20**:3–26
- Vermeij G.J. (1977) **The Mesozoic marine revolution: evidence from snails, predators and grazer** *Paleobiology* **3**:245–258
- Vinther J., Parry L.A. (2019) **Bilateral jaw elements in Amiskwia sagittiformis bridge the morphological gap between gnathiferans and chaetognaths** *Current Biology* **29**:881–888
- Wang D., Vannier J., Schumann I., Wang X., Yang X.G., Komiya T., Uesugi K., Sun J., Han J. (2019) **Origin of ecdysis: fossil evidence from 535-million-year-old scalidophoran worms** *Proceedings of the Royal Society B* **286**



Zhang H., Xiao S., Liu Y., Yuan X., Wan B., Muscente A.D., Shao T., Gong H., Cao G. (2015) **Armored kinorhynch-like scalidophoran animals from the early Cambrian** *Scientific Reports* 5

Zhang Z., Smith M.R., Shu D. (2015) **New reconstruction of the Wiwaxia scleritome, with data from Chengjiang juveniles** *Sci Rep* 5

## Article and author information

### Deng Wang

State Key Laboratory of Continental Dynamics, Shaanxi Key Laboratory of Early Life & Environments and Department of Geology, Northwest University; Xi'an, China  
ORCID iD: [0000-0002-4464-9632](https://orcid.org/0000-0002-4464-9632)

### Yaqin Qiang

School of Earth Science and Resources, Key Laboratory of Western China's Mineral Resources and Geological Engineering, Ministry of Education, Chang'an University; Xi'an, China

### Junfeng Guo

School of Earth Science and Resources, Key Laboratory of Western China's Mineral Resources and Geological Engineering, Ministry of Education, Chang'an University; Xi'an, China  
**For correspondence:** [junfengguo@chd.edu.cn](mailto:junfengguo@chd.edu.cn)

### Jean Vannier

Université de Lyon, Université Claude Bernard Lyon 1, ENS de Lyon, CNRS, Laboratoire de Géologie de Lyon: Terre, Planètes, Environnement (CNRS-UMR 5276), Villeurbanne 69622, France  
ORCID iD: [0000-0003-0998-1231](https://orcid.org/0000-0003-0998-1231)

### Zuchen Song

School of Earth Science and Resources, Key Laboratory of Western China's Mineral Resources and Geological Engineering, Ministry of Education, Chang'an University; Xi'an, China

### Jiaxin Peng

School of Earth Science and Resources, Key Laboratory of Western China's Mineral Resources and Geological Engineering, Ministry of Education, Chang'an University; Xi'an, China

### Boyao Zhang

School of Earth Science and Resources, Key Laboratory of Western China's Mineral Resources and Geological Engineering, Ministry of Education, Chang'an University; Xi'an, China

### Jie Sun

State Key Laboratory of Continental Dynamics, Shaanxi Key Laboratory of Early Life & Environments and Department of Geology, Northwest University; Xi'an, China, School of Earth Science and Resources, Key Laboratory of Western China's Mineral Resources and Geological Engineering, Ministry of Education, Chang'an University; Xi'an, China

### Yilun Yu

University of Chinese Academy of Sciences; Beijing, China, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences; Beijing, China

**Yiheng Zhang**

School of Information Science and Technology, Northwest University; Xi'an, China  
ORCID iD: [0009-0002-2426-0838](https://orcid.org/0009-0002-2426-0838)

**Tao Zhang**

School of Information Science and Technology, Northwest University; Xi'an, China

**Xiaoguang Yang**

State Key Laboratory of Continental Dynamics, Shaanxi Key Laboratory of Early Life & Environments and Department of Geology, Northwest University; Xi'an, China

**Jian Han**

State Key Laboratory of Continental Dynamics, Shaanxi Key Laboratory of Early Life & Environments and Department of Geology, Northwest University; Xi'an, China

**For correspondence:** [elihanj@nwu.edu.cn](mailto:elihanj@nwu.edu.cn)

ORCID iD: [0000-0002-2134-4078](https://orcid.org/0000-0002-2134-4078)

**Copyright**

© 2024, Wang et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

**Editors**

Reviewing Editor

**Shigehiro Kuraku**

National Institute of Genetics, Mishima, Japan

Senior Editor

**Claude Desplan**

New York University, New York, United States of America

**Reviewer #1 (Public Review):**

Summary:

Wang and co-workers characterise the fossil of *Beretella spinosa* from the early Cambrian, Yanjiahe Formation, South China. Combining morphological analyses with phylogenetic reconstructions, the authors conclude that *B. spinosa* is closely related to *Saccorhytus*, an enigmatic fossil recently ascribed to Ecdysozoa, or moulting animals, as an extinct "basal" lineage. Finding additional representatives of the clade Saccorhytida strengthens the idea that there existed a diversity of body plans previously underappreciated in Ecdysozoa, which may have implications for our understanding of the earliest steps in the evolution of this major animal group.

Strengths:

I'm not a paleobiologist; therefore, I cannot give an expert opinion on the descriptions of the fossils. However, the similarities with *Saccorhytus* seem evident, and the phylogenetic reconstructions are adequate. Evolutionary interpretations are generally justified, and the consolidation of Saccorhytida as the extinct sister lineage to extant Ecdysozoans will have significant implications for our understanding of this major animal clade.

**Weaknesses:**

While I generally agree with the author's interpretations, the idea of Saccorhytida as a divergent, simplified off-shot is slightly contradictory with a probably non-vermiform ecdysozoan ancestor. The author's analyses do not discard the possibility of a vermiform ecdysozoan ancestor (importantly, Supplementary Table 4 does not reconstruct that character), and outgroup comparison with Spiralia (and even Deuterostomia for Protostomia as a whole) indicates that a more or less anteroposteriorly elongated (i.e., vermiform) body is likely common and ancestral to all major bilaterian groups, including Ecdysozoa. Indeed, Figure 4b depicts the potential ancestor as a "worm". The authors argue that the simplification of Saccorhytida from a vermiform ancestor is unlikely "because it would involve considerable anatomical transformations such as the loss of vermiform organization, introvert, and pharynx in addition to that of the digestive system". However, their data support the introvert as a specialisation of Scalidophora (Figure 4a and Supplementary Table 4), and a pharyngeal structure cannot be ruled out in Saccorhytida. Likewise, loss of an anus is not uncommon in Bilateria. Moreover, this can easily become a semantics discussion (to what extent can an animal be defined as "vermiform"? Where is the limit?). Therefore, I suggest to leave the evolutionary scenario more open. Supporting Saccorhytida as a true group at the early steps of Ecdysozoa evolution is important and demonstrates that animal body plans are more plastic than previously appreciated. However, with the current data, it is unlikely that Saccorhytida represents the ancestral state for Ecdysozoa (as the authors admit), and a vermiform nature is not ruled out (and even likely) in this animal group. Suggesting that the ancestral Ecdysozoan might have been small and meiobenthic is perhaps more interesting and supported by the current data (phylogeny and outgroup comparison with Spiralia).

<https://doi.org/10.7554/eLife.94709.1.sa2>

**Reviewer #2 (Public Review):****Summary:**

This work provides important anatomical features of a new species from the Lower Cambrian, which helps advance our understanding of the evolutionary origins of animal body plans. The authors interpreted that the new species possessed a bilateral body covered with cuticular polygonal reticulation and a ventral mouth. Based on cladistic analyses using maximum likelihood, Bayesian, and parsimony, the new species was placed, along with Saccorhytus, in a sister group ("Saccorhytida") of the Ecdysozoa. The phylogenetic position of Saccorhytida suggests a new scenario of the evolutionary origin of the crown ecdysozoan body plan.

**Strengths:**

Although the new species reported in this paper show strange morphologies, the interpretation of anatomical features was based on detailed observations of multiple fossil specimens, thereby convincing at the moment. Morphological data about fossil taxa in the Ediacaran and Early Cambrian are quite important for our understanding of the evolution of body plans (and origins of phyla) in paleontology and evolutionary developmental biology, and this paper represents a valuable contribution to such research fields.

**Weaknesses:**

The preservations of the specimens, in particular on the putative ventral side, are not good, and the interpretation of the anatomical features needs to be tested with additional specimens in the future. The monophyly of Cycloneuralia (Nematoida + Scalidophora) was not necessarily well-supported by cladistic analyses, and the evolutionary scenario (Figure 4) also needs to be tested in future works.

### Reviewer #3 (Public Review):

#### Summary:

The authors of this manuscript identified the fossils of the newly designated species *Beretella spinosa* and analyzed its phylogenetic position in relation to the extinct described species and extant species. Their analysis placed the newly described species *Beretella spinosa* and *Saccorhytus* as an independent clade from the rest of the ecdysozoans. Remarkably, these species are non-vermiform, and the resulting evolutionary scenario assumes non-vermiform as early ecdysozoans.

#### Strengths:

The study presents outstanding, novel data and provides new insights into the evolution of animal forms especially regarding their morphological diversity after the Cambrian explosion.

#### Weaknesses:

I, as a paleontology non-expert, experienced several difficulties in reading the manuscript. This should be taken into consideration when assuming a wide range of readers including non-experts.

<https://doi.org/10.7554/eLife.94709.1.sa0>

### Author Response

#### Public reviews:

##### Reviewer 1:

#### Weaknesses:

*While I generally agree with the author's interpretations, the idea of Saccorhytida as a divergent, simplified off-shot is slightly contradictory with a probably non-vermiform ecdysozoan ancestor. The author's analyses do not discard the possibility of a vermiform ecdysozoan ancestor (importantly, Supplementary Table 4 does not reconstruct that character),*

Reply: Thanks for the comments. Saccorhytids are only known from the early Cambrian and their unique morphology has no equivalent among any extinct or extant ecdysozoan groups. This prompted us to consider them as a possible dead-end evolutionary off-shot. The nature of the last common ancestor of ecdysozoan (i.e. a vermiform or non-vermiform animal with capacities to renew its cuticle by molting) remains hypothetical. At present, palaeontological data do not allow us to resolve this question. The animal in Fig. 4b at the base of the tree is supposed to represent an ancestral soft-bodied form with no cuticle from which ecdysozoan evolved via major innovations (cuticular secretion and ecdysis). Its shape is hypothetical as indicated by a question mark. Our evolutionary model is clearly intended to be tested by further studies and hopefully new fossil discoveries.

*and outgroup comparison with Spiralia (and even Deuterostomia for Protostomia as a whole) indicates that a more or less anteroposteriorly elongated (i.e., vermiform) body is likely common and ancestral to all major bilaterian groups, including Ecdysozoa. Indeed, Figure 4b depicts the potential ancestor as a "worm". The authors argue that the simplification of Saccorhytida from a vermiform ancestor is unlikely "because it would*



*involve considerable anatomical transformations such as the loss of vermiform organization, introvert, and pharynx in addition to that of the digestive system". However, their data support the introvert as a specialisation of Scalidophora (Figure 4a and Supplementary Table 4), and a pharyngeal structure cannot be ruled out in Saccorhytida. Likewise, loss of an anus is not uncommon in Bilateria. Moreover, this can easily become a semantics discussion (to what extent can an animal be defined as "vermiform"? Where is the limit?).*

Reply: We agree with you that “vermiform” is an ill-defined term that should be avoided. “Elongated” might be a better term to designate the elongation of the body along the antero-posterior axis. Changes have been made in the text to solve this semantic problem. Priapulid worms or annelids are examples of extremely elongated, tubular animals. In saccorhytids, the antero-posterior elongation is present (as it is in the vast majority of bilaterians) but extremely reduced, Saccorhytus and Beretella having a sac-like or beret-shape, respectively. That such forms may have derived from elongated, tubular ancestors (e.g. comparable with scalidophoran worms) would require major anatomical transformations that have no equivalent among modern animals. We agree that further speculation about the nature of these transformations is unnecessary and should be deleted simply because the nature of these ancestors is purely hypothetical. We also agree that the loss of anus and the extreme simplification of the digestive system is common among extant bilaterians. The single opening seen in Saccorhytus and possibly Beretella may result from a comparable simplification process. In Figure 4b, the hypothetical pre-ecdysozoan animal is slightly elongated (antero-posterior axis and polarity) but in no way comparable with a very elongated and cylindrical ecdysozoan worm (e.g. extant or extinct priapulid).

*Therefore, I suggest to leave the evolutionary scenario more open. Supporting Saccorhytida as a true group at the early steps of Ecdysozoa evolution is important and demonstrates that animal body plans are more plastic than previously appreciated. However, with the current data, it is unlikely that Saccorhytida represents the ancestral state for Ecdysozoa (as the authors admit), and a vermiform nature is not ruled out (and even likely) in this animal group. Suggesting that the ancestral Ecdysozoan might have been small and meiobenthic is perhaps more interesting and supported by the current data (phylogeny and outgroup comparison with Spiralia).*

Reply: We agree the evolutionary scenario should be more open, especially the evolutionary process that gave rise to Saccorhytida. Again, we know nothing about the morphology of the ancestral ecdysozoan (typically the degree of body elongation, whether it had a differentiated introvert or not, whether it had a through gut or not). Simplification appears as one possible option, but which assumes that the ancestral ecdysozoan was an elongated animal with a through gut. Changes will be made in Fig.4A accordingly. Alternatively, the ancestral ecdysozoan might have been small and meiobenthic.

#### **Reviewer 2:**

##### *Weaknesses:*

*The preservations of the specimens, in particular on the putative ventral side, are not good, and the interpretation of the anatomical features needs to be tested with additional specimens in the future. The monophyly of Cycloneuralia (Nematoida + Scalidophora) was not necessarily well-supported by cladistic analyses, and the evolutionary scenario (Figure 4) also needs to be tested in future works.*

Reply: Yes, we agree that our MS is the first report on an enigmatic ecdysozoan. Whereas the dorsal side of the animal is well documented (sclerites), uncertainties remain concerning its ventral anatomy (typically the mouth location and shape). Additional better-preserved

specimens will hopefully provide the missing information. Concerning Cycloneuralia, their monophyly is generally better supported by analyses based on morphological characters than in molecular phylogenies. I

**Reviewer 3:**

*Weaknesses: I, as a paleontology non-expert, experienced several difficulties in reading the manuscript. This should be taken into consideration when assuming a wide range of readers including non-experts.*

Reply: We have ensured that the text is comprehensible to biologists. Our main results are summarized in relatively simple diagrams (e.g. Fig. 4). We are aware that technical descriptive terms may appear obscure to non-specialists. However, we think that our text-figures help the reader to understand the morphology of these ancient animals.