Genetic Relatedness

Magnus Dehli Vigeland


Solutions for exercise set I

Note: QuickPed usually gives the quickest solution for these exercises. R code is provided here.


library(pedsuite)

Exercise I-1


a) Grandaunt – grandnephew:



b) First cousins twice removed:




c) Half second cousins once removed:



Exercise I-2

- a) (Omitted)
- b) Adrian and Belinda are simultaneous half siblings and half first cousins.
- c) Solution in R:

```
x = halfSibStack(2) |> swapSex(8)
plot(x, hatched = leaves, labs = c(Adrian = 7, Belinda = 8))
```



d) No, there is no inbreeding in this pedigree.


Exercise I-3

Here is a way to recreate the pedigree in R:

Exercise I-4

a) Solution in R:

```
x = halfSibPed(nch1 = 1, nch2 = 2, sex1 = 1, sex2 = 2:1) >
addSon(4:5)
plot(x)
```



b) 6 is both an uncle and half-uncle of 7. In R, this can be verified using the verbalise function:

```
verbalise(x)
```

```
## Avuncular: 6 is an uncle of 7
## 6-[2,3]-5-7
## Half-avuncular: 6 is a half-uncle of 7
## 6-[2]-4-7
```

c) (Omitted)

Exercise I-5

Omitted; the answer is given in the exercise.

Exercise I-6

Quadruple first cousins!

```
x = nuclearPed(4, sex = c(1,2,1,2)) |>
   addSon(3:4) |>
   addDaughter(5:6)
plot(x, hatched = 7:8)
verbalise(x, 7:8)
## Quadruple first cousins
## 7-3-[1,2]-5-8
## 7-3-[1,2]-5-8
## 7-4-[1,2]-5-8
## 7-4-[1,2]-6-8
```