

## Foundations of Chabauty - Kim

$X/\mathbb{Q}$  sm. proj. curve of genus  $g \geq 2$

Thm: (Faltings 1983)

$$\#X(\mathbb{Q}) < \infty$$

Q: ("Effective Mordell")

- How to find  $X(\mathbb{Q})$ ?

Unknown whether a general algorithm exists.

- Bounds on  $\#X(\mathbb{Q})$ ?

↳ Uniformity Conjecture: (Caporaso-Harris-Mazur)

$$\exists N(g) \text{ s.t. } \#X(\mathbb{Q}) < N(g) \text{ for all } X \text{ of genus } g$$

Chabauty - Kim theory:  $p$ -adic approach addressing these, still largely conjectural

### 1. Method of Chabauty (1941) and Coleman (1985)

$J := \text{Jac}_X$  - ab. var /  $\mathbb{Q}$  of dim.  $g$

$J(\mathbb{Q})$  is f.g. ab. gp. (Mordell-Weil)

$$r := \text{rk}_{\mathbb{Z}} J(\mathbb{Q}) \quad \rightarrow \quad J(\mathbb{Q}) \cong \mathbb{Z}^r \oplus (\text{finite})$$

Assume  $X(\mathbb{Q}) \neq \emptyset$  and fix  $b \in X(\mathbb{Q})$ .

$$\rightarrow \text{Abel-Jacobi map } AJ: X \rightarrow J, \quad P \mapsto [P] - [b]$$

Choose auxiliary prime  $p$  of good reduction for  $X$ .

Have commutative diagram

$$\begin{array}{ccc} X(\mathbb{Q}) & \hookrightarrow & X(\mathbb{Q}_p) \\ AJ \downarrow & & \downarrow AJ_p \\ J(\mathbb{Q}) & \hookrightarrow & J(\mathbb{Q}_p) \end{array}$$

$$\Rightarrow X(\mathbb{Q}) \subseteq AJ_p^{-1}(J(\mathbb{Q})) \subseteq X(\mathbb{Q}_p)$$

Idea: find functions  $0 \neq f: \mathcal{J}(\mathbb{Q}_p) \rightarrow \mathbb{Q}_p$  which vanish on  $\mathcal{J}(\mathbb{Q})$ ,  
 then  $f \circ A\mathcal{J}_p: X(\mathbb{Q}_p) \rightarrow \mathbb{Q}_p$  vanishes on  $X(\mathbb{Q})$  2

This works if  $\boxed{r < g}$ . (\*)

Coleman integration: Fix  $\omega \in H^0(X_{\mathbb{Q}_p}, \Omega^1) \cong H^0(\mathcal{J}_{\mathbb{Q}_p}, \Omega^1)$ .

$\omega$  translation-inv. on  $\mathcal{J}_{\mathbb{Q}_p} \Rightarrow \exists!$  anti-derivative  $F_\omega: \mathcal{J}(\mathbb{Q}_p) \rightarrow \mathbb{Q}_p$  s.t.

- $F_\omega$  is a homom.
- near  $0 \in \mathcal{J}(\mathbb{Q}_p)$ ,  $F_\omega$  is given by a convergent power series with  $dF_\omega = \omega$

Write  $\int_0^D \omega := F_\omega(D)$ ,  $\int_P^Q \omega := \int_0^{[Q]-[P]} \omega$  for  $P, Q \in X(\mathbb{Q}_p)$ .

Get homom.  $\log: \mathcal{J}(\mathbb{Q}_p) \rightarrow H^0(\mathcal{J}_{\mathbb{Q}_p}, \Omega^1)^\vee$ ,  $D \mapsto \left[ \omega \mapsto \int_0^D \omega \right]$ .

If  $r < g \Rightarrow \log \overline{\mathcal{J}(\mathbb{Q})} \in H^0(\mathcal{J}_{\mathbb{Q}_p}, \Omega^1)^\vee$   $\mathbb{Z}_p$ -submodule of rank  $r < g$   
 $\Rightarrow \exists 0 \neq \omega \in H^0(\mathcal{J}_{\mathbb{Q}_p}, \Omega^1)$  s.t.  $F_\omega$  vanishes on  $\mathcal{J}(\mathbb{Q})$

$\Rightarrow F_\omega \circ A\mathcal{J}_p: X(\mathbb{Q}_p) \rightarrow \mathbb{Q}_p$ ,  $P \mapsto \int_b^P \omega$  vanishes on  $X(\mathbb{Q})$

$P \mapsto \int_b^P \omega$  locally analytic (given by convergent power series on residue discs)

$\Rightarrow$  only finitely many zeroes

$\Rightarrow X(\mathbb{Q})$  finite

Also get explicit bounds, e.g.  $\#X(\mathbb{Q}) \leq \#X(\mathbb{F}_p) + (2g-2)$  if  $p > 2g$ .

Problem: method fails if  $r \not< g$

Kim (2005): developed non-abelian generalisation to overcome this.

Central object:  $\mathbb{Q}_p$ -pro-unipotent étale fundamental group  $U^{\text{ét}}$  of  $X$

Chabauty-Kim diagram:

$$\begin{array}{ccc}
 X(\mathbb{Q}) & \longrightarrow & X(\mathbb{Q}_p) \\
 \downarrow j & & \downarrow j_p \\
 \text{Sel}_\infty(X) & \xrightarrow{\text{loc}_p} & H_f^1(G_p, U^{\text{ét}}) \xrightarrow[\sim]{\log_{\mathbb{Z}_p^k}} F^0 \setminus U^{\text{ét}}
 \end{array}
 \quad (\text{CK})$$

Properties:

- bottom row: affine  $\mathbb{Q}_p$ -schemes, maps are algebraic
- image of each residue disc  $\text{red}^{-1}(y) \subseteq X(\mathbb{Q}_p)$ ,  $y \in X(\mathbb{F}_p)$ , under  $j^{dR}$  is Zariski-dense
- Conjecturally,  $\text{loc}_p$  is non-dominant ( $\Leftarrow$  Bloch-Kato conj.)

Method works as follows:

If  $\text{loc}_p$  non-dominant  $\Rightarrow$  find  $0 \neq f \in \mathcal{O}(F^\circ \setminus U^{dR})$  vanishing on

image of  $H_f^1(G_{\mathbb{Q}}, U^{\text{ét}})$

$\Rightarrow f \circ j^{dR}: X(\mathbb{Q}_p) \rightarrow \mathbb{Q}_p$  non-trivial + analytic on residue discs

(given by iterated Coleman integrals)

$\Rightarrow V(f \circ j^{dR}) \subseteq X(\mathbb{Q}_p)$  finite set containing  $X(\mathbb{Q})$

Def: Chabauty-Kim locus

$$X(\mathbb{Q}_p)_\infty := j_p^{-1}(\text{loc}_p(\text{Sel}_\infty(X))) = \bigcap_{f \text{ as above}} V(f \circ j^{dR}) \subseteq X(\mathbb{Q}_p).$$

Note:  $X(\mathbb{Q}) \subseteq X(\mathbb{Q}_p)_\infty$

Conj.:  $X(\mathbb{Q}_p)_\infty$  is finite ( $\Leftarrow$  Bloch-Kato)

Conj.: (Kim's conjecture)

$$X(\mathbb{Q})_\infty = X(\mathbb{Q}).$$

Rmks: • replace  $U^{\text{ét}}$  with  $n$ -step nilpotent quotient  $U_n^{\text{ét}} \rightsquigarrow X(\mathbb{Q}_p)_n$

• roughly: "n=1"  $\rightarrow$  Chabauty-Coleman

"n=2"  $\rightarrow$  quadratic Chabauty

• variant for  $S$ -integral points on affine hyperbolic curves

2. Prounipotent étale fundamental group

$X/\mathbb{Q}$ ,  $b \in X(\mathbb{Q})$  as above

$\bar{\mathbb{Q}}/\mathbb{Q}$  alg. closure,  $G_{\mathbb{Q}} := \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$

$\leadsto \pi_1^{\text{ét}}(X_{\bar{Q}}, b)$  - "étale fundamental group"

It is a profinite group with cts.  $G_{\bar{Q}}$ -action.

$\text{Cov}(X_{\bar{Q}})$  - category of finite étale covers  $Y \rightarrow X_{\bar{Q}}$

$F_b: \text{Cov}(X_{\bar{Q}}) \rightarrow \text{FinSet}, (Y \xrightarrow{f} X_{\bar{Q}}) \mapsto f^{-1}(b)$  fibre functor

$\pi_1^{\text{ét}}(X_{\bar{Q}}, b) := \text{Aut}(F_b)$ , acts compatibly on  $f^{-1}(b)$  for all  $(Y \xrightarrow{f} X_{\bar{Q}}) \in \text{Cov}(X_{\bar{Q}})$

Comparison with topological  $\pi_1$ : (via Riemann's existence theorem)

$$\pi_1^{\text{ét}}(X_{\bar{Q}}, b) = \text{profinite completion of } \pi_1^{\text{top}}(X(\mathbb{C}), b)$$

$$\cong \langle a_1, b_1, \dots, a_g, b_g \mid \prod [a_i, b_i] = 1 \rangle^{\wedge}$$

Mal'cev completion:  $\Pi$  profinite gp.  $p$  prime

The  $\mathbb{Q}_p$ -pro-unipotent completion of  $\Pi$  is the universal pro-unipotent algebraic gp  $\Pi^{\mathbb{Q}_p}$  over  $\mathbb{Q}_p$  with a continuous homom.

$$\Pi \longrightarrow \Pi^{\mathbb{Q}_p}(\mathbb{Q}_p).$$

Think:  $\Pi^{\mathbb{Q}_p} = " \mathbb{Q}_p \otimes_{\mathbb{Z}} \Pi "$ .

Def:  $\pi_1^{\text{ét}, \mathbb{Q}_p}(X_{\bar{Q}}, b) := \pi_1^{\text{ét}}(X_{\bar{Q}}, b)^{\mathbb{Q}_p}$

" $\mathbb{Q}_p$ -pro-unipotent étale fundamental group"

Alternative definition via Tannakian formalism

$\text{Loc}_{\mathbb{Q}_p}^{\text{un}}(X_{\bar{Q}}) :=$  category of unipotent  $\mathbb{Q}_p$ -local systems on  $X_{\bar{Q}}$ ,

i.e. étale-locally constant sheaves of  $\mathbb{Q}_p$ -v.s. on  $X_{\bar{Q}}$  ét which

admit a filtration  $0 \leq \text{Fil}_0 E \leq \dots \leq \text{Fil}_m E = E$  s.t.

$\text{Fil}_i E / \text{Fil}_{i-1} E \cong$  direct sum of copies of  $\mathbb{Q}_p$ .

$\text{Fib}_b: \text{Loc}_{\mathbb{Q}_p}^{\text{un}}(X_{\bar{Q}}) \rightarrow \mathbb{Q}_p\text{-Vect}, E \mapsto E(b)$  fibre functor, preserves  $\otimes$

$\pi_1^{\text{ét}, \mathbb{Q}_p}(X_{\bar{Q}}, b) := \underline{\text{Aut}}^{\otimes}(\text{Fib}_b)$  tensor-preserving automorphisms

Notation:  $U^{\text{ét}} := \pi_1^{\text{ét}, \mathbb{Q}_p}(X_{\bar{Q}}, b)$ ,  $U_n^{\text{ét}} := n$ -step nilpotent quotient

$G_{\bar{Q}}$  acts on  $U_{(n)}^{\text{ét}}$ . Fix  $G_{\bar{Q}}$ -equivariant quotient  $U^{\text{ét}} \twoheadrightarrow U'$  e.g.  $U' = U_n^{\text{ét}}$ .

Path torsors: For  $x \in X(\mathbb{Q})$  have

$$P^{\text{ét}}(x) := \pi_1^{\text{ét}, \mathbb{Q}_p}(X_{\bar{\mathbb{Q}}}; b, x) := \underline{\text{Isom}}^{\otimes}(Fib_b, Fib_x) \quad \text{"path torsor"}$$

It is a  $G_{\mathbb{Q}}$ -equivariant  $U^{\text{ét}}$ -torsor over  $\mathbb{Q}_p$ .

Form pushout along  $U^{\text{ét}} \rightarrow U'$

$$\leadsto P'(x) := P^{\text{ét}}(x) / \ker(U^{\text{ét}} \rightarrow U'),$$

$G_{\mathbb{Q}}$ -equiv.  $U'$ -torsor

Can do the same locally for  $\mathbb{Q}_l$ -points ( $l$  any prime):

Fix  $\bar{\mathbb{Q}}_l / \mathbb{Q}_l$  with embedding  $\bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_l$

$\leadsto$  local Galois group  $G_l := \text{Gal}(\bar{\mathbb{Q}}_l / \mathbb{Q}_l) \leq G_{\mathbb{Q}}$

Have  $\pi_1^{\text{ét}, \mathbb{Q}_p}(X_{\bar{\mathbb{Q}}_l}, b) = \pi_1^{\text{ét}, \mathbb{Q}_p}(X_{\bar{\mathbb{Q}}}, b)$ .

Given  $x \in X(\mathbb{Q}_l) \rightarrow G_l$ -equivariant  $U'$ -torsor  $P'(x)$

### 3. Spaces of torsors

$G$  profinite gp,  $U / \mathbb{Q}_p$  pro-unipotent group,  $G \curvearrowright U$

Def:  $R$   $\mathbb{Q}_p$ -algebra. A  $G$ -equivariant  $U$ -torsor  $P$  over  $R$

is an  $R$ -scheme  $P$  with a  $G$ -action and a  $G$ -equivariant right  $U$ -action  $P \times U \rightarrow U$  such that  $P \rightarrow \text{Spec}(R)$  is faithfully flat and the map

$$P \times U \rightarrow P \times_R P, \quad (y, u) \mapsto (y, yu)$$

is an isomorphism.

Remark:  $U$  pronipotent implies that all  $U$ -torsors are trivial as mere torsors, but not necessarily  $G$ -equivariantly.

$c: G \rightarrow U(R)$  is a cocycle if  $c(\sigma\tau) = \sigma(c(\tau)) \cdot c(\sigma) \quad \forall \sigma, \tau \in G$ .

$c$  and  $c'$  are cohomologous ( $c \sim c'$ ) if  $\exists u \in U(R)$  s.t.  $c'(\sigma) = u^{-1} c(\sigma) \sigma u \quad \forall \sigma \in G$

$$H^1(G, U(R)) := \{\text{cocycles } G \rightarrow U(R)\} / \sim$$

Cohomology functor on  $\mathbb{Q}_p$ -algebras:

$$H^1(G, U) : R \mapsto H^1(G, U(R))$$

Prop:  $\left( \begin{array}{c} G\text{-equivariant } U\text{-torsors} \\ \text{over } R \end{array} \right) / \text{iso} \cong H^1(G, U(R))$

Pf:  $P$   $G$ -equiv.  $U$ -torsor

$U$  pro-unipotent  $\Rightarrow P(R) \neq \emptyset$

Choose  $p_0 \in P(R)$ . For  $\sigma \in G$   $\exists!$   $c(\sigma) \in U(R)$  s.t.  $\sigma p_0 = p_0 \cdot c(\sigma)$ .

Check:  $c: G \rightarrow U(R)$  is cocycle

• different choice of  $p_0$  gives cohomologous cocycle

Conversely, given  $c: G \rightarrow U(R)$  define torsor  $P_c := U$  with

$c$ -twisted  $G$ -action  $G \times U \rightarrow U, (\sigma, u) \mapsto \sigma u \cdot c(\sigma)$ . □

Rmk:  $1 \rightarrow A \rightarrow U \rightarrow U' \rightarrow 1$  central extension induces long exact sequence of pointed sets ending at  $H^2(G, A)$

Thm: Assume  $U$  admits a separated  $G$ -stable filtration

$$U = W_{-1}U \supseteq W_{-2}U \supseteq \dots$$

with  $[W_{-i}U, W_{-j}U] \subseteq W_{-i-j}U \quad \forall i, j \geq 1$ . Write  $V_n := \text{gr}_{-n}^W U$ .

Note  $V_n$  is a  $\mathbb{Q}_p$ -v.s. Assume:

(1)  $H^0(G, V_n) = 0$

(2)  $H^1(G, V_n)$  is fin. dim.

for all  $n \geq 1$ . Then  $H^1(G, U)$  is representable by an affine

$\mathbb{Q}_p$ -scheme which is non-canonically isomorphic to a closed subscheme of  $\prod_{n \geq 1} H^1(G, V_n)$ .

Proof of representability: (Sketch)

Let  $U_n := U/W_{-(n+1)}U$ , then  $U = \varprojlim_n U_n$ .

$H^1(G, U) \xrightarrow{\sim} \varprojlim_n H^1(G, U_n)$  iso  $\Rightarrow$  reduced to  $H^1(G, U_n)$

Use induction on  $n$ .

Case  $n=1$ :  $U_1 = V_1$  vector group,  $U_1(\mathbb{R}) = V_1 \otimes \mathbb{R}$ .

Show  $H^1(G, V_1) \otimes \mathbb{R} \rightarrow H^1(G, V_1 \otimes \mathbb{R})$  iso.

$\Rightarrow H^1(G, V_1)$  represented by vector scheme  $H^1(G, V_1)$

Inductive step  $n \rightarrow n+1$ .

Central extension

$$1 \rightarrow V_n \rightarrow U_n \rightarrow U_{n-1} \rightarrow 1$$

$\Rightarrow$  LES of functors on  $\mathbb{Q}_p$ -algebras

$$\begin{array}{c}
 1 \rightarrow H^0(G, V_n) \rightarrow H^0(G, U_n) \rightarrow H^0(G, U_{n-1}) \rightarrow \\
 \searrow \hspace{10em} \searrow \hspace{10em} \searrow \hspace{10em} \\
 \hookrightarrow H^1(G, V_n) \rightarrow H^1(G, U_n) \rightarrow H^1(G, U_{n-1}) \xrightarrow{\delta^1} \\
 \searrow \hspace{10em} \searrow \hspace{10em} \searrow \hspace{10em} \\
 \hookrightarrow H^2(G, V_n)
 \end{array}$$

- $H^1(G, U_{n-1})$  and  $H^1(G, V_n)$  repr.  
 $\Rightarrow K := \ker(\delta^1)$  repr. by affine  $\mathbb{Q}_p$ -scheme
- $H^1(G, U_n) \rightarrow K$  is surjective map of functors  
 $K = \text{Spec}(S)$ , then  $H^1(G, U_n)(S) \rightarrow K(S) = \text{Hom}(S, S) \ni \text{id}_S$  surj.  
 $\rightarrow$  get splitting  $s: K \rightarrow H^1(G, U_n)$
- $H^1(G, U_n) \rightarrow K$  is  $H^1(G, V_n)$ -torsor  
 $\Rightarrow K \times H^1(G, V_n) \xrightarrow{\sim} H^1(G, U_n)$  iso  
 $\Rightarrow H^1(G, U_n)$  representable

Cor:  $X/\mathbb{Q}$ ,  $U^{\text{ét}} \rightarrow U'$  as before,  $l$  prime

$$\Rightarrow R \mapsto \left( \begin{array}{c} G_l\text{-equiv. } U'\text{-torsors} \\ \text{over } R \end{array} \right) /_{\text{iso}} \cong H^1(G_l, U'(R))$$

is representable by affine  $\mathbb{Q}_p$ -scheme.

Denote it by  $H^1(G_l, U')$ .

Pf: (for  $l \neq p$ )

Check conditions of Theorem.

Define  $W_n U'$  via descending central series:

$$W_{-1} U' := U', \quad W_{-i+1} U' := [W_{-i} U', U']$$

$V_n := \text{gr}_n^W U'$ . Claim:  $V_n$  is pure of weight  $-n$ , i.e., the eigenvalues  $\alpha$  of any geom. Frobenius  $F \in G_p$  are alg. /  $\mathbb{Q}$  and  $|\iota(\alpha)| = l^{-n/2}$  for all embeddings  $\iota: \mathbb{Q}(\alpha) \hookrightarrow \mathbb{C}$ . Follows from:

- $V_1 = (U')^{\text{ab}} \leftarrow U^{\text{ét, ab}} = H_{\text{ét}}^1(X_{\overline{\mathbb{Q}_l}}, \mathbb{Q}_p)^*$ , pure of wt  $-1$  by Riemann hypothesis for curves over finite fields
- $V_1^{\otimes n} \rightarrow V_n$ ,  $v_1 \otimes \dots \otimes v_n \mapsto [v_1, [v_2, [\dots, v_n]] \dots]$   
 $\Rightarrow V_n$  pure of wt  $-n$

$$\text{Claim} \Rightarrow H^0(G_l, V_n) = 0 \quad \forall n$$

$G_l$  has "property (F)", i.e.  $\exists$  only fin. many open subgroups of any given index

$$\Rightarrow H^i(G_l, V_n) \text{ fin. dim. } \forall i$$

$$\Rightarrow H^1(G_l, U') \text{ representable by Thm}$$

□

Get diagram for every prime  $l$

$$\begin{array}{ccc} X(\mathbb{Q}) & \hookrightarrow & X(\mathbb{Q}_l) \\ \downarrow j & & \downarrow j_l \\ H^1(G_{\mathbb{Q}}, U') & \xrightarrow{\text{loc}_l} & H^1(G_l, U') \end{array}$$

$j, j_l$  are "non-abelian Kummer maps" or "higher Albanese maps"



Selmer functor:

$$\text{Sel}_{U'}(X): R \mapsto \{ \alpha \in H^1(G_{\mathbb{Q}}, U')(R) \mid \forall \ell: \text{loc}_{\ell}(\alpha) \in j_{\ell}(X(\mathbb{Q}_{\ell}))^{\text{zar}} \}$$

Rmk:

- if  $\ell \neq p$ :  $j_{\ell}(X(\mathbb{Q}_{\ell}))$  finite,  
if  $\ell$  of good reduction:  $j_{\ell}(X(\mathbb{Q}_{\ell})) = \{*\}$
- if  $\ell = p$ :  $j_p(X(\mathbb{Q}_p))^{\text{zar}} = H_f^1(G_p, U')$ ,  $R \mapsto \ker(H^1(G_p, U')(R) \rightarrow H^1(G_p, U'(R \otimes_{\mathbb{Q}_p} B_{\text{cris}})))$   
subscheme of "crystalline classes" Fontaine's period ring

Thm: The Selmer functor is representable by an affine  $\mathbb{Q}_p$ -scheme.

This is the Selmer scheme  $\text{Sel}_{U'}(X)$ .

(Pf sketch: for  $T$  finite set of primes,

$G_T :=$  Galois gp of max. ext'n of  $\mathbb{Q}$  which is unram. outside  $T$ ,

$G_T$  (unlike  $G_{\mathbb{Q}}$ ) has property (F),

$\text{Sel}_{U'}(X) \subseteq H^1(G_T, U')$  for some  $T$ )

#### 4. The de Rham fundamental group

$X/\mathbb{Q}_p$  for now

$\text{MIC}^{\text{un}}(X)$ : Tannaka category of unipotent vector bundles w/ integrable connection.

objects:  $(E, \nabla)$  with  $E$  vector bundle on  $X$ ,

$\nabla: E \rightarrow \Omega_{\mathbb{Q}_p}^1 \otimes_{\mathbb{Q}_p} E$  connection ( $\nabla^2 = 0$  automatically)

$\exists$  filtration  $E = E_n \supseteq \dots \supseteq E_0 = 0$  s.t.  $E_{i+1}/E_i \cong (\mathcal{O}_X, d)$  unit object

$b \in X(\mathbb{Q}_p)$  defines fibre functor  $\text{Fib}_b: \text{MIC}^{\text{un}}(X) \rightarrow \mathbb{Q}_p\text{-Mod}_{f.d.}$

$U^{\text{dR}} := \underline{\text{Aut}}^{\otimes}(\text{Fib}_b)$  de Rham fundamental group

$P^{\text{dR}}(x, y) := \underline{\text{Isom}}^{\otimes}(\text{Fib}_x, \text{Fib}_y)$  de Rham path space ( $x, y \in X(\mathbb{Q}_p)$ )

Iterated integrals: Let  $\omega_1, \dots, \omega_n \in H^0(X, \Omega^1)$ ,  $\gamma \in \mathcal{P}^{dR}(x, y)(\mathbb{R})$ ,  $\mathbb{R}$   $\mathbb{Q}_p$ -algebra. 10

$\rightsquigarrow$  define  $\int_{\gamma} \omega_1 \dots \omega_n \in \mathbb{R}$  as follows

$$E_{\underline{\omega}} := (\mathcal{O}_X^{n+1}, \nabla) \quad \text{with} \quad \nabla \begin{pmatrix} f_1 \\ \vdots \\ f_{n+1} \end{pmatrix} = \begin{pmatrix} df_1 \\ \vdots \\ df_{n+1} \end{pmatrix} - \begin{pmatrix} 0 & \omega_1 & & \\ & 0 & \omega_2 & \\ & & 0 & \dots & \omega_n \\ & & & & 0 \end{pmatrix} \begin{pmatrix} f_1 \\ \vdots \\ f_{n+1} \end{pmatrix}$$

$$\gamma_{E_{\underline{\omega}}}: \mathbb{R}^{n+1} = \text{Fib}_x(E_{\underline{\omega}}) \longrightarrow \text{Fib}_y(E_{\underline{\omega}}) = \mathbb{R}^{n+1}$$

$$\int_{\gamma} \omega_1 \dots \omega_n := e_1^v(\gamma_{E_{\underline{\omega}}}(e_{n+1})) \in \mathbb{R}$$

Fact: Every alg. function on  $\mathcal{P}^{dR}(x, y)$  is a  $\mathbb{Q}_p$ -linear combination of iterated integrals  $\gamma \mapsto \int_{\gamma} \omega_1 \dots \omega_n$ .

Iterated Coleman integrals:

$X_0 :=$  special fibre of unique smooth model of  $X$  over  $\mathbb{Z}_p$

$\rightarrow$  Tamaya category  $\text{Isoc}^{un}(X_0)$  of unipotent isocrystals on  $X_0$

$\rightsquigarrow$  crystalline fundamental gp  $U^{cris}$ , path space  $\mathcal{P}^{cris}(x, y)$

Comparison thm:  $\text{MIC}^{un}(X) \simeq \text{Isoc}^{un}(X_0)$

$$\Rightarrow U^{dR} \cong U^{cris}, \quad \mathcal{P}^{dR}(x, y) \cong \mathcal{P}^{cris}(\bar{x}, \bar{y})$$

$$\rightsquigarrow \text{Frob} \subset U^{dR}, \quad \mathcal{P}^{dR}(x, y)$$

Thm: (Besser)

For all  $x, y \in X(\mathbb{Q}_p)$ ,  $\exists!$  Frobenius-invariant path  $p^{cr} \in \mathcal{P}^{dR}(x, y)(\mathbb{Q}_p)$ .

Def: Let  $x, y \in X(\mathbb{Q}_p)$ ,  $\omega_1, \dots, \omega_n \in H^0(X, \Omega^1)$ .

$$\int_x^y \omega_1 \dots \omega_n := \int_{p^{cr}} \omega_1 \dots \omega_n \in \mathbb{Q}_p \quad \text{iterated Coleman integral}$$

Rank: For  $x, y$  in the same residue disc,  $t$  unif. at  $x$ , can compute  $\int_x^y \omega_1 \dots \omega_n$  by expanding  $\omega_i = f_i(t)dt$  as convergent power series and

$$\text{integrating iteratively: } \int_x^y \omega_1 \dots \omega_n = \dots \int f_{n+1}(t) \int f_n(t) dt \Big|_{t=t(y)}$$

de Rham Kummer map

$U^{dR}$  carries Hodge filtration  $F \cdot U^{dR}$  by subschemes and Frobenius autom.  
 $F_0 U^{dR} \subseteq U^{dR}$  subgp.  $\rightarrow F^0 \backslash U^{dR}$  right coset scheme

Define "admissible  $U^{dR}$ -torsors": carry compatible Hodge filtr. + Frob.

Thm:  $(F^0 \backslash U^{dR})(R) \cong \left( \begin{smallmatrix} \text{admissible } U^{dR}\text{-torsors} \\ \text{over } R \end{smallmatrix} \right) / \text{iso}$

Sketch:  $\mathcal{P}$  adm.  $U^{dR}$ -torsor /  $R$

$\rightarrow \exists \rho^H \in F_0 \mathcal{P}(R), \exists \text{ unique } \rho^{cr} \in \mathcal{P}(R)^{Frob}$

$\mathcal{P} \mapsto (\rho^H)^{-1} \cdot \rho^{cr} \in (F^0 \backslash U^{dR})(R)$

□

de Rham Kummer map:

$$j^{dR}: X(\mathbb{Q}_p) \rightarrow F^0 \backslash U^{dR} \quad x \mapsto [\rho^{dR}(b, x)] = (\rho^H)^{-1} \cdot \rho^{cr}$$

$\uparrow$   
Hodge path

$\uparrow$   
Frob.-inv. path

pulling back  $(\int \omega_1 \dots \omega_n) \in \mathcal{O}(F^0 \backslash U^{dR})$  along gives iterated Coleman integral

$$X(\mathbb{Q}_p) \rightarrow \mathbb{Q}_p \quad x \mapsto \int_b^x \omega_1 \dots \omega_n.$$