

The Clowder Project

2017–2024



The Clowder Project Authors

Version 01f3714, compiled on Apr 04, 2024.

Contents

Part I	Sets	1
1	Sets	2
1.1	Sets and Functions	2
1.2	The Enrichment of Sets in Classical Truth Values	4
1.A	Other Chapters	10
2	Constructions With Sets	12
2.1	Limits of Sets	13
2.2	Colimits of Sets	36
2.3	Operations With Sets	54
2.4	Powersets	75
2.A	Other Chapters	119
3	Pointed Sets	120
3.1	Pointed Sets	120
3.2	Limits of Pointed Sets	126
3.3	Colimits of Pointed Sets	139
3.4	Constructions With Pointed Sets	153
3.A	Other Chapters	159
4	Tensor Products of Pointed Sets	160
4.1	Bilinear Morphisms of Pointed Sets	162
4.2	Tensors and Cotensors of Pointed Sets by Sets	166
4.3	The Left Tensor Product of Pointed Sets	182
4.4	The Right Tensor Product of Pointed Sets	208
4.5	The Smash Product of Pointed Sets	233
4.6	Miscellany	270
4.A	Other Chapters	270

Part II	Relations	271
5	Relations	272
5.1	Relations	274
5.2	Categories of Relations	284
5.3	Properties of the 2-Category of Relations	299
5.4	The Left Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	323
5.5	The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	328
5.A	Other Chapters	333
6	Constructions With Relations	335
6.1	Co/Limits in the Category of Relations	336
6.2	Kan Extensions and Kan Lifts in the 2-Category of Relations	337
6.3	More Constructions With Relations	342
6.4	Functoriality of Powersets	366
6.A	Other Chapters	394
7	Equivalence Relations and Apartness Relations	395
7.1	Reflexive Relations	395
7.2	Symmetric Relations	399
7.3	Transitive Relations	401
7.4	Equivalence Relations	406
7.5	Quotients by Equivalence Relations	408
7.A	Other Chapters	414
Part III	Category Theory	415
8	Categories	416
8.1	Categories	418
8.2	The Quadruple Adjunction With Sets	431
8.3	Groupoids	438
8.4	Functors	446
8.5	Conditions on Functors	456
8.6	More Conditions on Functors	473
8.7	Even More Conditions on Functors	483
8.8	Natural Transformations	492
8.9	Categories of Categories	506
8.A	Other Chapters	513
Part IV	Bicategories	514
9	Types of Morphisms in Bicategories	515
9.1	Monomorphisms in Bicategories	516
9.2	Epimorphisms in Bicategories	527

9.A Other Chapters	538
Part V Extra Part	539
10 Miscellaneous Notes	540
10.1 To Do List	540
10.A Other Chapters	547
Index of Notation	557
Index of Set Theory	560
Index of Category Theory	564
Index of Higher Category Theory	566

Contents

Part I	Sets	1
1	Sets	2
1.1	Sets and Functions	2
1.1.1	Functions	2
1.2	The Enrichment of Sets in Classical Truth Values	4
1.2.1	(-2) -Categories	4
1.2.2	(-1) -Categories	4
1.2.3	0-Categories	8
1.2.4	Tables of Analogies Between Set Theory and Category Theory	9
1.A	Other Chapters	10
2	Constructions With Sets	12
2.1	Limits of Sets	13
2.1.1	The Terminal Set	13
2.1.2	Products of Families of Sets	14
2.1.3	Binary Products of Sets	16
2.1.4	Pullbacks	25
2.1.5	Equalisers	32
2.2	Colimits of Sets	36
2.2.1	The Initial Set	36
2.2.2	Coproducts of Families of Sets	37
2.2.3	Binary Coproducts	39
2.2.4	Pushouts	42
2.2.5	Coequalisers	49
2.3	Operations With Sets	54
2.3.1	The Empty Set	54
2.3.2	Singleton Sets	54
2.3.3	Pairings of Sets	54
2.3.4	Ordered Pairs	55
2.3.5	Sets of Maps	55

2.3.6	Unions of Families	56
2.3.7	Binary Unions	56
2.3.8	Intersections of Families	59
2.3.9	Binary Intersections	59
2.3.10	Differences	64
2.3.11	Complements	67
2.3.12	Symmetric Differences	70
2.4	Powersets	75
2.4.1	Characteristic Functions	75
2.4.2	The Yoneda Lemma for Sets	81
2.4.3	Powersets	82
2.4.4	Direct Images	99
2.4.5	Inverse Images	105
2.4.6	Direct Images With Compact Support	109
2.A	Other Chapters	119
3	Pointed Sets	120
3.1	Pointed Sets	120
3.1.1	Foundations	120
3.1.2	Morphisms of Pointed Sets	122
3.1.3	The Category of Pointed Sets	122
3.1.4	Elementary Properties of Pointed Sets	123
3.2	Limits of Pointed Sets	126
3.2.1	The Terminal Pointed Set	126
3.2.2	Products of Families of Pointed Sets	127
3.2.3	Products	129
3.2.4	Pullbacks	131
3.2.5	Equalisers	136
3.3	Colimits of Pointed Sets	139
3.3.1	The Initial Pointed Set	139
3.3.2	Coproducts of Families of Pointed Sets	140
3.3.3	Coproducts	142
3.3.4	Pushouts	146
3.3.5	Coequalisers	151
3.4	Constructions With Pointed Sets	153
3.4.1	Free Pointed Sets	153
3.A	Other Chapters	159
4	Tensor Products of Pointed Sets	160
4.1	Bilinear Morphisms of Pointed Sets	162
4.1.1	Left Bilinear Morphisms of Pointed Sets	162
4.1.2	Right Bilinear Morphisms of Pointed Sets	163
4.1.3	Bilinear Morphisms of Pointed Sets	164

4.2	Tensors and Cotensors of Pointed Sets by Sets	166
4.2.1	Tensors of Pointed Sets by Sets	166
4.2.2	Cotensors of Pointed Sets by Sets	174
4.3	The Left Tensor Product of Pointed Sets	182
4.3.1	Foundations	182
4.3.2	The Left Internal Hom of Pointed Sets	187
4.3.3	The Left Skew Unit	189
4.3.4	The Left Skew Associator	189
4.3.5	The Left Skew Left Unitor	192
4.3.6	The Left Skew Right Unitor	195
4.3.7	The Diagonal	197
4.3.8	The Left Skew Monoidal Structure on Pointed Sets Asso-	
	ciated to \triangleleft	199
4.3.9	Monoids With Respect to the Left Tensor Product of Pointed	
	Sets	203
4.4	The Right Tensor Product of Pointed Sets	208
4.4.1	Foundations	208
4.4.2	The Right Internal Hom of Pointed Sets	212
4.4.3	The Right Skew Unit	215
4.4.4	The Right Skew Associator	215
4.4.5	The Right Skew Left Unitor	218
4.4.6	The Right Skew Right Unitor	220
4.4.7	The Diagonal	223
4.4.8	The Right Skew Monoidal Structure on Pointed Sets As-	
	sociated to \triangleright	224
4.4.9	Monoids With Respect to the Right Tensor Product of	
	Pointed Sets	229
4.5	The Smash Product of Pointed Sets	233
4.5.1	Foundations	233
4.5.2	The Internal Hom of Pointed Sets	243
4.5.3	The Monoidal Unit	247
4.5.4	The Associator	247
4.5.5	The Left Unitor	250
4.5.6	The Right Unitor	253
4.5.7	The Symmetry	256
4.5.8	The Diagonal	258
4.5.9	The Monoidal Structure on Pointed Sets Associated to \wedge .	262
4.5.10	Universal Properties of the Smash Product of Pointed	
	Sets I	267
4.5.11	Universal Properties of the Smash Product of Pointed	
	Sets II	268
4.5.12	Monoids With Respect to the Smash Product of Pointed	
	Sets	269

4.5.13 Comonoids With Respect to the Smash Product of Pointed
 Sets 269
 4.6 Miscellany 270
 4.6.1 The Smash Product of a Family of Pointed Sets 270
 4.A Other Chapters 270

Part II Relations 271

5 Relations 272
 5.1 Relations 274
 5.1.1 Foundations 274
 5.1.2 Relations as Decategorifications of Profunctors 278
 5.1.3 Examples of Relations 279
 5.1.4 Functional Relations 282
 5.1.5 Total Relations 283
 5.2 Categories of Relations 284
 5.2.1 The Category of Relations 284
 5.2.2 The Closed Symmetric Monoidal Category of Relations ... 284
 5.2.3 The 2-Category of Relations 290
 5.2.4 The Double Category of Relations 291
 5.3 Properties of the 2-Category of Relations 299
 5.3.1 Self-Duality 299
 5.3.2 Isomorphisms and Equivalences in **Rel** 300
 5.3.3 Adjunctions in **Rel** 301
 5.3.4 Monads in **Rel** 304
 5.3.5 Comonads in **Rel** 305
 5.3.6 Co/Monoids in **Rel** 306
 5.3.7 Monomorphisms in **Rel** 306
 5.3.8 2-Categorical Monomorphisms in **Rel** 309
 5.3.9 Epimorphisms in **Rel** 313
 5.3.10 2-Categorical Epimorphisms in **Rel** 317
 5.3.11 Co/Limits in **Rel** 321
 5.3.12 Kan Extensions and Kan Lifts in **Rel** 321
 5.3.13 Closedness of **Rel** 321
 5.3.14 **Rel** as a Category of Free Algebras 322
 5.4 The Left Skew Monoidal Structure on **Rel**(A, B) 323
 5.4.1 The Left Skew Monoidal Product 323
 5.4.2 The Left Skew Monoidal Unit 324
 5.4.3 The Left Skew Associators 324
 5.4.4 The Left Skew Left Unitors 325
 5.4.5 The Left Skew Right Unitors 326
 5.4.6 The Left Skew Monoidal Structure on **Rel**(A, B) 327

5.5	The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	328
5.5.1	The Right Skew Monoidal Product.....	328
5.5.2	The Right Skew Monoidal Unit.....	329
5.5.3	The Right Skew Associators.....	330
5.5.4	The Right Skew Left Unitors.....	331
5.5.5	The Right Skew Right Unitors.....	331
5.5.6	The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	332
5.A	Other Chapters.....	333
6	Constructions With Relations.....	335
6.1	Co/Limits in the Category of Relations.....	336
6.2	Kan Extensions and Kan Lifts in the 2-Category of Relations.....	337
6.2.1	Left Kan Extensions in \mathbf{Rel}	337
6.2.2	Left Kan Lifts in \mathbf{Rel}	338
6.2.3	Right Kan Extensions in \mathbf{Rel}	339
6.2.4	Right Kan Lifts in \mathbf{Rel}	340
6.3	More Constructions With Relations.....	342
6.3.1	The Graph of a Function.....	342
6.3.2	The Inverse of a Function.....	346
6.3.3	Representable Relations.....	348
6.3.4	The Domain and Range of a Relation.....	348
6.3.5	Binary Unions of Relations.....	349
6.3.6	Unions of Families of Relations.....	351
6.3.7	Binary Intersections of Relations.....	352
6.3.8	Intersections of Families of Relations.....	353
6.3.9	Binary Products of Relations.....	354
6.3.10	Products of Families of Relations.....	356
6.3.11	The Inverse of a Relation.....	357
6.3.12	Composition of Relations.....	359
6.3.13	The Collage of a Relation.....	364
6.4	Functoriality of Powersets.....	366
6.4.1	Direct Images.....	366
6.4.2	Strong Inverse Images.....	372
6.4.3	Weak Inverse Images.....	378
6.4.4	Direct Images With Compact Support.....	384
6.4.5	Functoriality of Powersets.....	391
6.4.6	Functoriality of Powersets: Relations on Powersets.....	392
6.A	Other Chapters.....	394
7	Equivalence Relations and Apartness Relations.....	395
7.1	Reflexive Relations.....	395
7.1.1	Foundations.....	395
7.1.2	The Reflexive Closure of a Relation.....	397

7.2	Symmetric Relations	399
7.2.1	Foundations	399
7.2.2	The Symmetric Closure of a Relation	399
7.3	Transitive Relations	401
7.3.1	Foundations	401
7.3.2	The Transitive Closure of a Relation	403
7.4	Equivalence Relations	406
7.4.1	Foundations	406
7.4.2	The Equivalence Closure of a Relation	407
7.5	Quotients by Equivalence Relations	408
7.5.1	Equivalence Classes	408
7.5.2	Quotients of Sets by Equivalence Relations	409
7.A	Other Chapters	414
Part III Category Theory		415
8	Categories	416
8.1	Categories	418
8.1.1	Foundations	418
8.1.2	Examples of Categories	420
8.1.3	Posetal Categories	424
8.1.4	Subcategories	425
8.1.5	Skeletons of Categories	426
8.1.6	Precomposition and Postcomposition	428
8.2	The Quadruple Adjunction With Sets	431
8.2.1	Statement	431
8.2.2	Connected Components and Connected Categories	432
8.2.3	Discrete Categories	435
8.2.4	Indiscrete Categories	437
8.3	Groupoids	438
8.3.1	Foundations	439
8.3.2	The Groupoid Completion of a Category	439
8.3.3	The Core of a Category	443
8.4	Functors	446
8.4.1	Foundations	446
8.4.2	Contravariant Functors	450
8.4.3	Forgetful Functors	452
8.4.4	The Natural Transformation Associated to a Functor	454
8.5	Conditions on Functors	456
8.5.1	Faithful Functors	456
8.5.2	Full Functors	459
8.5.3	Fully Faithful Functors	462
8.5.4	Conservative Functors	466

8.5.5	Essentially Injective Functors	468
8.5.6	Essentially Surjective Functors	469
8.5.7	Equivalences of Categories	469
8.5.8	Isomorphisms of Categories	472
8.6	More Conditions on Functors	473
8.6.1	Dominant Functors	473
8.6.2	Monomorphisms of Categories	475
8.6.3	Epimorphisms of Categories	476
8.6.4	Pseudomononic Functors	478
8.6.5	Pseudoepic Functors	480
8.7	Even More Conditions on Functors	483
8.7.1	Injective on Objects Functors	483
8.7.2	Surjective on Objects Functors	484
8.7.3	Bijjective on Objects Functors	484
8.7.4	Functors Representably Faithful on Cores	484
8.7.5	Functors Representably Full on Cores	485
8.7.6	Functors Representably Fully Faithful on Cores	486
8.7.7	Functors Corepresentably Faithful on Cores	487
8.7.8	Functors Corepresentably Full on Cores	489
8.7.9	Functors Corepresentably Fully Faithful on Cores	490
8.8	Natural Transformations	492
8.8.1	Transformations	492
8.8.2	Natural Transformations	492
8.8.3	Vertical Composition of Natural Transformations	493
8.8.4	Horizontal Composition of Natural Transformations	497
8.8.5	Properties of Natural Transformations	503
8.8.6	Natural Isomorphisms	504
8.9	Categories of Categories	506
8.9.1	Functor Categories	506
8.9.2	The Category of Categories and Functors	510
8.9.3	The 2-Category of Categories, Functors, and Natural Trans-	
	formations	511
8.9.4	The Category of Groupoids	512
8.9.5	The 2-Category of Groupoids	512
8.A	Other Chapters	513

Part IV Bicategories 514

9	Types of Morphisms in Bicategories	515
9.1	Monomorphisms in Bicategories	516
9.1.1	Representably Faithful Morphisms	516
9.1.2	Representably Full Morphisms	517
9.1.3	Representably Fully Faithful Morphisms	518

9.1.4	Morphisms Representably Faithful on Cores	519
9.1.5	Morphisms Representably Full on Cores	520
9.1.6	Morphisms Representably Fully Faithful on Cores	521
9.1.7	Representably Essentially Injective Morphisms	522
9.1.8	Representably Conservative Morphisms	523
9.1.9	Strict Monomorphisms	524
9.1.10	Pseudomononic Morphisms	525
9.2	Epimorphisms in Bicategories	527
9.2.1	Corepresentably Faithful Morphisms	527
9.2.2	Corepresentably Full Morphisms	528
9.2.3	Corepresentably Fully Faithful Morphisms	529
9.2.4	Morphisms Corepresentably Faithful on Cores	530
9.2.5	Morphisms Corepresentably Full on Cores	531
9.2.6	Morphisms Corepresentably Fully Faithful on Cores	532
9.2.7	Corepresentably Essentially Injective Morphisms	533
9.2.8	Corepresentably Conservative Morphisms	534
9.2.9	Strict Epimorphisms	535
9.2.10	Pseudoeptic Morphisms	536
9.A	Other Chapters	538
Part V Extra Part		539
10	Miscellaneous Notes	540
10.1	To Do List	540
10.1.1	Omitted Proofs To Add	540
10.1.2	Things To Explore/Add	541
10.A	Other Chapters	547
Index of Notation		557
Index of Set Theory		560
Index of Category Theory		564
Index of Higher Category Theory		566

Part I

Sets

Chapter 1

Sets

0000 This chapter (will eventually) contain material on axiomatic set theory, as well as a couple other things.

Contents

1.1	Sets and Functions	2
1.1.1	Functions.....	2
1.2	The Enrichment of Sets in Classical Truth Values	4
1.2.1	(−2)-Categories.....	4
1.2.2	(−1)-Categories.....	4
1.2.3	0-Categories.....	8
1.2.4	Tables of Analogies Between Set Theory and Category Theory.....	9
1.A	Other Chapters	10

0001 1.1 Sets and Functions

0002 1.1.1 Functions

0003 DEFINITION 1.1.1.1 ► FUNCTIONS

A **function** is a functional and total relation.

0004 NOTATION 1.1.1.2 ► ADDITIONAL NOTATION FOR FUNCTIONS

Throughout this work, we will sometimes denote a function $f: X \rightarrow Y$ by

$$f \stackrel{\text{def}}{=} \llbracket x \mapsto f(x) \rrbracket.$$

1. For example, given a function

$$\Phi: \text{Hom}_{\text{Sets}}(X, Y) \rightarrow K$$

taking values on a set of functions such as $\text{Hom}_{\text{Sets}}(X, Y)$, we will sometimes also write

$$\Phi(f) \stackrel{\text{def}}{=} \Phi(\llbracket x \mapsto f(x) \rrbracket).$$

2. This notational choice is based on the lambda notation

$$f \stackrel{\text{def}}{=} (\lambda x. f(x)),$$

but uses a “ \mapsto ” symbol for better spacing and double brackets instead of either:

- (a) Square brackets $[x \mapsto f(x)]$;
- (b) Parentheses $(x \mapsto f(x))$;

hoping to improve readability when dealing with e.g.:

- (a) Equivalence classes, cf.:

- i. $\llbracket [x] \mapsto f([x]) \rrbracket$
- ii. $\llbracket [x] \mapsto f([x]) \rrbracket$
- iii. $(\lambda [x]. f([x]))$

- (b) Function evaluations, cf.:

- i. $\Phi(\llbracket x \mapsto f(x) \rrbracket)$
- ii. $\Phi((x \mapsto f(x)))$
- iii. $\Phi((\lambda x. f(x)))$

3. We will also sometimes write $-_1, -_2$, etc. for the arguments of a function. Some examples include:

- (a) Writing $f(-_1)$ for a function $f: A \rightarrow B$.
- (b) Writing $f(-_1, -_2)$ for a function $f: A \times B \rightarrow C$.
- (c) Given a function $f: A \times B \rightarrow C$, writing

$$f(a, -): B \rightarrow C$$

for the function $\llbracket b \mapsto f(a, b) \rrbracket$.

(d) Denoting a composition of the form

$$A \times B \xrightarrow{\phi \times \text{id}_B} A' \times B \xrightarrow{f} C$$

by $f(\phi(-_1), -_2)$.

4. Finally, given a function $f: A \rightarrow B$, we write

$$\text{ev}_a(f) \stackrel{\text{def}}{=} f(a)$$

for the value of f at some $a \in A$.

For an example of the above notations being used in practice, see the proof of the adjunction

$$(A \times - \dashv \text{Hom}_{\text{Sets}}(A, -)) : \text{Sets} \begin{array}{c} \xrightarrow{A \times -} \\ \perp \\ \xleftarrow{\text{Hom}_{\text{Sets}}(A, -)} \end{array} \text{Sets},$$

stated in **Item 2** of **Proposition 2.1.3.3**.

0005 1.2 The Enrichment of Sets in Classical Truth Values

0006 1.2.1 (-2) -Categories

0007 DEFINITION 1.2.1.1 \blacktriangleright (-2) -CATEGORIES

A (-2) -**category** is the “necessarily true” truth value.^{1,2,3}

¹Thus, there is only one (-2) -category.

²A $(-n)$ -category for $n = 3, 4, \dots$ is also the “necessarily true” truth value, coinciding with a (-2) -category.

³For motivation, see [BS10, p. 13].

0008 1.2.2 (-1) -Categories

0009 DEFINITION 1.2.2.1 \blacktriangleright (-1) -CATEGORIES

A (-1) -**category** is a classical truth value.

000A

REMARK 1.2.2.2 ► MOTIVATION FOR (-1) -CATEGORIES

¹ (-1) -categories should be thought of as being “categories enriched in (-2) -categories”, having a collection of objects and, for each pair of objects, a Hom-object $\text{Hom}(x, y)$ that is a (-2) -category (i.e. trivial).

Therefore, a (-1) -category C is either ([BS10, pp. 33–34]):

1. *Empty*, having no objects;
2. *Contractible*, having a collection of objects $\{a, b, c, \dots\}$, but with $\text{Hom}_C(a, b)$ being a (-2) -category (i.e. trivial) for all $a, b \in \text{Obj}(C)$, forcing all objects of C to be uniquely isomorphic to each other.

As such, there are only two (-1) -categories, up to equivalence:

- The (-1) -category false (the empty one);
- The (-1) -category true (the contractible one).

¹For more motivation, see [BS10, p. 13].

000B

DEFINITION 1.2.2.3 ► THE POSET OF TRUTH VALUES

The **poset of truth values**¹ is the poset $(\{\text{true}, \text{false}\}, \preceq)$ consisting of

- *The Underlying Set.* The set $\{\text{true}, \text{false}\}$ whose elements are the truth values true and false.
- *The Partial Order.* The partial order

$$\preceq: \{\text{true}, \text{false}\} \times \{\text{true}, \text{false}\} \rightarrow \{\text{true}, \text{false}\}$$

on $\{\text{true}, \text{false}\}$ defined by²

$$\begin{aligned} \text{false} \preceq \text{false} &\stackrel{\text{def}}{=} \text{true}, \\ \text{true} \preceq \text{false} &\stackrel{\text{def}}{=} \text{false}, \\ \text{false} \preceq \text{true} &\stackrel{\text{def}}{=} \text{true}, \\ \text{true} \preceq \text{true} &\stackrel{\text{def}}{=} \text{true}. \end{aligned}$$

¹*Further Terminology:* Also called the **poset of (-1) -categories**.

²This partial order coincides with logical implication.

000C

NOTATION 1.2.2.4 ► FURTHER NOTATION FOR THE POSET OF TRUTH VALUES

We also write $\{t, f\}$ for the poset $\{\text{true}, \text{false}\}$.

000D

PROPOSITION 1.2.2.5 ▶ CARTESIAN CLOSEDNESS OF THE POSET OF TRUTH VALUES

The poset of truth values $\{t, f\}$ is Cartesian closed with product given by¹

$$\begin{aligned} t \times t &= t, \\ t \times f &= f, \\ f \times t &= f, \\ f \times f &= f, \end{aligned}$$

and internal Hom $\mathbf{Hom}_{\{t,f\}}$ given by the partial order of $\{t, f\}$, i.e. by


$$\begin{aligned} \mathbf{Hom}_{\{t,f\}}(t, t) &= t, \\ \mathbf{Hom}_{\{t,f\}}(t, f) &= f, \\ \mathbf{Hom}_{\{t,f\}}(f, t) &= t, \\ \mathbf{Hom}_{\{t,f\}}(f, f) &= t. \end{aligned}$$

¹Note that \times coincides with the “and” operator, while $\mathbf{Hom}_{\{t,f\}}$ coincides with the logical implication operator.

PROOF 1.2.2.6 ▶ PROOF OF PROPOSITION 1.2.2.5

Existence of Products

We claim that the products $t \times t$, $t \times f$, $f \times t$, and $f \times f$ satisfy the universal property of the product in $\{t, f\}$. Indeed, consider the diagrams



Here:

1. If $P_1 = t$, then $p_1^1 = p_2^1 = \text{id}_t$, and there's indeed a unique morphism from P_1 to t making the diagram commute, namely id_t ;
2. If $P_1 = f$, then $p_1^1 = p_2^1$ are given by the unique morphism from f to t , and there's indeed a unique morphism from P_1 to t making the diagram commute, namely the unique morphism from f to t ;
3. If $P_2 = t$, then there is no morphism p_2^2 .
4. If $P_2 = f$, then p_1^2 is the unique morphism from f to t while $p_2^2 = \text{id}_f$, and

there's indeed a unique morphism from P_2 to f making the diagram commute, namely id_f ;

5. The proof for P_3 is similar to the one for P_2 ;
6. If $P_4 = t$, then there is no morphism p_1^4 or p_2^4 .
7. If $P_4 = f$, then $p_1^4 = p_2^4 = \text{id}_f$, and there's indeed a unique morphism from P_4 to f making the diagram commute, namely id_f .

Cartesian Closedness

We claim there's a bijection

$$\text{Hom}_{\{t,f\}}(A \times B, C) \cong \text{Hom}_{\{t,f\}}(A, \mathbf{Hom}_{\{t,f\}}(B, C))$$

natural in $A, B, C \in \{t, f\}$. Indeed:

- For $(A, B, C) = (t, t, t)$, we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(t \times t, t) &\cong \text{Hom}_{\{t,f\}}(t, t) \\ &= \{\text{id}_{\text{true}}\} \\ &\cong \text{Hom}_{\{t,f\}}(t, t) \\ &\cong \text{Hom}_{\{t,f\}}(t, \mathbf{Hom}_{\{t,f\}}(t, t)). \end{aligned}$$

- For $(A, B, C) = (t, t, f)$, we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(t \times t, f) &\cong \text{Hom}_{\{t,f\}}(t, f) \\ &= \emptyset \\ &\cong \text{Hom}_{\{t,f\}}(t, f) \\ &\cong \text{Hom}_{\{t,f\}}(t, \mathbf{Hom}_{\{t,f\}}(t, f)). \end{aligned}$$

- For $(A, B, C) = (t, f, t)$, we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(t \times f, t) &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong \text{pt} \\ &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong \text{Hom}_{\{t,f\}}(f, \mathbf{Hom}_{\{t,f\}}(f, t)). \end{aligned}$$

- For $(A, B, C) = (t, f, f)$, we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(t \times f, f) &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &\cong \{\text{id}_{\text{false}}\} \\ &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &\cong \text{Hom}_{\{t,f\}}(t, \mathbf{Hom}_{\{t,f\}}(f, f)). \end{aligned}$$

- For $(A, B, C) = (f, t, t)$, we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(f \times t, t) &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong \text{pt} \\ &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong \text{Hom}_{\{t,f\}}(f, \mathbf{Hom}_{\{t,f\}}(t, t)). \end{aligned}$$

- For $(A, B, C) = (f, t, f)$, we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(f \times t, f) &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &\cong \{\text{id}_{\text{false}}\} \\ &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &\cong \text{Hom}_{\{t,f\}}(f, \mathbf{Hom}_{\{t,f\}}(t, f)). \end{aligned}$$

- For $(A, B, C) = (f, f, t)$, we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(f \times f, t) &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong \text{pt} \\ &\cong \text{Hom}_{\{t,f\}}(f, t) \\ &\cong \text{Hom}_{\{t,f\}}(f, \mathbf{Hom}_{\{t,f\}}(f, t)). \end{aligned}$$

- For $(A, B, C) = (f, f, f)$, we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(f \times f, f) &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &= \{\text{id}_{\text{false}}\} \\ &\cong \text{Hom}_{\{t,f\}}(f, f) \\ &\cong \text{Hom}_{\{t,f\}}(f, \mathbf{Hom}_{\{t,f\}}(f, f)). \end{aligned}$$

The proof of naturality is omitted.



000E 1.2.3 0-Categories

000F DEFINITION 1.2.3.1 ► 0-CATEGORIES

A **0-category** is a poset.¹

¹Motivation: A 0-category is precisely a category enriched in the poset of (-1) -categories.

000G

DEFINITION 1.2.3.2 ► 0-GROUPOIDS

A **0-groupoid** is a 0-category in which every morphism is invertible.¹

¹That is, a *set*.

000H 1.2.4 Tables of Analogies Between Set Theory and Category Theory

Here we record some analogies between notions in set theory and category theory. Note that the analogies relating to presheaves relate equally well to copresheaves, as the opposite X^{op} of a set X is just X again.

Basics:

SET THEORY	CATEGORY THEORY
Enrichment in $\{\text{true}, \text{false}\}$	Enrichment in Sets
Set X	Category \mathcal{C}
Element $x \in X$	Object $X \in \text{Obj}(\mathcal{C})$
Function	Functor
Function $X \rightarrow \{\text{true}, \text{false}\}$	Functor $\mathcal{C} \rightarrow \text{Sets}$
Function $X \rightarrow \{\text{true}, \text{false}\}$	Presheaf $\mathcal{C}^{\text{op}} \rightarrow \text{Sets}$

Powersets and categories of presheaves:

SET THEORY	CATEGORY THEORY
Powerset $\mathcal{P}(X)$	Presheaf category $\text{PSh}(\mathcal{C})$
Characteristic function $\chi_{\{x\}}$	Representable presheaf h_X
Characteristic embedding $\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$	Yoneda embedding $\mathcal{Y} : \mathcal{C}^{\text{op}} \hookrightarrow \text{PSh}(\mathcal{C})$
Characteristic relation $\chi_X(-_1, -_2)$	Hom profunctor $\text{Hom}_{\mathcal{C}}(-_1, -_2)$
The Yoneda lemma for sets $\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_U) = \chi_U(x)$	The Yoneda lemma for categories $\text{Nat}(h_X, \mathcal{F}) \cong \mathcal{F}(X)$
The characteristic embedding is fully faithful, $\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_y) = \chi_X(x, y)$	The Yoneda embedding is fully faithful, $\text{Nat}(h_X, h_Y) \cong \text{Hom}_{\mathcal{C}}(X, Y)$
Subsets are unions of their elements $U = \bigcup_{x \in U} \{x\}$ or $\chi_U = \text{colim}_{\chi_x \in \text{Sets}(U, \{\text{t}, \text{f}\})} (\chi_x)$	Presheaves are colimits of representables, $\mathcal{F} \cong \text{colim}_{h_X \in \int_{\mathcal{C}} \mathcal{F}} (h_X)$

Categories of elements:

SET THEORY	CATEGORY THEORY
Assignment $U \mapsto \chi_U$	Assignment $\mathcal{F} \mapsto \int_C \mathcal{F}$ (the category of elements)
Assignment $U \mapsto \chi_U$ giving an isomorphism $\mathcal{P}(X) \cong \text{Sets}(X, \{t, f\})$	Assignment $\mathcal{F} \mapsto \int_C \mathcal{F}$ giving an equivalence $\text{PSh}(C) \cong^{\text{eq.}} \text{DFib}(C)$

Functions between powersets and functors between presheaf categories:

SET THEORY	CATEGORY THEORY
Direct image function $f_*: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$	Inverse image functor $f^{-1}: \text{PSh}(C) \rightarrow \text{PSh}(D)$
Inverse image function $f^{-1}: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$	Direct image functor $f_*: \text{PSh}(D) \rightarrow \text{PSh}(C)$
Direct image with compact support function $f_!: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$	Direct image with compact support functor $f_!: \text{PSh}(C) \rightarrow \text{PSh}(D)$

Relations and profunctors:

SET THEORY	CATEGORY THEORY
Relation $R: X \times Y \rightarrow \{t, f\}$	Profunctor $\mathfrak{p}: \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}$
Relation $R: X \rightarrow \mathcal{P}(Y)$	Profunctor $\mathfrak{p}: C \rightarrow \text{PSh}(D)$
Relation as a cocontinuous morphism of posets $R: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(Y), \subset)$	Profunctor as a colimit-preserving functor $\mathfrak{p}: \text{PSh}(C) \rightarrow \text{PSh}(D)$

Appendices

1.A Other Chapters

Sets

1. [Sets](#)
2. [Constructions With Sets](#)
3. [Pointed Sets](#)
4. [Tensor Products of Pointed Sets](#)

Relations

5. [Relations](#)
6. [Constructions With Relations](#)
7. [Equivalence Relations and Apartness Relations](#)

Category Theory

8. **Categories**
Bicategories

9. **Types of Morphisms in Bicat-**
egories

Chapter 2

Constructions With Sets

000J This chapter develops some material relating to constructions with sets with an eye towards its categorical and higher-categorical counterparts to be introduced later in this work. In particular, it contains:

1. Explicit descriptions of the major types of co/limits in Sets, including in particular explicit descriptions of pushouts and coequalisers (see [Definitions 2.2.4.1](#) and [2.2.5.1](#) and [Remarks 2.2.4.3](#) and [2.2.5.3](#)).
2. A discussion of powersets as decategorifications of categories of presheaves ([Remarks 2.4.1.2](#) and [2.4.3.2](#)), including a (-1) -categorical analogue of un/s-traightening, described in [Items 1](#) and [2](#) of [Proposition 2.4.3.9](#) and [Remark 2.4.3.11](#).
3. A lengthy discussion of the adjoint triple

$$f_* \dashv f^{-1} \dashv f_! : \mathcal{P}(A) \xrightarrow{\cong} \mathcal{P}(B)$$

of functors (morphisms of posets) between $\mathcal{P}(A)$ and $\mathcal{P}(B)$ induced by a map of sets $f: A \rightarrow B$, along with a discussion of the properties of f_* , f^{-1} , and $f_!$.

In line with the categorical viewpoint developed here, this adjoint triple may be described in terms of Kan extensions, and, as it turns out, it also shows up in some definitions and results in point-set topology, such as in e.g. notions of continuity for functions (??).

Contents

2.1	Limits of Sets	13
2.1.1	The Terminal Set	13
2.1.2	Products of Families of Sets	14
2.1.3	Binary Products of Sets	16
2.1.4	Pullbacks	25

2.1.5	Equalisers	32
2.2	Colimits of Sets	36
2.2.1	The Initial Set	36
2.2.2	Coproducts of Families of Sets	37
2.2.3	Binary Coproducts	39
2.2.4	Pushouts	42
2.2.5	Coequalisers	49
2.3	Operations With Sets	54
2.3.1	The Empty Set	54
2.3.2	Singleton Sets	54
2.3.3	Pairings of Sets	54
2.3.4	Ordered Pairs	55
2.3.5	Sets of Maps	55
2.3.6	Unions of Families	56
2.3.7	Binary Unions	56
2.3.8	Intersections of Families	59
2.3.9	Binary Intersections	59
2.3.10	Differences	64
2.3.11	Complements	67
2.3.12	Symmetric Differences	70
2.4	Powersets	75
2.4.1	Characteristic Functions	75
2.4.2	The Yoneda Lemma for Sets	81
2.4.3	Powersets	82
2.4.4	Direct Images	99
2.4.5	Inverse Images	105
2.4.6	Direct Images With Compact Support	109
2.A	Other Chapters	119

000K 2.1 Limits of Sets

000L 2.1.1 The Terminal Set

000M DEFINITION 2.1.1.1 ► THE TERMINAL SET

The **terminal set** is the pair $(pt, \{!_A\}_{A \in \text{Obj}(\text{Sets})})$ consisting of:

- *The Limit.* The punctual set $pt \stackrel{\text{def}}{=} \{\star\}$.

- *The Cone.* The collection of maps

$$\{!_A : A \rightarrow \text{pt}\}_{A \in \text{Obj}(\text{Sets})}$$

defined by

$$!_A(a) \stackrel{\text{def}}{=} \star$$

for each $a \in A$ and each $A \in \text{Obj}(\text{Sets})$.


PROOF 2.1.1.2 ► PROOF OF DEFINITION 2.1.1.1

We claim that pt is the terminal object of Sets . Indeed, suppose we have a diagram of the form

$$A \quad \text{pt}$$

in Sets . Then there exists a unique map $\phi : A \rightarrow \text{pt}$ making the diagram

$$A \xrightarrow[\exists!]{\phi} \text{pt}$$

commute, namely $!_A$. 

000N 2.1.2 Products of Families of Sets

Let $\{A_i\}_{i \in I}$ be a family of sets.

000P DEFINITION 2.1.2.1 ► THE PRODUCT OF A FAMILY OF SETS

The **product**¹ of $\{A_i\}_{i \in I}$ is the pair $(\prod_{i \in I} A_i, \{\text{pr}_i\}_{i \in I})$ consisting of:

- *The Limit.* The set $\prod_{i \in I} A_i$ defined by²

$$\prod_{i \in I} A_i \stackrel{\text{def}}{=} \left\{ f \in \text{Sets} \left(I, \bigcup_{i \in I} A_i \right) \mid \text{for each } i \in I, \text{ we have } f(i) \in A_i \right\}.$$

- *The Cone.* The collection

$$\left\{ \text{pr}_i : \prod_{i \in I} A_i \rightarrow A_i \right\}_{i \in I}$$

of maps given by

$$\text{pr}_i(f) \stackrel{\text{def}}{=} f(i)$$

for each $f \in \prod_{i \in I} A_i$ and each $i \in I$.

¹*Further Terminology:* Also called the **Cartesian product** of $\{A_i\}_{i \in I}$.

²Less formally, $\prod_{i \in I} A_i$ is the set whose elements are I -indexed collections $(a_i)_{i \in I}$ with $a_i \in A_i$ for each $i \in I$. The projection maps

$$\left\{ \text{pr}_i : \prod_{i \in I} A_i \rightarrow A_i \right\}_{i \in I}$$

are then given by

$$\text{pr}_i \left((a_j)_{j \in I} \right) \stackrel{\text{def}}{=} a_i$$

for each $(a_j)_{j \in I} \in \prod_{i \in I} A_i$ and each $i \in I$.

PROOF 2.1.2.2 ► PROOF OF DEFINITION 2.1.2.1

We claim that $\prod_{i \in I} A_i$ is the categorical product of $\{A_i\}_{i \in I}$ in Sets. Indeed, suppose we have, for each $i \in I$, a diagram of the form

$$\begin{array}{ccc} P & & \\ & \searrow^{p_i} & \\ \prod_{i \in I} A_i & \xrightarrow{\text{pr}_i} & A_i \end{array}$$

in Sets. Then there exists a unique map $\phi : P \rightarrow \prod_{i \in I} A_i$ making the diagram

$$\begin{array}{ccc} P & & \\ \downarrow \phi \exists! & \searrow^{p_i} & \\ \prod_{i \in I} A_i & \xrightarrow{\text{pr}_i} & A_i \end{array}$$

commute, being uniquely determined by the condition $\text{pr}_i \circ \phi = p_i$ for each $i \in I$ via

$$\phi(x) = (p_i(x))_{i \in I}$$

for each $x \in P$. □

000Q

PROPOSITION 2.1.2.3 ► PROPERTIES OF PRODUCTS OF FAMILIES OF SETS

Let $\{A_i\}_{i \in I}$ be a family of sets.

000R

1. *Functoriality.* The assignment $\{A_i\}_{i \in I} \mapsto \prod_{i \in I} A_i$ defines a functor

$$\prod_{i \in I}: \text{Fun}(I_{\text{disc}}, \text{Sets}) \rightarrow \text{Sets}$$

where

- *Action on Objects.* For each $(A_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$, we have

$$\left[\prod_{i \in I} \right] ((A_i)_{i \in I}) \stackrel{\text{def}}{=} \prod_{i \in I} A_i$$

- *Action on Morphisms.* For each $(A_i)_{i \in I}, (B_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$, the action on Hom-sets

$$\left(\prod_{i \in I} \right)_{(A_i)_{i \in I}, (B_i)_{i \in I}} : \text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I}) \rightarrow \text{Sets} \left(\prod_{i \in I} A_i, \prod_{i \in I} B_i \right)$$

of $\prod_{i \in I}$ at $((A_i)_{i \in I}, (B_i)_{i \in I})$ is defined by sending a map

$$\{f_i: A_i \rightarrow B_i\}_{i \in I}$$

in $\text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I})$ to the map of sets

$$\prod_{i \in I} f_i: \prod_{i \in I} A_i \rightarrow \prod_{i \in I} B_i$$

defined by

$$\left[\prod_{i \in I} f_i \right] ((a_i)_{i \in I}) \stackrel{\text{def}}{=} (f_i(a_i))_{i \in I}$$

for each $(a_i)_{i \in I} \in \prod_{i \in I} A_i$.

PROOF 2.1.2.4 ► PROOF OF PROPOSITION 2.1.2.3

Item 1: Functoriality

This follows from ?? of ??.



000S 2.1.3 Binary Products of Sets

Let A and B be sets.

000T

DEFINITION 2.1.3.1 ► PRODUCTS OF SETS

The **product**¹ of **A and B** is the pair $(A \times B, \{pr_1, pr_2\})$ consisting of:

- *The Limit.* The set $A \times B$ defined by²

$$\begin{aligned} A \times B &\stackrel{\text{def}}{=} \prod_{z \in \{A, B\}} z \\ &\stackrel{\text{def}}{=} \{f \in \text{Sets}(\{0, 1\}, A \cup B) \mid \text{we have } f(0) \in A \text{ and } f(1) \in B\} \\ &\cong \{ \{ \{a\}, \{a, b\} \} \in \mathcal{P}(\mathcal{P}(A \cup B)) \mid \text{we have } a \in A \text{ and } b \in B \}. \end{aligned}$$

- *The Cone.* The maps

$$\begin{aligned} pr_1 &: A \times B \rightarrow A, \\ pr_2 &: A \times B \rightarrow B \end{aligned}$$

defined by

$$\begin{aligned} pr_1(a, b) &\stackrel{\text{def}}{=} a, \\ pr_2(a, b) &\stackrel{\text{def}}{=} b \end{aligned}$$

for each $(a, b) \in A \times B$.

¹*Further Terminology:* Also called the **Cartesian product of A and B** or the **binary Cartesian product of A and B**, for emphasis.

This can also be thought of as the $(\mathbb{E}_{-1}, \mathbb{E}_{-1})$ -**tensor product of A and B**.

²In other words, $A \times B$ is the set whose elements are ordered pairs (a, b) with $a \in A$ and $b \in B$ as in [Definition 2.3.4.1](#)

PROOF 2.1.3.2 ► PROOF OF DEFINITION 2.1.3.1

We claim that $A \times B$ is the categorical product of A and B in Sets. Indeed, suppose we have a diagram of the form

$$\begin{array}{ccc} & P & \\ p_1 \swarrow & & \searrow p_2 \\ A & \longleftarrow A \times B \longrightarrow & B \\ pr_1 \longleftarrow & & \longrightarrow pr_2 \end{array}$$

in Sets. Then there exists a unique map $\phi: P \rightarrow A \times B$ making the diagram

$$\begin{array}{ccc}
 & P & \\
 p_1 \swarrow & \downarrow \phi \downarrow \exists! & \searrow p_2 \\
 A & \xleftarrow{\text{pr}_1} A \times B \xrightarrow{\text{pr}_2} & B
 \end{array}$$


commute, being uniquely determined by the conditions

$$\text{pr}_1 \circ \phi = p_1,$$

$$\text{pr}_2 \circ \phi = p_2$$

via

$$\phi(x) = (p_1(x), p_2(x))$$

for each $x \in P$. 

000U PROPOSITION 2.1.3.3 ► PROPERTIES OF PRODUCTS OF SETS

Let A, B, C , and X be sets.

000V 1. *Functoriality.* The assignments $A, B, (A, B) \mapsto A \times B$ define functors

$$A \times -: \text{Sets} \rightarrow \text{Sets},$$

$$- \times B: \text{Sets} \rightarrow \text{Sets},$$

$$-_1 \times -_2: \text{Sets} \times \text{Sets} \rightarrow \text{Sets},$$

where $-_1 \times -_2$ is the functor where

• *Action on Objects.* For each $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$, we have

$$[-_1 \times -_2](A, B) \stackrel{\text{def}}{=} A \times B.$$

• *Action on Morphisms.* For each $(A, B), (X, Y) \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$\times_{(A,B),(X,Y)}: \text{Sets}(A, X) \times \text{Sets}(B, Y) \rightarrow \text{Sets}(A \times B, X \times Y)$$

of \times at $((A, B), (X, Y))$ is defined by sending (f, g) to the function

$$f \times g: A \times B \rightarrow X \times Y$$

defined by

$$[f \times g](a, b) \stackrel{\text{def}}{=} (f(a), g(b))$$

for each $(a, b) \in A \times B$.

and where $A \times -$ and $- \times B$ are the partial functors of $-_1 \times -_2$ at $A, B \in \text{Obj}(\text{Sets})$.

000W

2. *Adjointness.* We have adjunctions

$$(A \times - \dashv \text{Hom}_{\text{Sets}}(A, -)) : \text{Sets} \begin{array}{c} \xrightarrow{A \times -} \\ \perp \\ \xleftarrow{\text{Hom}_{\text{Sets}}(A, -)} \end{array} \text{Sets},$$

$$(- \times B \dashv \text{Hom}_{\text{Sets}}(B, -)) : \text{Sets} \begin{array}{c} \xrightarrow{- \times B} \\ \perp \\ \xleftarrow{\text{Hom}_{\text{Sets}}(B, -)} \end{array} \text{Sets},$$

witnessed by bijections

$$\text{Hom}_{\text{Sets}}(A \times B, C) \cong \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(B, C)),$$

$$\text{Hom}_{\text{Sets}}(A \times B, C) \cong \text{Hom}_{\text{Sets}}(B, \text{Hom}_{\text{Sets}}(A, C)),$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

000X

3. *Associativity.* We have an isomorphism of sets

$$(A \times B) \times C \cong A \times (B \times C),$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

000Y

4. *Unitality.* We have isomorphisms of sets

$$\text{pt} \times A \cong A,$$

$$A \times \text{pt} \cong A,$$

natural in $A \in \text{Obj}(\text{Sets})$.

000Z

5. *Commutativity.* We have an isomorphism of sets

$$A \times B \cong B \times A,$$

natural in $A, B \in \text{Obj}(\text{Sets})$.

0010 6. *Annihilation With the Empty Set.* We have isomorphisms of sets

$$A \times \emptyset \cong \emptyset,$$

$$\emptyset \times A \cong \emptyset,$$

natural in $A \in \text{Obj}(\text{Sets})$.

0011 7. *Distributivity Over Unions.* We have isomorphisms of sets

$$A \times (B \cup C) = (A \times B) \cup (A \times C),$$

$$(A \cup B) \times C = (A \times C) \cup (B \times C).$$

0012 8. *Distributivity Over Intersections.* We have isomorphisms of sets

$$A \times (B \cap C) = (A \times B) \cap (A \times C),$$

$$(A \cap B) \times C = (A \times C) \cap (B \times C).$$

0013 9. *Middle-Four Exchange with Respect to Intersections.* We have an isomorphism of sets

$$(A \times B) \cap (C \times D) \cong (A \cap C) \times (B \cap D).$$

0014 10. *Distributivity Over Differences.* We have isomorphisms of sets

$$A \times (B \setminus C) = (A \times B) \setminus (A \times C),$$

$$(A \setminus B) \times C = (A \times C) \setminus (B \times C),$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

0015 11. *Distributivity Over Symmetric Differences.* We have isomorphisms of sets

$$A \times (B \Delta C) = (A \times B) \Delta (A \times C),$$

$$(A \Delta B) \times C = (A \times C) \Delta (B \times C),$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

0016 12. *Symmetric Monoidality.* The triple $(\text{Sets}, \times, \text{pt})$ is a symmetric monoidal category.

0017 13. *Symmetric Bimonoidality.* The quintuple $(\text{Sets}, \coprod, \emptyset, \times, \text{pt})$ is a symmetric bimonoidal category.

PROOF 2.1.3.4 ► PROOF OF PROPOSITION 2.1.3.3

Item 1: Functoriality

This follows from ?? of ??.

Item 2: Adjointness

We prove only that there's an adjunction $- \times B \dashv \text{Hom}_{\text{Sets}}(B, -)$, witnessed by a bijection

$$\text{Hom}_{\text{Sets}}(A \times B, C) \cong \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(B, C)),$$

natural in $B, C \in \text{Obj}(\text{Sets})$, as the proof of the existence of the adjunction $A \times - \dashv \text{Hom}_{\text{Sets}}(A, -)$ follows almost exactly in the same way.

- *Map I.* We define a map

$$\Phi_{B,C}: \text{Hom}_{\text{Sets}}(A \times B, C) \rightarrow \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(B, C)),$$

by sending a function

$$\xi: A \times B \rightarrow C$$

to the function

$$\begin{aligned} \xi^\dagger: A &\rightarrow \text{Hom}_{\text{Sets}}(B, C), \\ a &\mapsto (\xi_a^\dagger: B \rightarrow C), \end{aligned}$$

where we define

$$\xi_a^\dagger(b) \stackrel{\text{def}}{=} \xi(a, b)$$

for each $b \in B$. In terms of the $\llbracket a \mapsto f(a) \rrbracket$ notation of [Notation 1.1.1.2](#), we have

$$\xi^\dagger \stackrel{\text{def}}{=} \llbracket a \mapsto \llbracket b \mapsto \xi(a, b) \rrbracket \rrbracket.$$

- *Map II.* We define a map

$$\Psi_{B,C}: \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(B, C)) \rightarrow \text{Hom}_{\text{Sets}}(A \times B, C)$$

given by sending a function

$$\begin{aligned} \xi: A &\rightarrow \text{Hom}_{\text{Sets}}(B, C), \\ a &\mapsto (\xi_a: B \rightarrow C), \end{aligned}$$

to the function

$$\xi^\dagger: A \times B \rightarrow C$$

defined by

$$\begin{aligned} \xi^\dagger(a, b) &\stackrel{\text{def}}{=} \text{ev}_b(\text{ev}_a(\xi)) \\ &\stackrel{\text{def}}{=} \text{ev}_b(\xi_a) \\ &\stackrel{\text{def}}{=} \xi_a(b) \end{aligned}$$

for each $(a, b) \in A \times B$.

• *Invertibility I.* We claim that

$$\Psi_{A,B} \circ \Phi_{A,B} = \text{id}_{\text{Hom}_{\text{Sets}}(A \times B, C)}.$$

Indeed, given a function $\xi: A \times B \rightarrow C$, we have

$$\begin{aligned} [\Psi_{A,B} \circ \Phi_{A,B}](\xi) &= \Psi_{A,B}(\Phi_{A,B}(\xi)) \\ &= \Psi_{A,B}(\Phi_{A,B}(\llbracket (a, b) \mapsto \xi(a, b) \rrbracket\rrbracket)) \\ &= \Psi_{A,B}(\llbracket a \mapsto \llbracket b \mapsto \xi(a, b) \rrbracket \rrbracket\rrbracket) \\ &= \Psi_{A,B}(\llbracket a' \mapsto \llbracket b' \mapsto \xi(a', b') \rrbracket \rrbracket\rrbracket) \\ &= \llbracket (a, b) \mapsto \text{ev}_b(\text{ev}_a(\llbracket a' \mapsto \llbracket b' \mapsto \xi(a', b') \rrbracket \rrbracket\rrbracket)) \rrbracket\rrbracket \\ &= \llbracket (a, b) \mapsto \text{ev}_b(\llbracket b' \mapsto \xi(a, b') \rrbracket) \rrbracket\rrbracket \\ &= \llbracket (a, b) \mapsto \xi(a, b) \rrbracket \\ &= \xi. \end{aligned}$$

• *Invertibility II.* We claim that

$$\Phi_{A,B} \circ \Psi_{A,B} = \text{id}_{\text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(B, C))}.$$

Indeed, given a function

$$\begin{aligned} \xi: A &\rightarrow \text{Hom}_{\text{Sets}}(B, C), \\ a &\mapsto (\xi_a: B \rightarrow C), \end{aligned}$$

we have

$$\begin{aligned}
[\Phi_{A,B} \circ \Psi_{A,B}](\xi) &\stackrel{\text{def}}{=} \Phi_{A,B}(\Psi_{A,B}(\xi)) \\
&\stackrel{\text{def}}{=} \Phi_{A,B}(\llbracket (a, b) \mapsto \xi_a(b) \rrbracket) \\
&\stackrel{\text{def}}{=} \Phi_{A,B}(\llbracket (a', b') \mapsto \xi_{a'}(b') \rrbracket) \\
&\stackrel{\text{def}}{=} \llbracket a \mapsto \llbracket b \mapsto \text{ev}_{(a,b)}(\llbracket (a', b') \mapsto \xi_{a'}(b') \rrbracket) \rrbracket \rrbracket \\
&\stackrel{\text{def}}{=} \llbracket a \mapsto \llbracket b \mapsto \xi_a(b) \rrbracket \rrbracket \\
&\stackrel{\text{def}}{=} \llbracket a \mapsto \xi_a \rrbracket \\
&\stackrel{\text{def}}{=} \xi.
\end{aligned}$$

- *Naturality for Φ , Part I.* We need to show that, given a function $g: B \rightarrow B'$, the diagram

$$\begin{array}{ccc}
\text{Hom}_{\text{Sets}}(A \times B', C) & \xrightarrow{\Phi_{B',C}} & \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(B', C)), \\
\text{id}_A \times g^* \downarrow & & \downarrow (g^*)_* \\
\text{Hom}_{\text{Sets}}(A \times B, C) & \xrightarrow{\Phi_{B,C}} & \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(B, C))
\end{array}$$

commutes. Indeed, given a function

$$\xi: A \times B' \rightarrow C,$$

we have

$$\begin{aligned}
[\Phi_{B,C} \circ (\text{id}_A \times g^*)](\xi) &= \Phi_{B,C}([\text{id}_A \times g^*](\xi)) \\
&= \Phi_{B,C}(\xi(-1, g(-2))) \\
&= [\xi(-1, g(-2))]^\dagger \\
&= \xi_{-1}^\dagger(g(-2)) \\
&= (g^*)_* (\xi^\dagger) \\
&= (g^*)_* (\Phi_{B',C}(\xi)) \\
&= [(g^*)_* \circ \Phi_{B',C}](\xi).
\end{aligned}$$

Alternatively, using the $\llbracket a \mapsto f(a) \rrbracket$ notation of [Notation 1.1.1.2](#), we

have

$$\begin{aligned}
[\Phi_{B,C} \circ (\text{id}_A \times g^*)](\xi) &= \Phi_{B,C}([\text{id}_A \times g^*](\xi)) \\
&= \Phi_{B,C}([\text{id}_A \times g^*](\llbracket (a, b') \mapsto \xi(a, b') \rrbracket)) \\
&= \Phi_{B,C}(\llbracket (a, b) \mapsto \xi(a, g(b)) \rrbracket) \\
&= \llbracket a \mapsto \llbracket b \mapsto \xi(a, g(b)) \rrbracket \rrbracket \\
&= \llbracket a \mapsto g^*(\llbracket b' \mapsto \xi(a, b') \rrbracket) \rrbracket \\
&= (g^*)_* (\llbracket a \mapsto \llbracket b' \mapsto \xi(a, b') \rrbracket \rrbracket) \\
&= (g^*)_* (\Phi_{B',C}(\llbracket (a, b') \mapsto \xi(a, b') \rrbracket)) \\
&= (g^*)_* (\Phi_{B',C}(\xi)) \\
&= [(g^*)_* \circ \Phi_{B',C}](\xi).
\end{aligned}$$

- *Naturality for Φ , Part II.* We need to show that, given a function $h: C \rightarrow C'$, the diagram

$$\begin{array}{ccc}
\text{Hom}_{\text{Sets}}(A \times B, C) & \xrightarrow{\Phi_{B,C}} & \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(B, C)), \\
\downarrow h_* & & \downarrow (h_*)_* \\
\text{Hom}_{\text{Sets}}(A \times B, C') & \xrightarrow{\Phi_{B,C'}} & \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(B, C'))
\end{array}$$

commutes. Indeed, given a function

$$\xi: A \times B \rightarrow C,$$

we have

$$\begin{aligned}
[\Phi_{B,C} \circ h_*](\xi) &= \Phi_{B,C}(h_*(\xi)) \\
&= \Phi_{B,C}(h_*(\llbracket (a, b) \mapsto \xi(a, b) \rrbracket)) \\
&= \Phi_{B,C}(\llbracket (a, b) \mapsto h(\xi(a, b)) \rrbracket) \\
&= \llbracket a \mapsto \llbracket b \mapsto h(\xi(a, b)) \rrbracket \rrbracket \\
&= \llbracket a \mapsto h_* (\llbracket b \mapsto \xi(a, b) \rrbracket) \rrbracket \\
&= (h_*)_* (\llbracket a \mapsto \llbracket b \mapsto \xi(a, b) \rrbracket \rrbracket) \\
&= (h_*)_* (\Phi_{B,C}(\llbracket (a, b) \mapsto \xi(a, b) \rrbracket)) \\
&= (h_*)_* (\Phi_{B,C}(\xi)) \\
&= [(h_*)_* \circ \Phi_{B,C}](\xi).
\end{aligned}$$

- *Naturality for Ψ* . Since Φ is natural in each argument and Φ is a componentwise inverse to Ψ in each argument, it follows from [Item 2 of Proposition 8.8.6.2](#) that Ψ is also natural in each argument.

Item 3: Associativity

See [[Pro24a](#)].

Item 4: Unitality

Clear.

Item 5: Commutativity

See [[Pro24b](#)].

Item 6: Annihilation With the Empty Set

See [[Pro24f](#)].

Item 7: Distributivity Over Unions

See [[Pro24e](#)].

Item 8: Distributivity Over Intersections

See [[Pro24g](#), Corollary 1].

Item 9: Middle-Four Exchange With Respect to Intersections

See [[Pro24g](#), Corollary 1].

Item 10: Distributivity Over Differences

See [[Pro24c](#)].

Item 11: Distributivity Over Symmetric Differences

See [[Pro24d](#)].

Item 12: Symmetric Monoidality

See [[MO 382264](#)].

Item 13: Symmetric Bimonoidality

Omitted. 

0018 2.1.4 Pullbacks

Let A , B , and C be sets and let $f: A \rightarrow C$ and $g: B \rightarrow C$ be functions.

0019 DEFINITION 2.1.4.1 ► PULLBACKS OF SETS

The **pullback of A and B over C along f and g** ¹ is the pair² $(A \times_C B, \{pr_1, pr_2\})$ consisting of:

- *The Limit.* The set $A \times_C B$ defined by

$$A \times_C B \stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid f(a) = g(b)\}.$$

- *The Cone.* The maps

$$\text{pr}_1 : A \times_C B \rightarrow A,$$

$$\text{pr}_2 : A \times_C B \rightarrow B$$

defined by

$$\text{pr}_1(a, b) \stackrel{\text{def}}{=} a,$$

$$\text{pr}_2(a, b) \stackrel{\text{def}}{=} b$$

for each $(a, b) \in A \times_C B$.

¹Further Terminology: Also called the **fibre product of A and B over C along f and g** .

²Further Notation: Also written $A \times_{f,C,g} B$.

PROOF 2.1.4.2 ► PROOF OF DEFINITION 2.1.4.1


We claim that $A \times_C B$ is the categorical pullback of A and B over C with respect to (f, g) in Sets. First we need to check that the relevant pullback diagram commutes, i.e. that we have

$$f \circ \text{pr}_1 = g \circ \text{pr}_2, \quad \begin{array}{ccc} A \times_C B & \xrightarrow{\text{pr}_2} & B \\ \text{pr}_1 \downarrow & & \downarrow g \\ A & \xrightarrow{f} & C. \end{array}$$


Indeed, given $(a, b) \in A \times_C B$, we have

$$\begin{aligned} [f \circ \text{pr}_1](a, b) &= f(\text{pr}_1(a, b)) \\ &= f(a) \\ &= g(b) \\ &= g(\text{pr}_2(a, b)) \\ &= [g \circ \text{pr}_2](a, b), \end{aligned}$$

where $f(a) = g(b)$ since $(a, b) \in A \times_C B$. Next, we prove that $A \times_C B$ satisfies the universal property of the pullback. Suppose we have a diagram of the form



in Sets. Then there exists a unique map $\phi: P \rightarrow A \times_C B$ making the diagram



commute, being uniquely determined by the conditions

$$\text{pr}_1 \circ \phi = p_1,$$

$$\text{pr}_2 \circ \phi = p_2$$

via

$$\phi(x) = (p_1(x), p_2(x))$$

for each $x \in P$, where we note that $(p_1(x), p_2(x)) \in A \times B$ indeed lies in $A \times_C B$ by the condition

$$f \circ p_1 = g \circ p_2,$$

which gives

$$f(p_1(x)) = g(p_2(x))$$

for each $x \in P$, so that $(p_1(x), p_2(x)) \in A \times_C B$. ▢

001A **EXAMPLE 2.1.4.3 ▶ EXAMPLES OF PULLBACKS OF SETS**

Here are some examples of pullbacks of sets.

001B 1. *Unions via Intersections.* Let $A, B \subset X$. We have a bijection of sets

$$A \cap B \cong A \times_{A \cup B} B,$$


$$\begin{array}{ccc} A \cap B & \longrightarrow & B \\ \downarrow \lrcorner & & \downarrow i_B \\ A & \xrightarrow{i_A} & A \cup B \end{array}$$

PROOF 2.1.4.4 ▶ PROOF OF EXAMPLE 2.1.4.3

Item 1: Unions via Intersections

Indeed, we have

$$\begin{aligned} A \times_{A \cup B} B &\cong \{(x, y) \in A \times B \mid x = y\} \\ &\cong A \cap B. \end{aligned}$$

This finishes the proof. 


001C **PROPOSITION 2.1.4.5 ▶ PROPERTIES OF PULLBACKS OF SETS**

Let A, B, C , and X be sets.

001D 1. *Functoriality.* The assignment $(A, B, C, f, g) \mapsto A \times_{f, C, g} B$ defines a functor


$$-_1 \times_{-3} -_1: \text{Fun}(\mathcal{P}, \text{Sets}) \rightarrow \text{Sets},$$

where \mathcal{P} is the category that looks like this:



In particular, the action on morphisms of $-_1 \times_{-3} -_1$ is given by sending


a morphism



in $\text{Fun}(\mathcal{P}, \text{Sets})$ to the map $\xi: A \times_C B \xrightarrow{\exists!} A' \times_C B'$ given by

$$\xi(a, b) \stackrel{\text{def}}{=} (\phi(a), \psi(b))$$


for each $(a, b) \in A \times_C B$, which is the unique map making the diagram



commute.

001E


2. *Associativity.* Given a diagram




in Sets , we have isomorphisms of sets

$$(A \times_X B) \times_Y C \cong (A \times_X B) \times_B (B \times_Y C) \cong A \times_X (B \times_Y C),$$


where these pullbacks are built as in the diagrams




001F 3. *Unitality.* We have isomorphisms of sets




001G 4. *Commutativity.* We have an isomorphism of sets



001H 5. *Annihilation With the Empty Set.* We have isomorphisms of sets



001J 6. *Interaction With Products.* We have an isomorphism of sets



001K 7. *Symmetric Monoidality.* The triple (Sets, \times_X , X) is a symmetric monoidal category.

PROOF 2.1.4.6 ► PROOF OF PROPOSITION 2.1.4.5

Item 1: Functoriality

This is a special case of functoriality of co/limits, ?? of ??, with the explicit expression for ξ following from the commutativity of the cube pullback diagram.

Item 2: Associativity

Indeed, we have

$$\begin{aligned}
 (A \times_X B) \times_Y C &\cong \{(a, b), c\} \in (A \times_X B) \times C \mid h(b) = k(c)\} \\
 &\cong \{(a, b), c\} \in (A \times B) \times C \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\
 &\cong \{(a, (b, c)) \in A \times (B \times C) \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\
 &\cong \{(a, (b, c)) \in A \times (B \times_Y C) \mid f(a) = g(b)\} \\
 &\cong A \times_X (B \times_Y C)
 \end{aligned}$$

and

$$\begin{aligned}
 (A \times_X B) \times_B (B \times_Y C) &\cong \{(a, b), (b', c)\} \in (A \times_X B) \times (B \times_Y C) \mid b = b'\} \\
 &\cong \left\{ \{(a, b), (b', c)\} \in (A \times B) \times (B \times C) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } h(b') = k(c) \end{array} \right\} \\
 &\cong \left\{ (a, (b, (b', c))) \in A \times (B \times (B \times C)) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } h(b') = k(c) \end{array} \right\} \\
 &\cong \left\{ (a, ((b, b'), c)) \in A \times ((B \times B) \times C) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } h(b') = k(c) \end{array} \right\} \\
 &\cong \left\{ (a, ((b, b'), c)) \in A \times ((B \times_B B) \times C) \mid \begin{array}{l} f(a) = g(b) \text{ and } \\ h(b') = k(c) \end{array} \right\} \\
 &\cong \{(a, (b, c)) \in A \times (B \times C) \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\
 &\cong A \times_X (B \times_Y C),
 \end{aligned}$$

where we have used **Item 3** for the isomorphism $B \times_B B \cong B$.

Item 3: Unitality

Indeed, we have

$$\begin{aligned}
 X \times_X A &\cong \{(x, a) \in X \times A \mid f(a) = x\}, \\
 A \times_X X &\cong \{(a, x) \in X \times A \mid f(a) = x\},
 \end{aligned}$$

which are isomorphic to A via the maps $(x, a) \mapsto a$ and $(a, x) \mapsto a$.

Item 4: Commutativity

Clear.

Item 5: Annihilation With the Empty Set

Clear.

Item 6: Interaction With Products

Clear.

Item 7: Symmetric Monoidality

Omitted. 

001L 2.1.5 Equalisers

Let A and B be sets and let $f, g: A \rightrightarrows B$ be functions.

001M DEFINITION 2.1.5.1 ► EQUALISERS OF SETS

The **equaliser of f and g** is the pair $(\text{Eq}(f, g), \text{eq}(f, g))$ consisting of:

- *The Limit.* The set $\text{Eq}(f, g)$ defined by

$$\text{Eq}(f, g) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = g(a)\}.$$

- *The Cone.* The inclusion map

$$\text{eq}(f, g): \text{Eq}(f, g) \hookrightarrow A.$$

PROOF 2.1.5.2 ► PROOF OF DEFINITION 2.1.5.1

We claim that $\text{Eq}(f, g)$ is the categorical equaliser of f and g in Sets . First we need to check that the relevant equaliser diagram commutes, i.e. that we have

$$f \circ \text{eq}(f, g) = g \circ \text{eq}(f, g),$$

which indeed holds by the definition of the set $\text{Eq}(f, g)$. Next, we prove that $\text{Eq}(f, g)$ satisfies the universal property of the equaliser. Suppose we have a diagram of the form

$$\begin{array}{ccc} \text{Eq}(f, g) & \xrightarrow{\text{eq}(f, g)} & A & \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} & B \\ & \nearrow e & & & \\ E & & & & \end{array}$$

in Sets. Then there exists a unique map $\phi: E \rightarrow \text{Eq}(f, g)$ making the diagram

$$\begin{array}{ccc} \text{Eq}(f, g) & \xrightarrow{\text{eq}(f, g)} & A \xrightarrow[f]{g} B \\ \uparrow \phi \exists! & \nearrow e & \\ E & & \end{array}$$

commute, being uniquely determined by the condition

$$\text{eq}(f, g) \circ \phi = e$$

via

$$\phi(x) = e(x)$$

for each $x \in E$, where we note that $e(x) \in A$ indeed lies in $\text{Eq}(f, g)$ by the condition

$$f \circ e = g \circ e,$$

which gives

$$f(e(x)) = g(e(x))$$

for each $x \in E$, so that $e(x) \in \text{Eq}(f, g)$. ▢

001N PROPOSITION 2.1.5.3 ► PROPERTIES OF EQUALISERS OF SETS

Let A, B , and C be sets.

001P 1. *Associativity.* We have isomorphisms of sets¹

$$\underbrace{\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h))}_{=\text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))} \cong \text{Eq}(f, g, h) \cong \underbrace{\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g))}_{=\text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))},$$

where $\text{Eq}(f, g, h)$ is the limit of the diagram

$$\begin{array}{ccc} & f & \\ & \rightarrow & \\ A & \xrightarrow{g} & B \\ & \xrightarrow{h} & \end{array}$$

in Sets, being explicitly given by

$$\text{Eq}(f, g, h) \cong \{a \in A \mid f(a) = g(a) = h(a)\}.$$

001Q 2. *Unitality.* We have an isomorphism of sets

$$\text{Eq}(f, f) \cong A.$$

001R

3. *Commutativity.* We have an isomorphism of sets

$$\text{Eq}(f, g) \cong \text{Eq}(g, f).$$

001S

4. *Interaction With Composition.* Let

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B \begin{array}{c} \xrightarrow{h} \\ \xrightarrow{k} \end{array} C$$

be functions. We have an inclusion of sets

$$\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g)) \subset \text{Eq}(h \circ f, k \circ g),$$

where $\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g))$ is the equaliser of the composition

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B \begin{array}{c} \xrightarrow{h} \\ \xrightarrow{k} \end{array} C.$$

¹That is, the following three ways of forming “the” equaliser of (f, g, h) agree:

(a) Take the equaliser of (f, g, h) , i.e. the limit of the diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \\ \xrightarrow{h} \end{array} B$$

in Sets.

(b) First take the equaliser of f and g , forming a diagram

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B$$

and then take the equaliser of the composition

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{h} \end{array} B,$$

obtaining a subset

$$\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)) = \text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))$$

of $\text{Eq}(f, g)$.

(c) First take the equaliser of g and h , forming a diagram

$$\text{Eq}(g, h) \xrightarrow{\text{eq}(g, h)} A \begin{array}{c} \xrightarrow{g} \\ \xrightarrow{h} \end{array} B$$

and then take the equaliser of the composition

$$\text{Eq}(g, h) \xrightarrow{\text{eq}(g, h)} A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B,$$

obtaining a subset

$$\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) = \text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))$$

of $\text{Eq}(g, h)$.

PROOF 2.1.5.4 ► PROOF OF PROPOSITION 2.1.5.3

Item 1: Associativity

We first prove that $\text{Eq}(f, g, h)$ is indeed given by

$$\text{Eq}(f, g, h) \cong \{a \in A \mid f(a) = g(a) = h(a)\}.$$

Indeed, suppose we have a diagram of the form

$$\begin{array}{ccc} \text{Eq}(f, g, h) & \xrightarrow{\text{eq}(f, g, h)} & A \\ & \searrow e & \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \\ \xrightarrow{h} \end{array} \\ E & & B \end{array}$$

in Sets. Then there exists a unique map $\phi: E \rightarrow \text{Eq}(f, g, h)$, uniquely determined by the condition

$$\text{eq}(f, g) \circ \phi = e$$

being necessarily given by

$$\phi(x) = e(x)$$

for each $x \in E$, where we note that $e(x) \in A$ indeed lies in $\text{Eq}(f, g, h)$ by the condition

$$f \circ e = g \circ e = h \circ e,$$

which gives

$$f(e(x)) = g(e(x)) = h(e(x))$$

for each $x \in E$, so that $e(x) \in \text{Eq}(f, g, h)$.

We now check the equalities

$$\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) \cong \text{Eq}(f, g, h) \cong \text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)).$$

Indeed, we have

$$\begin{aligned} \text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) &\cong \{x \in \text{Eq}(g, h) \mid [f \circ \text{eq}(g, h)](a) = [g \circ \text{eq}(g, h)](a)\} \\ &\cong \{x \in \text{Eq}(g, h) \mid f(a) = g(a)\} \\ &\cong \{x \in A \mid f(a) = g(a) \text{ and } g(a) = h(a)\} \\ &\cong \{x \in A \mid f(a) = g(a) = h(a)\} \\ &\cong \text{Eq}(f, g, h). \end{aligned}$$

Similarly, we have

$$\begin{aligned}
 \text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)) &\cong \{x \in \text{Eq}(f, g) \mid [f \circ \text{eq}(f, g)](a) = [h \circ \text{eq}(f, g)](a)\} \\
 &\cong \{x \in \text{Eq}(f, g) \mid f(a) = h(a)\} \\
 &\cong \{x \in A \mid f(a) = h(a) \text{ and } f(a) = g(a)\} \\
 &\cong \{x \in A \mid f(a) = g(a) = h(a)\} \\
 &\cong \text{Eq}(f, g, h).
 \end{aligned}$$

Item 2: Unitality

Clear.

Item 3: Commutativity

Clear.


Item 4: Interaction With Composition

Indeed, we have

$$\begin{aligned}
 \text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g)) &\cong \{a \in \text{Eq}(f, g) \mid h(f(a)) = k(g(a))\} \\
 &\cong \{a \in A \mid f(a) = g(a) \text{ and } h(f(a)) = k(g(a))\}.
 \end{aligned}$$

and

$$\text{Eq}(h \circ f, k \circ g) \cong \{a \in A \mid h(f(a)) = k(g(a))\},$$

and thus there's an inclusion from $\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g))$ to $\text{Eq}(h \circ f, k \circ g)$. 

001T 2.2 Colimits of Sets

001U 2.2.1 The Initial Set

001V DEFINITION 2.2.1.1 ► THE INITIAL SET

The **initial set** is the pair $(\emptyset, \{\iota_A\}_{A \in \text{Obj}(\text{Sets})})$ consisting of:

- *The Limit.* The empty set \emptyset of [Definition 2.3.1.1](#).
- *The Cone.* The collection of maps

$$\{\iota_A: \emptyset \rightarrow A\}_{A \in \text{Obj}(\text{Sets})}$$

given by the inclusion maps from \emptyset to A .


PROOF 2.2.1.2 ► PROOF OF DEFINITION 2.2.1.1

We claim that \emptyset is the initial object of Sets. Indeed, suppose we have a diagram of the form

$$\emptyset \quad A$$

in Sets. Then there exists a unique map $\phi: \emptyset \rightarrow A$ making the diagram

$$\emptyset \xrightarrow[\exists!]{\phi} A$$

commute, namely the inclusion map ι_A . 

001W 2.2.2 Coproducts of Families of Sets

Let $\{A_i\}_{i \in I}$ be a family of sets.

001X DEFINITION 2.2.2.1 ► DISJOINT UNIONS OF FAMILIES

The **disjoint union of the family** $\{A_i\}_{i \in I}$ is the pair $(\coprod_{i \in I} A_i, \{\text{inj}_i\}_{i \in I})$ consisting of:

- *The Colimit.* The set $\coprod_{i \in I} A_i$ defined by

$$\coprod_{i \in I} A_i \stackrel{\text{def}}{=} \left\{ (i, x) \in I \times \left(\bigcup_{i \in I} A_i \right) \mid x \in A_i \right\}.$$

- *The Cocone.* The collection

$$\left\{ \text{inj}_i: A_i \rightarrow \coprod_{i \in I} A_i \right\}_{i \in I}$$

of maps given by

$$\text{inj}_i(x) \stackrel{\text{def}}{=} (i, x)$$

for each $x \in A_i$ and each $i \in I$.

PROOF 2.2.2.2 ► PROOF OF DEFINITION 2.2.2.1

We claim that $\coprod_{i \in I} A_i$ is the categorical coproduct of $\{A_i\}_{i \in I}$ in Sets. Indeed, suppose we have, for each $i \in I$, a diagram of the form

$$\begin{array}{ccc} & & C \\ & \nearrow \iota_i & \\ A_i & \xrightarrow{\text{inj}_i} & \coprod_{i \in I} A_i \end{array}$$

in Sets. Then there exists a unique map $\phi: \coprod_{i \in I} A_i \rightarrow C$ making the diagram

$$\begin{array}{ccc} & & C \\ & \nearrow \iota_i & \uparrow \phi \exists! \\ A_i & \xrightarrow{\text{inj}_i} & \coprod_{i \in I} A_i \end{array}$$

commute, being uniquely determined by the condition $\phi \circ \text{inj}_i = \iota_i$ for each $i \in I$ via

$$\phi((i, x)) = \iota_i(x)$$

for each $(i, x) \in \coprod_{i \in I} A_i$. ▢

001Y

PROPOSITION 2.2.2.3 ► PROPERTIES OF COPRODUCTS OF FAMILIES OF SETS

Let $\{A_i\}_{i \in I}$ be a family of sets.

001Z

1. *Functoriality.* The assignment $\{A_i\}_{i \in I} \mapsto \coprod_{i \in I} A_i$ defines a functor

$$\coprod_{i \in I}: \text{Fun}(I_{\text{disc}}, \text{Sets}) \rightarrow \text{Sets}$$

where

• *Action on Objects.* For each $(A_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$, we have

$$\left[\coprod_{i \in I} \right] ((A_i)_{i \in I}) \stackrel{\text{def}}{=} \coprod_{i \in I} A_i$$

• *Action on Morphisms.* For each $(A_i)_{i \in I}, (B_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$

$\text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$, the action on Hom-sets

$$\left(\coprod_{i \in I} \right)_{(A_i)_{i \in I}, (B_i)_{i \in I}} : \text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I}) \rightarrow \text{Sets} \left(\coprod_{i \in I} A_i, \coprod_{i \in I} B_i \right)$$

of $\coprod_{i \in I}$ at $((A_i)_{i \in I}, (B_i)_{i \in I})$ is defined by sending a map

$$\{f_i : A_i \rightarrow B_i\}_{i \in I}$$

in $\text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I})$ to the map of sets

$$\coprod_{i \in I} f_i : \coprod_{i \in I} A_i \rightarrow \coprod_{i \in I} B_i$$

defined by

$$\left[\coprod_{i \in I} f_i \right] (i, a) \stackrel{\text{def}}{=} f_i(a)$$

for each $(i, a) \in \coprod_{i \in I} A_i$.

PROOF 2.2.2.4 ► PROOF OF PROPOSITION 2.2.2.3

Item 1: Functoriality

This follows from ?? of ??.



0020 **2.2.3 Binary Coproducts**

Let A and B be sets.

0021 **DEFINITION 2.2.3.1 ► COPRODUCTS OF SETS**

The **coproduct**¹ of A and B is the pair $(A \amalg B, \{\text{inj}_1, \text{inj}_2\})$ consisting of:

- *The Colimit.* The set $A \amalg B$ defined by

$$\begin{aligned} A \amalg B &\stackrel{\text{def}}{=} \coprod_{z \in \{A, B\}} z \\ &\cong \{(0, a) \mid a \in A\} \cup \{(1, b) \mid b \in B\}. \end{aligned}$$

• *The Cocone.* The maps

$$\text{inj}_1: A \rightarrow A \amalg B,$$

$$\text{inj}_2: B \rightarrow A \amalg B,$$

given by

$$\text{inj}_1(a) \stackrel{\text{def}}{=} (0, a),$$

$$\text{inj}_2(b) \stackrel{\text{def}}{=} (1, b),$$

for each $a \in A$ and each $b \in B$.

¹*Further Terminology:* Also called the **disjoint union of A and B** , or the **binary disjoint union of A and B** , for emphasis.

PROOF 2.2.3.2 ► PROOF OF DEFINITION 2.2.3.1

We claim that $A \amalg B$ is the categorical coproduct of A and B in Sets. Indeed, suppose we have a diagram of the form

$$\begin{array}{ccc} & C & \\ \iota_A \nearrow & & \nwarrow \iota_B \\ A & \xrightarrow{\text{inj}_A} & A \amalg B \xleftarrow{\text{inj}_B} B \end{array}$$

in Sets. Then there exists a unique map $\phi: A \amalg B \rightarrow C$ making the diagram

$$\begin{array}{ccc} & C & \\ \iota_A \nearrow & \uparrow \phi \exists! & \nwarrow \iota_B \\ A & \xrightarrow{\text{inj}_A} & A \amalg B \xleftarrow{\text{inj}_B} B \end{array}$$

commute, being uniquely determined by the conditions

$$\phi \circ \text{inj}_A = \iota_A,$$

$$\phi \circ \text{inj}_B = \iota_B$$

via

$$\phi(x) = \begin{cases} \iota_A(a) & \text{if } x = (0, a), \\ \iota_B(b) & \text{if } x = (1, b) \end{cases}$$

for each $x \in A \amalg B$. ▢

0022 **PROPOSITION 2.2.3.3 ► PROPERTIES OF COPRODUCTS OF SETS**

Let A, B, C , and X be sets.

0023 1. *Functoriality.* The assignment $A, B, (A, B) \mapsto A \amalg B$ defines functors

$$\begin{aligned} A \amalg - &: \text{Sets} \rightarrow \text{Sets}, \\ - \amalg B &: \text{Sets} \rightarrow \text{Sets}, \\ -_1 \amalg -_2 &: \text{Sets} \times \text{Sets} \rightarrow \text{Sets}, \end{aligned}$$

where $-_1 \amalg -_2$ is the functor where

· *Action on Objects.* For each $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$, we have

$$[-_1 \amalg -_2](A, B) \stackrel{\text{def}}{=} A \amalg B.$$

· *Action on Morphisms.* For each $(A, B), (X, Y) \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$\amalg_{(A,B),(X,Y)} : \text{Sets}(A, X) \times \text{Sets}(B, Y) \rightarrow \text{Sets}(A \amalg B, X \amalg Y)$$

of \amalg at $((A, B), (X, Y))$ is defined by sending (f, g) to the function

$$f \amalg g : A \amalg B \rightarrow X \amalg Y$$

defined by

$$[f \amalg g](x) \stackrel{\text{def}}{=} \begin{cases} (0, f(a)) & \text{if } x = (0, a), \\ (1, g(b)) & \text{if } x = (1, b), \end{cases}$$

for each $x \in A \amalg B$.

and where $A \amalg -$ and $- \amalg B$ are the partial functors of $-_1 \amalg -_2$ at $A, B \in \text{Obj}(\text{Sets})$.

0024 2. *Associativity.* We have an isomorphism of sets

$$(A \amalg B) \amalg C \cong A \amalg (B \amalg C),$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

0025 3. *Unitality.* We have isomorphisms of sets

$$\begin{aligned} A \amalg \emptyset &\cong A, \\ \emptyset \amalg A &\cong A, \end{aligned}$$

natural in $A \in \text{Obj}(\text{Sets})$.

0026

4. *Commutativity.* We have an isomorphism of sets

$$A \amalg B \cong B \amalg A,$$

natural in $A, B \in \text{Obj}(\text{Sets})$.

0027

5. *Symmetric Monoidality.* The triple $(\text{Sets}, \amalg, \emptyset)$ is a symmetric monoidal category.

PROOF 2.2.3.4 ► PROOF OF PROPOSITION 2.2.3.3

Item 1: Functoriality

This follows from ?? of ??.

Item 2: Associativity

Clear.

Item 3: Unitality

Clear.

Item 4: Commutativity

Clear.

Item 5: Symmetric Monoidality

Omitted. 

0028 2.2.4 Pushouts

Let A, B , and C be sets and let $f: C \rightarrow A$ and $g: C \rightarrow B$ be functions.

0029

DEFINITION 2.2.4.1 ► PUSHOUTS OF SETS

The **pushout of A and B over C along f and g** ¹ is the pair² $(A \amalg_C B, \{\text{inj}_1, \text{inj}_2\})$ consisting of:

- *The Colimit.* The set $A \amalg_C B$ defined by

$$A \amalg_C B \stackrel{\text{def}}{=} A \amalg B / \sim_C,$$

where \sim_C is the equivalence relation on $A \amalg B$ generated by $(0, f(c)) \sim_C (1, g(c))$.

• *The Cocone.* The maps

$$\text{inj}_1 : A \rightarrow A \amalg_C B,$$

$$\text{inj}_2 : B \rightarrow A \amalg_C B$$

given by

$$\text{inj}_1(a) \stackrel{\text{def}}{=} [(0, a)]$$

$$\text{inj}_2(b) \stackrel{\text{def}}{=} [(1, b)]$$

for each $a \in A$ and each $b \in B$.

¹*Further Terminology:* Also called the **fibre coproduct of A and B over C along f and g .**

²*Further Notation:* Also written $A \amalg_{f,C,g} B$.

PROOF 2.2.4.2 ► PROOF OF DEFINITION 2.2.4.1

We claim that $A \amalg_C B$ is the categorical pushout of A and B over C with respect to (f, g) in Sets. First we need to check that the relevant pushout diagram commutes, i.e. that we have

$$\text{inj}_1 \circ f = \text{inj}_2 \circ g,$$


$$\begin{array}{ccc} A \amalg_C B & \xleftarrow{\text{inj}_2} & B \\ \text{inj}_1 \uparrow & & \uparrow g \\ A & \xleftarrow{f} & C. \end{array}$$

Indeed, given $c \in C$, we have


$$\begin{aligned} [\text{inj}_1 \circ f](c) &= \text{inj}_1(f(c)) \\ &= [(0, f(c))] \\ &= [(1, g(c))] \\ &= \text{inj}_2(g(c)) \\ &= [\text{inj}_2 \circ g](c), \end{aligned}$$

where $[(0, f(c))] = [(1, g(c))]$ by the definition of the relation \sim on $A \amalg B$. Next, we prove that $A \amalg_C B$ satisfies the universal property of the pushout.

Suppose we have a diagram of the form



in Sets. Then there exists a unique map $\phi: A \amalg_C B \rightarrow P$ making the diagram



commute, being uniquely determined by the conditions

$$\begin{aligned}\phi \circ \text{inj}_1 &= \iota_1, \\ \phi \circ \text{inj}_2 &= \iota_2\end{aligned}$$

via

$$\phi(x) = \begin{cases} \iota_1(a) & \text{if } x = [(0, a)], \\ \iota_2(b) & \text{if } x = [(1, b)] \end{cases}$$

for each $x \in A \amalg_C B$, where the well-definedness of ϕ is guaranteed by the equality $\iota_1 \circ f = \iota_2 \circ g$ and the definition of the relation \sim on $A \amalg B$ as follows:

1. *Case 1:* Suppose we have $x = [(0, a)] = [(0, a')]$ for some $a, a' \in A$. Then, by [Remark 2.2.4.3](#), we have a sequence

$$(0, a) \sim' x_1 \sim' \dots \sim' x_n \sim' (0, a').$$

2. *Case 2:* Suppose we have $x = [(1, b)] = [(1, b')]$ for some $b, b' \in B$. Then, by [Remark 2.2.4.3](#), we have a sequence

$$(1, b) \sim' x_1 \sim' \dots \sim' x_n \sim' (1, b').$$

3. *Case 3:* Suppose we have $x = [(0, a)] = [(1, b)]$ for some $a \in A$ and $b \in B$. Then, by [Remark 2.2.4.3](#), we have a sequence

$$(0, a) \sim' x_1 \sim' \cdots \sim' x_n \sim' (1, b).$$

In all these cases, we declare $x \sim' y$ iff there exists some $c \in C$ such that $x = (0, f(c))$ and $y = (1, g(c))$ or $x = (1, g(c))$ and $y = (0, f(c))$. Then, the equality $\iota_1 \circ f = \iota_2 \circ g$ gives


$$\begin{aligned} \phi([x]) &= \phi([(0, f(c))]) \\ &\stackrel{\text{def}}{=} \iota_1(f(c)) \\ &= \iota_2(g(c)) \\ &\stackrel{\text{def}}{=} \phi([(1, g(c))]) \\ &= \phi([y]), \end{aligned}$$

with the case where $x = (1, g(c))$ and $y = (0, f(c))$ similarly giving $\phi([x]) = \phi([y])$. Thus, if $x \sim' y$, then $\phi([x]) = \phi([y])$. Applying this equality pairwise to the sequences

$$\begin{aligned} (0, a) \sim' x_1 \sim' \cdots \sim' x_n \sim' (0, a'), \\ (1, b) \sim' x_1 \sim' \cdots \sim' x_n \sim' (1, b'), \\ (0, a) \sim' x_1 \sim' \cdots \sim' x_n \sim' (1, b) \end{aligned}$$

gives

$$\begin{aligned} \phi([(0, a)]) &= \phi([(0, a')]), \\ \phi([(1, b)]) &= \phi([(1, b')]), \\ \phi([(0, a)]) &= \phi([(1, b)]), \end{aligned}$$

showing ϕ to be well-defined. 

002A

REMARK 2.2.4.3 ► UNWINDING DEFINITION 2.2.4.1

In detail, by [Construction 7.4.2.2](#), the relation \sim of [Definition 2.2.4.1](#) is given by declaring $a \sim b$ iff one of the following conditions is satisfied:

- We have $a, b \in A$ and $a = b$;
- We have $a, b \in B$ and $a = b$;
- There exist $x_1, \dots, x_n \in A \amalg B$ such that $a \sim' x_1 \sim' \cdots \sim' x_n \sim' b$,

where we declare $x \sim' y$ if one of the following conditions is satisfied:

1. There exists $c \in C$ such that $x = (0, f(c))$ and $y = (1, g(c))$.
2. There exists $c \in C$ such that $x = (1, g(c))$ and $y = (0, f(c))$.

That is: we require the following condition to be satisfied:

- (★) There exist $x_1, \dots, x_n \in A \amalg B$ satisfying the following conditions:
1. There exists $c_0 \in C$ satisfying one of the following conditions:
 - (a) We have $a = f(c_0)$ and $x_1 = g(c_0)$.
 - (b) We have $a = g(c_0)$ and $x_1 = f(c_0)$.
 2. For each $1 \leq i \leq n - 1$, there exists $c_i \in C$ satisfying one of the following conditions:
 - (a) We have $x_i = f(c_i)$ and $x_{i+1} = g(c_i)$.
 - (b) We have $x_i = g(c_i)$ and $x_{i+1} = f(c_i)$.
 3. There exists $c_n \in C$ satisfying one of the following conditions:
 - (a) We have $x_n = f(c_n)$ and $b = g(c_n)$.
 - (b) We have $x_n = g(c_n)$ and $b = f(c_n)$.

002B **EXAMPLE 2.2.4.4 ► EXAMPLES OF PUSHOUTS OF SETS**

Here are some examples of pushouts of sets.

002C 1. *Wedge Sums of Pointed Sets.* The wedge sum of two pointed sets of [Definition 3.3.3.1](#) is an example of a pushout of sets.

002D 2. *Intersections via Unions.* Let $A, B \subset X$. We have a bijection of sets

$$A \cup B \cong A \amalg_{A \cap B} B,$$

$$\begin{array}{ccc} A \cup B & \longleftarrow & B \\ \uparrow \Gamma & & \uparrow \\ A & \longleftarrow & A \cap B \end{array}$$

PROOF 2.2.4.5 ► PROOF OF EXAMPLE 2.2.4.4

Item 1: Wedge Sums of Pointed Sets

Follows by definition.

Item 2: Intersections via Unions

Indeed, $A \coprod_{A \cap B} B$ is the quotient of $A \coprod B$ by the equivalence relation obtained by declaring $(0, a) \sim (1, b)$ iff $a = b \in A \cap B$, which is in bijection with $A \cup B$ via the map with $[(0, a)] \mapsto a$ and $[(1, b)] \mapsto b$. ▢


002E PROPOSITION 2.2.4.6 ► PROPERTIES OF PUSHOUTS OF SETS

Let A, B, C , and X be sets.


- 002F 1. *Functoriality.* The assignment $(A, B, C, f, g) \mapsto A \coprod_{f, C, g} B$ defines a functor

$$-_1 \coprod_{-3} -_1 : \text{Fun}(\mathcal{P}, \text{Sets}) \rightarrow \text{Sets},$$

where \mathcal{P} is the category that looks like this:




In particular, the action on morphisms of $-_1 \coprod_{-3} -_1$ is given by sending a morphism



in $\text{Fun}(\mathcal{P}, \text{Sets})$ to the map $\xi : A \coprod_C B \xrightarrow{\exists!} A' \coprod_{C'} B'$ given by

$$\xi(x) \stackrel{\text{def}}{=} \begin{cases} \phi(a) & \text{if } x = [(0, a)], \\ \psi(b) & \text{if } x = [(1, b)] \end{cases}$$


for each $x \in A \amalg_C B$, which is the unique map making the diagram



commute.

002G


2. *Associativity.* Given a diagram



in Sets, we have isomorphisms of sets


$$(A \amalg_X B) \amalg_Y C \cong (A \amalg_X B) \amalg_B (B \amalg_Y C) \cong A \amalg_X (B \amalg_Y C),$$

where these pullbacks are built as in the diagrams



002H

3. *Unitality.* We have isomorphisms of sets



002J

4. *Commutativity.* We have an isomorphism of sets

$$\begin{array}{ccc}
 A \amalg_X B & \longleftarrow & B \\
 \uparrow \ulcorner & & \uparrow g \\
 A & \xleftarrow{f} & X,
 \end{array}
 \quad A \amalg_X B \cong B \amalg_X A
 \quad
 \begin{array}{ccc}
 B \amalg_X A & \longleftarrow & A \\
 \uparrow \ulcorner & & \uparrow f \\
 B & \xleftarrow{g} & X.
 \end{array}$$

002K

5. *Interaction With Coproducts.* We have

$$A \amalg_{\emptyset} B \cong A \amalg B,
 \quad
 \begin{array}{ccc}
 A \amalg B & \longleftarrow & B \\
 \uparrow \ulcorner & & \uparrow i_B \\
 A & \xleftarrow{i_A} & \emptyset.
 \end{array}$$

002L

6. *Symmetric Monoidality.* The triple $(\text{Sets}, \amalg_X, X)$ is a symmetric monoidal category.

PROOF 2.2.4.7 ► PROOF OF PROPOSITION 2.2.4.6

Item 1: Functoriality

This is a special case of functoriality of co/limits, ?? of ??, with the explicit expression for ξ following from the commutativity of the cube pushout diagram.

Item 2: Associativity

Omitted.

Item 3: Unitality

Omitted.

Item 4: Commutativity

Clear.

Item 5: Interaction With Coproducts

Clear.

Item 6: Symmetric Monoidality

Omitted. 

002M 2.2.5 Coequalisers

Let A and B be sets and let $f, g: A \rightrightarrows B$ be functions.

002N

DEFINITION 2.2.5.1 ► COEQUALISERS OF SETS

The **coequaliser of f and g** is the pair $(\text{CoEq}(f, g), \text{coeq}(f, g))$ consisting of:

- *The Colimit.* The set $\text{CoEq}(f, g)$ defined by

$$\text{CoEq}(f, g) \stackrel{\text{def}}{=} B/\sim,$$

where \sim is the equivalence relation on B generated by $f(a) \sim g(a)$.

- *The Cocone.* The map

$$\text{coeq}(f, g): B \rightarrow \text{CoEq}(f, g)$$

given by the quotient map $\pi: B \rightarrow B/\sim$ with respect to the equivalence relation generated by $f(a) \sim g(a)$.

PROOF 2.2.5.2 ► PROOF OF DEFINITION 2.2.5.1

We claim that $\text{CoEq}(f, g)$ is the categorical coequaliser of f and g in Sets. First we need to check that the relevant coequaliser diagram commutes, i.e. that we have

$$\text{coeq}(f, g) \circ f = \text{coeq}(f, g) \circ g.$$

Indeed, we have

$$\begin{aligned} [\text{coeq}(f, g) \circ f](a) &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](f(a)) \\ &\stackrel{\text{def}}{=} [f(a)] \\ &= [g(a)] \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](g(a)) \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g) \circ g](a) \end{aligned}$$

for each $a \in A$. Next, we prove that $\text{CoEq}(f, g)$ satisfies the universal property of the coequaliser. Suppose we have a diagram of the form

$$\begin{array}{ccc} A & \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} & B & \xrightarrow{\text{coeq}(f, g)} & \text{CoEq}(f, g) \\ & & & \searrow c & \\ & & & & C \end{array}$$

in Sets. Then, since $c(f(a)) = c(g(a))$ for each $a \in A$, it follows from **Items 4 and 5 of Proposition 7.5.2.3** that there exists a unique map $\text{CoEq}(f, g) \xrightarrow{\exists!} C$

making the diagram

$$\begin{array}{ccccc}
 A & \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} & B & \xrightarrow{\text{coeq}(f,g)} & \text{CoEq}(f,g) \\
 & & & \searrow c & \downarrow \exists! \\
 & & & & C
 \end{array}$$

commute. □

002P

REMARK 2.2.5.3 ▶ UNWINDING DEFINITION 2.2.5.1

In detail, by [Construction 7.4.2.2](#), the relation \sim of [Definition 2.2.5.1](#) is given by declaring $a \sim b$ iff one of the following conditions is satisfied:

- We have $a = b$;
- There exist $x_1, \dots, x_n \in B$ such that $a \sim' x_1 \sim' \dots \sim' x_n \sim' b$, where we declare $x \sim' y$ if one of the following conditions is satisfied:
 1. There exists $z \in A$ such that $x = f(z)$ and $y = g(z)$.
 2. There exists $z \in A$ such that $x = g(z)$ and $y = f(z)$.

That is: we require the following condition to be satisfied:

- (★) There exist $x_1, \dots, x_n \in B$ satisfying the following conditions:
 1. There exists $z_0 \in A$ satisfying one of the following conditions:
 - (a) We have $a = f(z_0)$ and $x_1 = g(z_0)$.
 - (b) We have $a = g(z_0)$ and $x_1 = f(z_0)$.
 2. For each $1 \leq i \leq n - 1$, there exists $z_i \in A$ satisfying one of the following conditions:
 - (a) We have $x_i = f(z_i)$ and $x_{i+1} = g(z_i)$.
 - (b) We have $x_i = g(z_i)$ and $x_{i+1} = f(z_i)$.
 3. There exists $z_n \in A$ satisfying one of the following conditions:
 - (a) We have $x_n = f(z_n)$ and $b = g(z_n)$.
 - (b) We have $x_n = g(z_n)$ and $b = f(z_n)$.

002Q **EXAMPLE 2.2.5.4 ► EXAMPLES OF COEQUALISERS OF SETS**

Here are some examples of coequalisers of sets.

- 002R 1. *Quotients by Equivalence Relations.* Let R be an equivalence relation on a set X . We have a bijection of sets

$$X/\sim_R \cong \text{CoEq}\left(R \hookrightarrow X \times X \begin{matrix} \xrightarrow{\text{pr}_1} \\ \xrightarrow{\text{pr}_2} \end{matrix} X\right).$$

PROOF 2.2.5.5 ► PROOF OF EXAMPLE 2.2.5.4

Item 1: Quotients by Equivalence Relations

See [Pro24ad]. 

002S **PROPOSITION 2.2.5.6 ► PROPERTIES OF COEQUALISERS OF SETS**

Let A, B , and C be sets.

- 002T 1. *Associativity.* We have isomorphisms of sets¹

$$\underbrace{\text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h)}_{=\text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ h)} \cong \text{CoEq}(f, g, h) \cong \underbrace{\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ g)}_{=\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ h)}$$

where $\text{CoEq}(f, g, h)$ is the colimit of the diagram

$$A \begin{matrix} \xrightarrow{f} \\ \xrightarrow{g} \\ \xrightarrow{h} \end{matrix} B$$

in Sets.

- 002U 2. *Unitality.* We have an isomorphism of sets

$$\text{CoEq}(f, f) \cong B.$$

- 002V 3. *Commutativity.* We have an isomorphism of sets

$$\text{CoEq}(f, g) \cong \text{CoEq}(g, f).$$

- 002W 4. *Interaction With Composition.* Let

$$A \begin{matrix} \xrightarrow{f} \\ \xrightarrow{g} \end{matrix} B \begin{matrix} \xrightarrow{h} \\ \xrightarrow{k} \end{matrix} C$$

be functions. We have a surjection

$$\text{CoEq}(h \circ f, k \circ g) \twoheadrightarrow \text{CoEq}(\text{coeq}(h, k) \circ h \circ f, \text{coeq}(h, k) \circ k \circ g)$$

exhibiting $\text{CoEq}(\text{coeq}(h, k) \circ h \circ f, \text{coeq}(h, k) \circ k \circ g)$ as a quotient of $\text{CoEq}(h \circ f, k \circ g)$ by the relation generated by declaring $h(y) \sim k(y)$ for each $y \in B$.

¹That is, the following three ways of forming “the” coequaliser of (f, g, h) agree:

- (a) Take the coequaliser of (f, g, h) , i.e. the colimit of the diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \\ \xrightarrow{h} \end{array} B$$

in Sets.

- (b) First take the coequaliser of f and g , forming a diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B \xrightarrow{\text{coeq}(f, g)} \text{CoEq}(f, g)$$

and then take the coequaliser of the composition

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \\ \xrightarrow{h} \end{array} B \xrightarrow{\text{coeq}(f, g)} \text{CoEq}(f, g),$$

obtaining a quotient

$$\text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h) = \text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ h)$$

of $\text{CoEq}(f, g)$

- (c) First take the coequaliser of g and h , forming a diagram

$$A \begin{array}{c} \xrightarrow{g} \\ \xrightarrow{h} \end{array} B \xrightarrow{\text{coeq}(g, h)} \text{CoEq}(g, h)$$

and then take the coequaliser of the composition

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B \xrightarrow{\text{coeq}(g, h)} \text{CoEq}(g, h),$$

obtaining a quotient

$$\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ g) = \text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ h)$$


of $\text{CoEq}(g, h)$.

PROOF 2.2.5.7 ► PROOF OF PROPOSITION 2.2.5.6

Item 1: Associativity
Omitted.

Item 2: Unitality
Clear.

Item 3: Commutativity
Clear.

Item 4: Interaction With Composition
Omitted. 

002X 2.3 Operations With Sets

002Y 2.3.1 The Empty Set

002Z DEFINITION 2.3.1.1 ► THE EMPTY SET

The **empty set** is the set \emptyset defined by

$$\emptyset \stackrel{\text{def}}{=} \{x \in X \mid x \neq x\},$$

where A is the set in the set existence axiom, ?? of ??.

0030 2.3.2 Singleton Sets

Let X be a set.

0031 DEFINITION 2.3.2.1 ► SINGLETON SETS

The **singleton set containing X** is the set $\{X\}$ defined by

$$\{X\} \stackrel{\text{def}}{=} \{X, X\},$$

where $\{X, X\}$ is the pairing of X with itself ([Definition 2.3.3.1](#)).

0032 2.3.3 Pairings of Sets

Let X and Y be sets.

0033 DEFINITION 2.3.3.1 ► PAIRINGS OF SETS

The **pairing of X and Y** is the set $\{X, Y\}$ defined by

$$\{X, Y\} \stackrel{\text{def}}{=} \{x \in A \mid x = X \text{ or } x = Y\},$$

where A is the set in the axiom of pairing, ?? of ??.

0034 2.3.4 Ordered Pairs

Let A and B be sets.

0035 DEFINITION 2.3.4.1 ► ORDERED PAIRS

The **ordered pair associated to A and B** is the set (A, B) defined by

$$(A, B) \stackrel{\text{def}}{=} \{\{A\}, \{A, B\}\}.$$

0036 PROPOSITION 2.3.4.2 ► PROPERTIES OF ORDERED PAIRS

Let A and B be sets.

- 0037 1. *Uniqueness.* Let $A, B, C,$ and D be sets. The following conditions are equivalent:
- 0038 (a) We have $(A, B) = (C, D)$.
- 0039 (b) We have $A = C$ and $B = D$.

PROOF 2.3.4.3 ► PROOF OF PROPOSITION 2.3.4.2

Item 1: Uniqueness

See [Cie97, Theorem 1.2.3]. 

003A 2.3.5 Sets of Maps

Let A and B be sets.

003B DEFINITION 2.3.5.1 ► SETS OF MAPS

The **set of maps from A to B** ¹ is the set $\text{Hom}_{\text{Sets}}(A, B)$ ² whose elements are the functions from A to B .

¹Further Terminology: Also called the **Hom set from A to B** .

²Further Notation: Also written $\text{Sets}(A, B)$.

003C PROPOSITION 2.3.5.2 ► PROPERTIES OF SETS OF MAPS


Let A and B be sets.

003D 1. *Functoriality.* The assignments $X, Y, (X, Y) \mapsto \text{Hom}_{\text{Sets}}(X, Y)$ define functors

$$\begin{aligned}\text{Hom}_{\text{Sets}}(X, -) &: \text{Sets} \rightarrow \text{Sets}, \\ \text{Hom}_{\text{Sets}}(-, Y) &: \text{Sets}^{\text{op}} \rightarrow \text{Sets}, \\ \text{Hom}_{\text{Sets}}(-, -) &: \text{Sets}^{\text{op}} \times \text{Sets} \rightarrow \text{Sets}.\end{aligned}$$

PROOF 2.3.5.3 ► PROOF OF PROPOSITION 2.3.5.2

Item 1: Functoriality

This follows from **Items 2 and 5** of **Proposition 8.1.6.2**. 

003E 2.3.6 Unions of Families

Let $\{A_i\}_{i \in I}$ be a family of sets.

003F DEFINITION 2.3.6.1 ► UNIONS OF FAMILIES

The **union of the family** $\{A_i\}_{i \in I}$ is the set $\bigcup_{i \in I} A_i$ defined by

$$\bigcup_{i \in I} A_i \stackrel{\text{def}}{=} \{x \in F \mid \text{there exists some } i \in I \text{ such that } x \in A_i\},$$

where F is the set in the axiom of union, ?? of ??.

003G 2.3.7 Binary Unions

Let A and B be sets.

003H DEFINITION 2.3.7.1 ► BINARY UNIONS

The **union¹ of A and B** is the set $A \cup B$ defined by

$$A \cup B \stackrel{\text{def}}{=} \bigcup_{z \in \{A, B\}} z.$$

¹*Further Terminology:* Also called the **binary union of A and B** , for emphasis.

003J **PROPOSITION 2.3.7.2 ► PROPERTIES OF BINARY UNIONS**

Let X be a set.

003K 1. *Functoriality.* The assignments $U, V, (U, V) \mapsto U \cup V$ define functors

$$\begin{aligned} U \cup -: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ - \cup V: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ -_1 \cup -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) &\rightarrow (\mathcal{P}(X), \subset), \end{aligned}$$

where $-_1 \cup -_2$ is the functor where

· *Action on Objects.* For each $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(X)$, we have

$$[-_1 \cup -_2](U, V) \stackrel{\text{def}}{=} U \cup V.$$

· *Action on Morphisms.* For each pair of morphisms

$$\begin{aligned} \iota_U: U &\hookrightarrow U', \\ \iota_V: V &\hookrightarrow V' \end{aligned}$$

of $\mathcal{P}(X) \times \mathcal{P}(X)$, the image

$$\iota_U \cup \iota_V: U \cup V \hookrightarrow U' \cup V'$$

of (ι_U, ι_V) by \cup is the inclusion

$$U \cup V \subset U' \cup V'$$

i.e. where we have

(★) If $U \subset U'$ and $V \subset V'$, then $U \cup V \subset U' \cup V'$.

and where $U \cup -$ and $- \cup V$ are the partial functors of $-_1 \cup -_2$ at $U, V \in \mathcal{P}(X)$.

003L 2. *Via Intersections and Symmetric Differences.* We have an equality of sets

$$U \cup V = (U \Delta V) \Delta (U \cap V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

003M

3. *Associativity.* We have an equality of sets

$$(U \cup V) \cup W = U \cup (V \cup W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

003N

4. *Unitality.* We have equalities of sets

$$U \cup \emptyset = U,$$

$$\emptyset \cup U = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

003P

5. *Commutativity.* We have an equality of sets

$$U \cup V = V \cup U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

003Q

6. *Idempotency.* We have an equality of sets

$$U \cup U = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

003R

7. *Distributivity Over Intersections.* We have equalities of sets

$$U \cup (V \cap W) = (U \cup V) \cap (U \cup W),$$

$$(U \cap V) \cup W = (U \cup W) \cap (V \cup W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

003S

8. *Interaction With Characteristic Functions I.* We have

$$\chi_{U \cup V} = \max(\chi_U, \chi_V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

003T

9. *Interaction With Characteristic Functions II.* We have

$$\chi_{U \cup V} = \chi_U + \chi_V - \chi_{U \cap V}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

003U

10. *Interaction With Powersets and Semirings.* The quintuple $(\mathcal{P}(X), \cup, \cap, \emptyset, X)$ is an idempotent commutative semiring.

PROOF 2.3.7.3 ► PROOF OF PROPOSITION 2.3.7.2

Item 1: Functoriality

See [Pro24ar].

Item 2: Via Intersections and Symmetric Differences

See [Pro24bc].

Item 3: Associativity

See [Pro24be].

Item 4: Unitality

This follows from [Pro24bh] and Item 5.

Item 5: Commutativity

See [Pro24bf].

Item 6: Idempotency

See [Pro24aq].

Item 7: Distributivity Over Intersections

See [Pro24bd].


Item 8: Interaction With Characteristic Functions I

See [Pro24k].

Item 9: Interaction With Characteristic Functions II

See [Pro24k].

Item 10: Interaction With Powersets and Semirings

This follows from Items 3 to 6 and Items 3 to 5, 7 and 8 of Proposition 2.3.9.2. 

003V 2.3.8 Intersections of Families

Let \mathcal{F} be a family of sets.

003W DEFINITION 2.3.8.1 ► INTERSECTIONS OF FAMILIES

The **intersection of a family \mathcal{F} of sets** is the set $\bigcap_{X \in \mathcal{F}} X$ defined by

$$\bigcap_{X \in \mathcal{F}} X \stackrel{\text{def}}{=} \left\{ z \in \bigcup_{X \in \mathcal{F}} X \mid \text{for each } X \in \mathcal{F}, \text{ we have } z \in X \right\}.$$

003X 2.3.9 Binary Intersections

Let X and Y be sets.

003Y DEFINITION 2.3.9.1 ► BINARY INTERSECTIONS

The **intersection**¹ of X and Y is the set $X \cap Y$ defined by

$$X \cap Y \stackrel{\text{def}}{=} \bigcap_{z \in \{X, Y\}} z.$$

¹Further Terminology: Also called the **binary intersection of X and Y** , for emphasis.

003Z PROPOSITION 2.3.9.2 ► PROPERTIES OF BINARY INTERSECTIONS

Let X be a set.

0040 1. *Functoriality.* The assignments $U, V, (U, V) \mapsto U \cap V$ define functors

$$\begin{aligned} U \cap -: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ - \cap V: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ -_1 \cap -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) &\rightarrow (\mathcal{P}(X), \subset), \end{aligned}$$

where $-_1 \cap -_2$ is the functor where

- *Action on Objects.* For each $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(X)$, we have

$$[-_1 \cap -_2](U, V) \stackrel{\text{def}}{=} U \cap V.$$

- *Action on Morphisms.* For each pair of morphisms

$$\begin{aligned} \iota_U: U &\hookrightarrow U', \\ \iota_V: V &\hookrightarrow V' \end{aligned}$$

of $\mathcal{P}(X) \times \mathcal{P}(X)$, the image

$$\iota_U \cap \iota_V: U \cap V \hookrightarrow U' \cap V'$$

of (ι_U, ι_V) by \cap is the inclusion

$$U \cap V \subset U' \cap V'$$

i.e. where we have

$$(\star) \text{ If } U \subset U' \text{ and } V \subset V', \text{ then } U \cap V \subset U' \cap V'.$$

and where $U \cap -$ and $- \cap V$ are the partial functors of $-_1 \cap -_2$ at $U, V \in \mathcal{P}(X)$.

0041

2. *Adjointness.* We have adjunctions

$$(U \cap - \dashv \mathbf{Hom}_{\mathcal{P}(X)}(U, -)) : \mathcal{P}(X) \begin{array}{c} \xrightarrow{U \cap -} \\ \perp \\ \xleftarrow{\mathbf{Hom}_{\mathcal{P}(X)}(U, -)} \end{array} \mathcal{P}(X),$$

$$(- \cap V \dashv \mathbf{Hom}_{\mathcal{P}(X)}(V, -)) : \mathcal{P}(X) \begin{array}{c} \xrightarrow{- \cap V} \\ \perp \\ \xleftarrow{\mathbf{Hom}_{\mathcal{P}(X)}(V, -)} \end{array} \mathcal{P}(X),$$

where

$$\mathbf{Hom}_{\mathcal{P}(X)}(-_1, -_2) : \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$$

is the bifunctor defined by¹

$$\mathbf{Hom}_{\mathcal{P}(X)}(U, V) \stackrel{\text{def}}{=} (X \setminus U) \cup V$$

witnessed by bijections

$$\mathbf{Hom}_{\mathcal{P}(X)}(U \cap V, W) \cong \mathbf{Hom}_{\mathcal{P}(X)}(U, \mathbf{Hom}_{\mathcal{P}(X)}(V, W)),$$

$$\mathbf{Hom}_{\mathcal{P}(X)}(U \cap V, W) \cong \mathbf{Hom}_{\mathcal{P}(X)}(V, \mathbf{Hom}_{\mathcal{P}(X)}(U, W)),$$

natural in $U, V, W \in \mathcal{P}(X)$, i.e. where:

(a) The following conditions are equivalent:

- i. We have $U \cap V \subset W$.
- ii. We have $U \subset \mathbf{Hom}_{\mathcal{P}(X)}(V, W)$.
- iii. We have $U \subset (X \setminus V) \cup W$.

(b) The following conditions are equivalent:

- i. We have $V \cap U \subset W$.
- ii. We have $V \subset \mathbf{Hom}_{\mathcal{P}(X)}(U, W)$.
- iii. We have $V \subset (X \setminus U) \cup W$.

0042

3. *Associativity.* We have an equality of sets

$$(U \cap V) \cap W = U \cap (V \cap W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0043

4. *Unitality.* Let X be a set and let $U \in \mathcal{P}(X)$. We have equalities of sets

$$X \cap U = U,$$

$$U \cap X = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

0044 5. *Commutativity.* We have an equality of sets

$$U \cap V = V \cap U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

0045 6. *Idempotency.* We have an equality of sets

$$U \cap U = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

0046 7. *Distributivity Over Unions.* We have equalities of sets

$$\begin{aligned} U \cap (V \cup W) &= (U \cap V) \cup (U \cap W), \\ (U \cup V) \cap W &= (U \cap W) \cup (V \cap W) \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0047 8. *Annihilation With the Empty Set.* We have an equality of sets

$$\begin{aligned} \emptyset \cap X &= \emptyset, \\ X \cap \emptyset &= \emptyset \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

0048 9. *Interaction With Characteristic Functions I.* We have

$$\chi_{U \cap V} = \chi_U \chi_V$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

0049 10. *Interaction With Characteristic Functions II.* We have

$$\chi_{U \cap V} = \min(\chi_U, \chi_V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

004A 11. *Interaction With Powersets and Monoids With Zero.* The quadruple $((\mathcal{P}(X), \emptyset), \cap, X)$ is a commutative monoid with zero.

004B 12. *Interaction With Powersets and Semirings.* The quintuple $(\mathcal{P}(X), \cup, \cap, \emptyset, X)$ is an idempotent commutative semiring.

¹For intuition regarding the expression defining $\text{Hom}_{\mathcal{P}(X)}(U, V)$, see Remark 2.3.9.4.

PROOF 2.3.9.3 ► PROOF OF PROPOSITION 2.3.9.2

Item 1: Functoriality

See [Pro24ap].

Item 2: Adjointness

See [MSE 267469].

Item 3: Associativity

See [Pro24v].

Item 4: Unitality

This follows from [Pro24z] and Item 5.

Item 5: Commutativity

See [Pro24w].

Item 6: Idempotency

See [Pro24ao].

Item 7: Distributivity Over Unions

See [Pro24an].

Item 8: Annihilation With the Empty Set

This follows from [Pro24x] and Item 5.

Item 9: Interaction With Characteristic Functions I

See [Pro24h].

Item 10: Interaction With Characteristic Functions II

See [Pro24h].

Item 11: Interaction With Powersets and Monoids With Zero

This follows from Items 3 to 5 and 8.

Item 12: Interaction With Powersets and Semirings

This follows from Items 3 to 6 and Items 3 to 5, 7 and 8 of Proposition 2.3.9.2. 

004C

REMARK 2.3.9.4 ► INTUITION FOR THE INTERNAL HOM OF $\mathcal{P}(X)$

Since intersections are the products in $\mathcal{P}(X)$ (Item 1 of Proposition 2.4.3.3), the left adjoint $\mathbf{Hom}_{\mathcal{P}(X)}(U, V)$ may be thought of as a function type $[U, V]$. Then, under the Curry–Howard correspondence, the function type $[U, V]$ corresponds to implication $U \implies V$, which is logically equivalent to the statement $\neg U \vee V$. This in turn corresponds to the set $U^c \vee V = (X \setminus U) \cup V$.

004D **2.3.10 Differences**

Let X and Y be sets.

004E **DEFINITION 2.3.10.1 ► DIFFERENCES**

The **difference of X and Y** is the set $X \setminus Y$ defined by

$$X \setminus Y \stackrel{\text{def}}{=} \{a \in X \mid a \notin Y\}.$$

004F **PROPOSITION 2.3.10.2 ► PROPERTIES OF DIFFERENCES**

Let X be a set.

004G 1. *Functoriality.* The assignments $U, V, (U, V) \mapsto U \cap V$ define functors

$$U \setminus -: (\mathcal{P}(X), \supset) \rightarrow (\mathcal{P}(X), \subset),$$

$$- \setminus V: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \supset),$$

$$-_1 \setminus -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \supset) \rightarrow (\mathcal{P}(X), \subset),$$

where $-_1 \setminus -_2$ is the functor where

- *Action on Objects.* For each $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(X)$, we have

$$[-_1 \setminus -_2](U, V) \stackrel{\text{def}}{=} U \setminus V.$$

- *Action on Morphisms.* For each pair of morphisms

$$\iota_A: A \hookrightarrow B,$$

$$\iota_U: U \hookrightarrow V$$

of $\mathcal{P}(X) \times \mathcal{P}(X)$, the image

$$\iota_U \setminus \iota_V: A \setminus V \hookrightarrow B \setminus U$$

of (ι_U, ι_V) by \setminus is the inclusion

$$A \setminus V \subset B \setminus U$$

i.e. where we have

$$(\star) \text{ If } A \subset B \text{ and } U \subset V, \text{ then } A \setminus V \subset B \setminus U.$$

and where $U \setminus -$ and $- \setminus V$ are the partial functors of $-_1 \setminus -_2$ at $U, V \in \mathcal{P}(X)$.

004H

2. *De Morgan's Laws.* We have equalities of sets

$$X \setminus (U \cup V) = (X \setminus U) \cap (X \setminus V),$$

$$X \setminus (U \cap V) = (X \setminus U) \cup (X \setminus V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

004J

3. *Interaction With Unions I.* We have equalities of sets

$$U \setminus (V \cup W) = (U \setminus V) \cap (U \setminus W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

004K

4. *Interaction With Unions II.* We have equalities of sets

$$(U \setminus V) \cup W = (U \cup W) \setminus (V \setminus W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

004L

5. *Interaction With Unions III.* We have equalities of sets

$$U \setminus (V \cup W) = (U \cup W) \setminus (V \cup W)$$

$$= (U \setminus V) \setminus W$$

$$= (U \setminus W) \setminus V$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

004M

6. *Interaction With Unions IV.* We have equalities of sets

$$(U \cup V) \setminus W = (U \setminus W) \cup (V \setminus W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

004N

7. *Interaction With Intersections.* We have equalities of sets

$$(U \setminus V) \cap W = (U \cap W) \setminus V$$

$$= U \cap (W \setminus V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

004P

8. *Interaction With Complements.* We have an equality of sets

$$U \setminus V = U \cap V^c$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

004Q 9. *Interaction With Symmetric Differences.* We have an equality of sets

$$U \setminus V = U \Delta (U \cap V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

004R 10. *Triple Differences.* We have

$$U \setminus (V \setminus W) = (U \cap W) \cup (U \setminus V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

004S 11. *Left Annihilation.* We have

$$\emptyset \setminus U = \emptyset$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

004T 12. *Right Unitality.* We have

$$U \setminus \emptyset = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

004U 13. *Invertibility.* We have

$$U \setminus U = \emptyset$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

004V 14. *Interaction With Containment.* The following conditions are equivalent:

004W (a) We have $V \setminus U \subset W$.

004X (b) We have $V \setminus W \subset U$.

004Y 15. *Interaction With Characteristic Functions.* We have

$$\chi_{U \setminus V} = \chi_U - \chi_{U \cap V}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

PROOF 2.3.10.3 ► PROOF OF PROPOSITION 2.3.10.2

Item 1: Functoriality

See [Pro24ah] and [Pro24al].

Item 2: De Morgan's Laws

See [Pro24p].

Item 3: Interaction With Unions I

See [Pro24q].

Item 4: Interaction With Unions II

Omitted.

Item 5: Interaction With Unions III

See [Pro24am].

Item 6: Interaction With Unions IV

See [Pro24ag].

Item 7: Interaction With Intersections

See [Pro24y].

Item 8: Interaction With Complements

See [Pro24ae].

Item 9: Interaction With Symmetric Differences

See [Pro24af].

Item 10: Triple Differences

See [Pro24ak].

Item 11: Left Annihilation

Clear.

Item 12: Right Unitality

See [Pro24ai].

Item 13: Invertibility

See [Pro24aj].

Item 14: Interaction With Containment

Omitted.

Item 15: Interaction With Characteristic Functions

See [Pro24i].

**004Z 2.3.11 Complements**

Let X be a set and let $U \in \mathcal{P}(X)$.

0050

The **complement** of U is the set U^c defined by

$$\begin{aligned} U^c &\stackrel{\text{def}}{=} X \setminus U \\ &\stackrel{\text{def}}{=} \{a \in X \mid a \notin U\}. \end{aligned}$$

0051 **PROPOSITION 2.3.11.2 ► PROPERTIES OF COMPLEMENTS**

Let X be a set.

0052 1. *Functoriality.* The assignment $U \mapsto U^c$ defines a functor

$$(-)^c: \mathcal{P}(X)^{\text{op}} \rightarrow \mathcal{P}(X),$$

where

· *Action on Objects.* For each $U \in \mathcal{P}(X)$, we have

$$[(-)^c](U) \stackrel{\text{def}}{=} U^c.$$

· *Action on Morphisms.* For each morphism $\iota_U: U \hookrightarrow V$ of $\mathcal{P}(X)$, the image

$$\iota_U^c: V^c \hookrightarrow U^c$$

of ι_U by $(-)^c$ is the inclusion

$$V^c \subset U^c$$

i.e. where we have

(★) If $U \subset V$, then $V^c \subset U^c$.

0053 2. *De Morgan's Laws.* We have equalities of sets

$$(U \cup V)^c = U^c \cap V^c,$$

$$(U \cap V)^c = U^c \cup V^c$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

0054 3. *Involutority.* We have

$$(U^c)^c = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

0055 4. *Interaction With Characteristic Functions.* We have

$$\chi_{U^c} = 1 - \chi_U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.


PROOF 2.3.11.3 ► PROOF OF PROPOSITION 2.3.11.2

Item 1: Functoriality
This follows from **Item 1** of **Proposition 2.3.10.2**.

Item 2: De Morgan's Laws
See [**Pro24p**].

Item 3: Involutority
See [**Pro24l**].

Item 4: Interaction With Characteristic Functions

Clear. 

0056 2.3.12 Symmetric Differences

Let A and B be sets.

0057 DEFINITION 2.3.12.1 ► SYMMETRIC DIFFERENCES

The **symmetric difference of A and B** is the set $A \Delta B$ defined by

$$A \Delta B \stackrel{\text{def}}{=} (A \setminus B) \cup (B \setminus A).$$

0058 PROPOSITION 2.3.12.2 ► PROPERTIES OF SYMMETRIC DIFFERENCES

Let X be a set.

0059 1. *Lack of Functoriality.* The assignment $(U, V) \mapsto U \Delta V$ **need not** define functors

$$U \Delta -: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset),$$

$$- \Delta V: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset),$$

$$-_1 \Delta -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) \rightarrow (\mathcal{P}(X), \subset).$$

005A 2. *Via Unions and Intersections.* We have¹

$$U \Delta V = (U \cup V) \setminus (U \cap V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

005B 3. *Associativity.* We have²

$$(U \Delta V) \Delta W = U \Delta (V \Delta W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

005C

4. *Commutativity.* We have

$$U \Delta V = V \Delta U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

005D

5. *Unitality.* We have

$$\begin{aligned} U \Delta \emptyset &= U, \\ \emptyset \Delta U &= U \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

005E

6. *Invertibility.* We have

$$U \Delta U = \emptyset$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

005F

7. *Interaction With Unions.* We have

$$(U \Delta V) \cup (V \Delta T) = (U \cup V \cup W) \setminus (U \cap V \cap W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

005G

8. *Interaction With Complements I.* We have

$$U \Delta U^c = X$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

005H

9. *Interaction With Complements II.* We have

$$\begin{aligned} U \Delta X &= U^c, \\ X \Delta U &= U^c \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

005J

10. *Interaction With Complements III.* We have

$$U^c \Delta V^c = U \Delta V$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

005K

11. "Transitivity". We have

$$(U \Delta V) \Delta (V \Delta W) = U \Delta W$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

005L

12. *The Triangle Inequality for Symmetric Differences.* We have

$$U \Delta W \subset U \Delta V \cup V \Delta W$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

005M

13. *Distributivity Over Intersections.* We have

$$\begin{aligned} U \cap (V \Delta W) &= (U \cap V) \Delta (U \cap W), \\ (U \Delta V) \cap W &= (U \cap W) \Delta (V \cap W) \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

005N

14. *Interaction With Characteristic Functions.* We have

$$\chi_{U \Delta V} = \chi_U + \chi_V - 2\chi_{U \cap V}$$

and thus, in particular, we have

$$\chi_{U \Delta V} \equiv \chi_U + \chi_V \pmod{2}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

005P

15. *Bijection.* Given $A, B \subset \mathcal{P}(X)$, the maps

$$\begin{aligned} A \Delta -: \mathcal{P}(X) &\rightarrow \mathcal{P}(X), \\ - \Delta B: \mathcal{P}(X) &\rightarrow \mathcal{P}(X) \end{aligned}$$

are bijections with inverses given by

$$\begin{aligned} (A \Delta -)^{-1} &= - \cup (A \cap -), \\ (- \Delta B)^{-1} &= - \cup (B \cap -). \end{aligned}$$

Moreover, the map

$$C \mapsto C \Delta (A \Delta B)$$

is a bijection of $\mathcal{P}(X)$ onto itself sending A to B and B to A .

005Q

16. *Interaction With Powersets and Groups.* Let X be a set.

005R

(a) The quadruple $(\mathcal{P}(X), \Delta, \emptyset, \text{id}_{\mathcal{P}(X)})$ is an abelian group.³

005S

(b) Every element of $\mathcal{P}(X)$ has order 2 with respect to Δ , and thus $\mathcal{P}(X)$ is a *Boolean group* (i.e. an abelian 2-group).

005T

17. *Interaction With Powersets and Vector Spaces I.* The pair $(\mathcal{P}(X), \alpha_{\mathcal{P}(X)})$ consisting of

- The group $\mathcal{P}(X)$ of ??;
- The map $\alpha_{\mathcal{P}(X)}: \mathbb{F}_2 \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ defined by

$$\begin{aligned} 0 \cdot U &\stackrel{\text{def}}{=} \emptyset, \\ 1 \cdot U &\stackrel{\text{def}}{=} U; \end{aligned}$$

is an \mathbb{F}_2 -vector space.

005U

18. *Interaction With Powersets and Vector Spaces II.* If X is finite, then:(a) The set of singletons sets on the elements of X forms a basis for the \mathbb{F}_2 -vector space $(\mathcal{P}(X), \alpha_{\mathcal{P}(X)})$ of [Item 17](#).

(b) We have

$$\dim(\mathcal{P}(X)) = \#X.$$

005V

19. *Interaction With Powersets and Rings.* The quintuple $(\mathcal{P}(X), \Delta, \cap, \emptyset, X)$ is a commutative ring.⁴¹Illustration:²Illustration:³Here are some examples:i. When $X = \emptyset$, we have an isomorphism of groups between $\mathcal{P}(\emptyset)$ and the trivial group:


$$(\mathcal{P}(\emptyset), \Delta, \emptyset, \text{id}_{\mathcal{P}(\emptyset)}) \cong \text{pt.}$$

ii. When $X = \text{pt}$, we have an isomorphism of groups between $\mathcal{P}(\text{pt})$ and $\mathbb{Z}/2$:

$$(\mathcal{P}(\text{pt}), \Delta, \emptyset, \text{id}_{\mathcal{P}(\text{pt})}) \cong \mathbb{Z}/2.$$

iii. When $X = \{0, 1\}$, we have an isomorphism of groups between $\mathcal{P}(\{0, 1\})$ and $\mathbb{Z}/2 \times \mathbb{Z}/2$:

$$\left(\mathcal{P}(\{0, 1\}), \Delta, \emptyset, \text{id}_{\mathcal{P}(\{0, 1\})} \right) \cong \mathbb{Z}/2 \times \mathbb{Z}/2.$$

 *Warning:* The analogous statement replacing intersections by unions (i.e. that the quintuple $(\mathcal{P}(X), \Delta, \cup, \emptyset, X)$ is a ring) is false, however. See [Pro24ba] for a proof.
END TEXTDBEND

PROOF 2.3.12.3 ► PROOF OF PROPOSITION 2.3.12.2

Item 1: Lack of Functoriality

Omitted.

Item 2: Via Unions and Intersections

See [Pro24r].

Item 3: Associativity

See [Pro24as].

Item 4: Commutativity

See [Pro24at].

Item 5: Unitality

This follows from Item 4 and [Pro24ax].

Item 6: Invertibility

See [Pro24az].

Item 7: Interaction With Unions

See [Pro24bg].

Item 8: Interaction With Complements I

See [Pro24aw].

Item 9: Interaction With Complements II

This follows from Item 4 and [Pro24bb].

Item 10: Interaction With Complements III

See [Pro24au].

Item 11: “Transitivity”

We have

$$\begin{aligned}
 (U \Delta V) \Delta (V \Delta W) &= U \Delta (V \Delta (V \Delta W)) && \text{(by Item 3)} \\
 &= U \Delta ((V \Delta V) \Delta W) && \text{(by Item 3)} \\
 &= U \Delta (\emptyset \Delta W) && \text{(by Item 6)} \\
 &= U \Delta W && \text{(by Item 5)}
 \end{aligned}$$

Item 12: The Triangle Inequality for Symmetric Differences

This follows from [Items 2](#) and [11](#).

Item 13: Distributivity Over Intersections

See [[Pro24u](#)].

Item 14: Interaction With Characteristic Functions

See [[Pro24j](#)].

Item 15: Bijectivity

Clear.

Item 16: Interaction With Powersets and Groups

[Item 16a](#) follows from¹ [Items 3](#) to [6](#), while [Item 16b](#) follows from [Item 6](#).

Item 17: Interaction With Powersets and Vector Spaces I

Clear.

Item 18: Interaction With Powersets and Vector Spaces II

Omitted.

Item 19: Interaction With Powersets and Rings

This follows from [Items 8](#) and [11](#) of [Proposition 2.3.9.2](#) and [Items 13](#) and [16](#).² 

¹Reference: [[Pro24av](#)].

²Reference: [[Pro24ay](#)].

005W 2.4 Powersets

005X 2.4.1 Characteristic Functions

Let X be a set.

005Y DEFINITION 2.4.1.1 ► CHARACTERISTIC FUNCTIONS

Let $U \subset X$ and let $x \in X$.

- 005Z 1. The **characteristic function of U** ¹ is the function²

$$\chi_U : X \rightarrow \{t, f\}$$

defined by

$$\chi_U(x) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x \in U, \\ \text{false} & \text{if } x \notin U \end{cases}$$

for each $x \in X$.

- 0060 2. The **characteristic function of x** is the function³

$$\chi_x : X \rightarrow \{t, f\}$$

defined by

$$\chi_x \stackrel{\text{def}}{=} \chi_{\{x\}},$$

i.e. by

$$\chi_x(y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each $y \in X$.

- 0061 3. The **characteristic relation on X** ⁴ is the relation⁵

$$\chi_X(-, -) : X \times X \rightarrow \{t, f\}$$

on X defined by⁶

$$\chi_X(x, y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each $x, y \in X$.

- 0062 4. The **characteristic embedding**⁷ of X into $\mathcal{P}(X)$ is the function

$$\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$$

defined by

$$\chi_{(-)}(x) \stackrel{\text{def}}{=} \chi_x$$

for each $x \in X$.

- ¹*Further Terminology:* Also called the **indicator function of U** .
²*Further Notation:* Also written $\chi_X(U, -)$ or $\chi_X(-, U)$.
³*Further Notation:* Also written χ^x , $\chi_X(x, -)$, or $\chi_X(-, x)$.
⁴*Further Terminology:* Also called the **identity relation on X** .
⁵*Further Notation:* Also written χ_{-2}^{-1} , or \sim_{id} in the context of relations.
⁶As a subset of $X \times X$, the relation χ_X corresponds to the diagonal $\Delta_X \subset X \times X$ of X .
⁷The name "characteristic embedding" comes from the fact that there is an analogue of fully faithfulness for $\chi(-)$: given a set X , we have

$$\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_y) = \chi_X(x, y),$$

for each $x, y \in X$.

REMARK 2.4.1.2 ► CHARACTERISTIC FUNCTIONS AS DECATEGORIFICATIONS OF PRESHEAVES

0063

The definitions in [Definition 2.4.1.1](#) are decategorifications of co/presheaves, representable co/presheaves, Hom profunctors, and the Yoneda embedding:¹

0064

1. A function

$$f: X \rightarrow \{\text{t}, \text{f}\}$$

is a decategorification of a presheaf

$$\mathcal{F}: C^{\text{op}} \rightarrow \text{Sets},$$

with the characteristic functions χ_U of the subsets of X being the primordial examples (and, in fact, all examples) of these.

0065

2. The characteristic function

$$\chi_x: X \rightarrow \{\text{t}, \text{f}\}$$

of an *element* x of X is a decategorification of the representable presheaf

$$h_x: C^{\text{op}} \rightarrow \text{Sets}$$

of an *object* x of a category C .

0066

3. The characteristic relation

$$\chi_X(-_1, -_2): X \times X \rightarrow \{\text{t}, \text{f}\}$$

of X is a decategorification of the Hom profunctor

$$\text{Hom}_C(-_1, -_2): C^{\text{op}} \times C \rightarrow \text{Sets}$$

of a category C .

0067

4. The characteristic embedding

$$\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$$

of X into $\mathcal{P}(X)$ is a decategorification of the Yoneda embedding

$$\mathcal{Y} : C^{\text{op}} \hookrightarrow \text{PSh}(C)$$

of a category C into $\text{PSh}(C)$.

0068

5. There is also a direct parallel between unions and colimits:

- An element of $\mathcal{P}(X)$ is a union of elements of X , viewed as one-point subsets $\{x\} \in \mathcal{P}(A)$.
- An object of $\text{PSh}(C)$ is a colimit of objects of C , viewed as representable presheaves $h_X \in \text{Obj}(\text{PSh}(C))$.

¹These statements can be made precise by using the embeddings

$$\begin{aligned} (-)_{\text{disc}} : \text{Sets} &\hookrightarrow \text{Cats}, \\ (-)_{\text{disc}} : \{\text{t}, \text{f}\}_{\text{disc}} &\hookrightarrow \text{Sets} \end{aligned}$$

of sets into categories and of classical truth values into sets.

For instance, in this approach the characteristic function

$$\chi_x : X \rightarrow \{\text{t}, \text{f}\}$$

of an element x of X , defined by

$$\chi_x(y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each $y \in X$, is recovered as the representable presheaf

$$\text{Hom}_{X_{\text{disc}}}(-, x) : X_{\text{disc}} \rightarrow \text{Sets}$$

of the corresponding object x of X_{disc} , defined on objects by

$$\text{Hom}_{X_{\text{disc}}}(y, x) \stackrel{\text{def}}{=} \begin{cases} \text{pt} & \text{if } x = y, \\ \emptyset & \text{if } x \neq y \end{cases}$$

for each $y \in \text{Obj}(X_{\text{disc}})$.

0069

PROPOSITION 2.4.1.3 ► PROPERTIES OF CHARACTERISTIC FUNCTIONS

Let X be a set.

006A

1. *The Inclusion of Characteristic Relations Associated to a Function.* Let

$f: A \rightarrow B$ be a function. We have an inclusion¹

$$\chi_B \circ (f \times f) \subset \chi_A,$$

$$\begin{array}{ccc} A \times A & \xrightarrow{f \times f} & B \times B \\ \searrow \chi_A & \supset & \swarrow \chi_B \\ & & \{t, f\}. \end{array}$$

006B 2. *Interaction With Unions I.* We have

$$\chi_{U \cup V} = \max(\chi_U, \chi_V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

006C 3. *Interaction With Unions II.* We have

$$\chi_{U \cup V} = \chi_U + \chi_V - \chi_{U \cap V}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

006D 4. *Interaction With Intersections I.* We have

$$\chi_{U \cap V} = \chi_U \chi_V$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

006E 5. *Interaction With Intersections II.* We have

$$\chi_{U \cap V} = \min(\chi_U, \chi_V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

006F 6. *Interaction With Differences.* We have

$$\chi_{U \setminus V} = \chi_U - \chi_{U \cap V}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

006G 7. *Interaction With Complements.* We have

$$\chi_{U^c} = 1 - \chi_U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

006H

8. *Interaction With Symmetric Differences.* We have

$$\chi_{U\Delta V} = \chi_U + \chi_V - 2\chi_{U\cap V}$$

and thus, in particular, we have

$$\chi_{U\Delta V} \equiv \chi_U + \chi_V \pmod{2}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

006J

9. *Interaction Between the Characteristic Embedding and Morphisms.* Let $f: X \rightarrow Y$ be a map of sets. The diagram

$$f_* \circ \chi_X = \chi_{X'} \circ f,$$

$$\begin{array}{ccc} X & \xrightarrow{f} & X' \\ \chi_X \downarrow & & \downarrow \chi_{X'} \\ \mathcal{P}(X) & \xrightarrow{f_*} & \mathcal{P}(X'). \end{array}$$

commutes.

¹This is the 0-categorical version of [Definition 8.4.4.1](#).

PROOF 2.4.1.4 ► PROOF OF PROPOSITION 2.4.1.3

Item 1: The Inclusion of Characteristic Relations Associated to a Function

The inclusion $\chi_B(f(a), f(b)) \subset \chi_A(a, b)$ is equivalent to the statement “if $a = b$, then $f(a) = f(b)$ ”, which is true.

Item 2: Interaction With Unions I

This is a repetition of [Item 8](#) of [Proposition 2.3.7.2](#) and is proved there.

Item 3: Interaction With Unions II

This is a repetition of [Item 9](#) of [Proposition 2.3.7.2](#) and is proved there.

Item 4: Interaction With Intersections I

This is a repetition of [Item 9](#) of [Proposition 2.3.9.2](#) and is proved there.

Item 5: Interaction With Intersections II

This is a repetition of [Item 10](#) of [Proposition 2.3.9.2](#) and is proved there.

Item 6: Interaction With Differences

This is a repetition of [Item 15](#) of [Proposition 2.3.10.2](#) and is proved there.

Item 7: Interaction With Complements

This is a repetition of **Item 4** of **Proposition 2.3.11.2** and is proved there.

Item 8: Interaction With Symmetric Differences

This is a repetition of **Item 14** of **Proposition 2.3.12.2** and is proved there.

Item 9: Interaction Between the Characteristic Embedding and Morphisms

Indeed, we have

$$\begin{aligned} [f_* \circ \chi_X](x) &\stackrel{\text{def}}{=} f_*(\chi_X(x)) \\ &\stackrel{\text{def}}{=} f_*({x}) \\ &= \{f(x)\} \\ &\stackrel{\text{def}}{=} \chi_{X'}(f(x)) \\ &\stackrel{\text{def}}{=} [\chi_{X'} \circ f](x), \end{aligned}$$

for each $x \in X$, showing the desired equality. 

006K 2.4.2 The Yoneda Lemma for Sets

Let X be a set and let $U \subset X$ be a subset of X .

006L PROPOSITION 2.4.2.1 ► THE YONEDA LEMMA FOR SETS

We have

$$\chi_{\mathcal{P}(X)}(\chi_x, \chi_U) = \chi_U(x)$$

for each $x \in X$, giving an equality of functions

$$\chi_{\mathcal{P}(X)}(\chi_{(-)}, \chi_U) = \chi_U.$$

PROOF 2.4.2.2 ► PROOF OF PROPOSITION 2.4.2.1

Clear. 

006M COROLLARY 2.4.2.3 ► THE CHARACTERISTIC EMBEDDING IS FULLY FAITHFUL

The characteristic embedding is fully faithful, i.e., we have

$$\chi_{\mathcal{P}(X)}(\chi_x, \chi_y) = \chi_X(x, y)$$

for each $x, y \in X$.

PROOF 2.4.2.4 ► PROOF OF COROLLARY 2.4.2.3

This follows from [Proposition 2.4.2.1](#). 

006N 2.4.3 Powersets

Let X be a set.

006P DEFINITION 2.4.3.1 ► POWERSETS

The **powerset of X** is the set $\mathcal{P}(X)$ defined by

$$\mathcal{P}(X) \stackrel{\text{def}}{=} \{U \in P \mid U \subset X\},$$

where P is the set in the axiom of powerset, ?? of ??.

006Q

REMARK 2.4.3.2 ► POWERSETS AS DECATEGORIFICATIONS OF CO/PRESHEAF CATEGORIES

The powerset of a set is a decategorification of the category of presheaves of a category: while¹

- The powerset of a set X is equivalently ([Items 1 and 2 of Proposition 2.4.3.9](#)) the set

$$\text{Sets}(X, \{t, f\})$$

of functions from X to the set $\{t, f\}$ of classical truth values.

- The category of presheaves on a category C is the category

$$\text{Fun}(C^{\text{op}}, \text{Sets})$$

of functors from C^{op} to the category Sets of sets.

¹This parallel is based on the following comparison:

- A category is enriched over the category

$$\text{Sets} \stackrel{\text{def}}{=} \text{Cats}_0$$

of sets (i.e. “0-categories”), with presheaves taking values on it.

- A set is enriched over the set

$$\{t, f\} \stackrel{\text{def}}{=} \text{Cats}_{-1}$$

of classical truth values (i.e. “(-1)-categories”), with characteristic functions taking values on it.

006R PROPOSITION 2.4.3.3 ► PROPERTIES OF POWERSETS: AS CATEGORIES

Let X be a set.

- 006S 1. *Co/Completeness.* The (posetal) category (associated to) $(\mathcal{P}(X), \subset)$ is complete and cocomplete:
- (a) *Products.* The products in $\mathcal{P}(X)$ are given by intersection of subsets.
 - (b) *Coproducts.* The coproducts in $\mathcal{P}(X)$ are given by union of subsets.
 - (c) *Co/Equalisers.* Being a posetal category, $\mathcal{P}(X)$ only has at most one morphisms between any two objects, so co/equalisers are trivial.
- 006T 2. *Cartesian Closedness.* The category $\mathcal{P}(X)$ is Cartesian closed with internal Hom

$$\mathbf{Hom}_{\mathcal{P}(X)}(-_1, -_2) : \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$$

given by¹

$$\mathbf{Hom}_{\mathcal{P}(X)}(U, V) \stackrel{\text{def}}{=} (X \setminus U) \cup V$$

for each $U, V \in \text{Obj}(\mathcal{P}(X))$.

¹For intuition regarding the expression defining $\mathbf{Hom}_{\mathcal{P}(X)}(U, V)$, see Remark 2.3.9.4.

PROOF 2.4.3.4 ► PROOF OF PROPOSITION 2.4.3.3

Item 1: Co/Completeness

Clear.

Item 2: Cartesian Closedness

This follows from Item 2 of Proposition 2.3.9.2. 

006U PROPOSITION 2.4.3.5 ► PROPERTIES OF POWERSETS: FUNCTORIALITY AND ADJOINTNESS

Let X be a set.

- 006V 1. *Functoriality I.* The assignment $X \mapsto \mathcal{P}(X)$ defines a functor

$$\mathcal{P}_* : \text{Sets} \rightarrow \text{Sets},$$

where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$\mathcal{P}_*(A) \stackrel{\text{def}}{=} \mathcal{P}(A).$$

- *Action on Morphisms.* For each $A, B \in \text{Obj}(\text{Sets})$, the action on morphisms

$$\mathcal{P}_{*|A,B}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B))$$

of \mathcal{P}_* at (A, B) is the map defined by sending a map of sets $f: A \rightarrow B$ to the map

$$\mathcal{P}_*(f): \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\mathcal{P}_*(f) \stackrel{\text{def}}{=} f_*$$

as in [Definition 2.4.4.1](#).

006W

2. *Functoriality II.* The assignment $X \mapsto \mathcal{P}(X)$ defines a functor

$$\mathcal{P}^{-1}: \text{Sets}^{\text{op}} \rightarrow \text{Sets},$$

where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$\mathcal{P}^{-1}(A) \stackrel{\text{def}}{=} \mathcal{P}(A).$$

- *Action on Morphisms.* For each $A, B \in \text{Obj}(\text{Sets})$, the action on morphisms

$$\mathcal{P}_{A,B}^{-1}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(B), \mathcal{P}(A))$$

of \mathcal{P}^{-1} at (A, B) is the map defined by sending a map of sets $f: A \rightarrow B$ to the map

$$\mathcal{P}^{-1}(f): \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by

$$\mathcal{P}^{-1}(f) \stackrel{\text{def}}{=} f^{-1},$$

as in [Definition 2.4.5.1](#).

006X

3. *Functoriality III.* The assignment $X \mapsto \mathcal{P}(X)$ defines a functor

$$\mathcal{P}_! : \mathbf{Sets} \rightarrow \mathbf{Sets},$$

where

- *Action on Objects.* For each $A \in \mathbf{Obj}(\mathbf{Sets})$, we have

$$\mathcal{P}_!(A) \stackrel{\text{def}}{=} \mathcal{P}(A).$$

- *Action on Morphisms.* For each $A, B \in \mathbf{Obj}(\mathbf{Sets})$, the action on morphisms

$$\mathcal{P}_!|_{A,B} : \mathbf{Sets}(A, B) \rightarrow \mathbf{Sets}(\mathcal{P}(A), \mathcal{P}(B))$$

of $\mathcal{P}_!$ at (A, B) is the map defined by sending a map of sets $f: A \rightarrow B$ to the map

$$\mathcal{P}_!(f) : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\mathcal{P}_!(f) \stackrel{\text{def}}{=} f_!$$

as in [Definition 2.4.6.1](#).

006Y

4. *Adjointness I.* We have an adjunction

$$(\mathcal{P}^{-1} \dashv \mathcal{P}^{-1, \text{op}}) : \mathbf{Sets}^{\text{op}} \begin{array}{c} \xrightarrow{\mathcal{P}^{-1}} \\ \perp \\ \xleftarrow{\mathcal{P}^{-1, \text{op}}} \end{array} \mathbf{Sets},$$

witnessed by a bijection

$$\underbrace{\mathbf{Sets}^{\text{op}}(\mathcal{P}(A), B)}_{\stackrel{\text{def}}{=} \mathbf{Sets}(B, \mathcal{P}(A))} \cong \mathbf{Sets}(A, \mathcal{P}(B)),$$

natural in $A \in \mathbf{Obj}(\mathbf{Sets})$ and $B \in \mathbf{Obj}(\mathbf{Sets}^{\text{op}})$.

006Z

5. *Adjointness II.* We have an adjunction

$$(\text{Gr} \dashv \mathcal{P}_*) : \mathbf{Sets} \begin{array}{c} \xrightarrow{\text{Gr}} \\ \perp \\ \xleftarrow{\mathcal{P}_*} \end{array} \mathbf{Rel},$$

witnessed by a bijection of sets

$$\mathbf{Rel}(\text{Gr}(A), B) \cong \mathbf{Sets}(A, \mathcal{P}(B))$$

natural in $A \in \mathbf{Obj}(\mathbf{Sets})$ and $B \in \mathbf{Obj}(\mathbf{Rel})$, where Gr is the graph functor of [Item 1 of Proposition 6.3.1.2](#) and \mathcal{P}_* is the functor of [Proposition 6.4.5.1](#).

PROOF 2.4.3.6 ► PROOF OF PROPOSITION 2.4.3.5

Item 1: Functoriality I

This follows from **Items 3 and 4** of **Proposition 2.4.4.6**.

Item 2: Functoriality II

This follows **Items 3 and 4** of **Proposition 2.4.5.5**.

Item 3: Functoriality III

This follows **Items 3 and 4** of **Proposition 2.4.6.8**.

Item 4: Adjointness I

We have

$$\begin{aligned}
 \text{Sets}^{\text{op}}(\mathcal{P}(A), B) &\stackrel{\text{def}}{=} \text{Sets}(B, \mathcal{P}(A)) \\
 &\cong \text{Sets}(B, \text{Sets}(A, \{t, f\})) && \text{(by Item 1 of Proposition 2.4.3.9)} \\
 &\cong \text{Sets}(A \times B, \{t, f\}) && \text{(by Item 2 of Proposition 2.1.3.3)} \\
 &\cong \text{Sets}(A, \text{Sets}(B, \{t, f\})) && \text{(by Item 2 of Proposition 2.1.3.3)} \\
 &\cong \text{Sets}(A, \mathcal{P}(B)) && \text{(by Item 1 of Proposition 2.4.3.9)}
 \end{aligned}$$

with all bijections natural in A and B (where we use **Item 2** of **Proposition 2.4.3.9** here).

Item 5: Adjointness II

We have

$$\begin{aligned}
 \text{Rel}(\text{Gr}(A), B) &\cong \mathcal{P}(A \times B) \\
 &\cong \text{Sets}(A \times B, \{t, f\}) && \text{(by Item 1 of Proposition 2.4.3.9)} \\
 &\cong \text{Sets}(A, \text{Sets}(B, \{t, f\})) && \text{(by Item 2 of Proposition 2.1.3.3)} \\
 &\cong \text{Sets}(A, \mathcal{P}(B)) && \text{(by Item 1 of Proposition 2.4.3.9)}
 \end{aligned}$$

with all bijections natural in A (where we use **Item 2** of **Proposition 2.4.3.9** here). Explicitly, this isomorphism is given by sending a relation $R: \text{Gr}(A) \rightarrow B$ to the map $R^\dagger: A \rightarrow \mathcal{P}(B)$ sending a to the subset $R(a)$ of B , as in **Remark 5.1.1.4**.

Naturality in B is then the statement that given a relation $R: B \rightarrow B'$, the diagram

$$\begin{array}{ccc} \text{Rel}(\text{Gr}(A), B) & \xrightarrow{R \circ -} & \text{Rel}(\text{Gr}(A), B') \\ \downarrow \wr & & \downarrow \wr \\ \text{Sets}(A, \mathcal{P}(B)) & \xrightarrow{R_*} & \text{Sets}(A, \mathcal{P}(B')) \end{array}$$

commutes, which follows from [Remark 6.4.1.2](#). 

0070 PROPOSITION 2.4.3.7 ► PROPERTIES OF POWERSETS: MONOIDALITY

Let X be a set.

- 0071 1. *Symmetric Strong Monoidality With Respect to Coproducts I.* The powerset functor \mathcal{P}_* of [Item 1](#) of [Proposition 2.4.3.5](#) has a symmetric strong monoidal structure

$$\left(\mathcal{P}_*, \mathcal{P}_* \amalg, \mathcal{P}_{*\mathbb{1}} \amalg \right): (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}, \amalg, \emptyset)$$

being equipped with isomorphisms

$$\begin{aligned} \mathcal{P}_{*|X,Y}^{\amalg} &: \mathcal{P}(X) \times \mathcal{P}(Y) \xrightarrow{\cong} \mathcal{P}(X \amalg Y), \\ \mathcal{P}_{*\mathbb{1}}^{\amalg} &: \text{pt} \xrightarrow{\cong} \mathcal{P}(\emptyset), \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

- 0072 2. *Symmetric Strong Monoidality With Respect to Coproducts II.* The powerset functor \mathcal{P}^{-1} of [Item 2](#) of [Proposition 2.4.3.5](#) has a symmetric strong monoidal structure

$$\left(\mathcal{P}^{-1}, \mathcal{P}^{-1} \amalg, \mathcal{P}_{\mathbb{1}}^{-1} \amalg \right): (\text{Sets}^{\text{op}}, \times^{\text{op}}, \text{pt}) \rightarrow (\text{Sets}, \amalg, \emptyset)$$

being equipped with isomorphisms

$$\begin{aligned} \mathcal{P}_{X,Y}^{-1 \amalg} &: \mathcal{P}(X) \times \mathcal{P}(Y) \xrightarrow{\cong} \mathcal{P}(X \amalg Y), \\ \mathcal{P}_{\mathbb{1}}^{-1 \amalg} &: \text{pt} \xrightarrow{\cong} \mathcal{P}(\emptyset), \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

0073

3. *Symmetric Strong Monoidality With Respect to Coproducts III.* The powerset functor $\mathcal{P}_!$ of [Item 3](#) of [Proposition 2.4.3.5](#) has a symmetric strong monoidal structure

$$\left(\mathcal{P}_!, \mathcal{P}_!^{\amalg}, \mathcal{P}_!^{\amalg}\right): (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}, \amalg, \emptyset)$$

being equipped with isomorphisms

$$\begin{aligned} \mathcal{P}_!^{\amalg} : \mathcal{P}(X) \times \mathcal{P}(Y) &\xrightarrow{\cong} \mathcal{P}(X \amalg Y), \\ \mathcal{P}_!^{\amalg} : \text{pt} &\xrightarrow{\cong} \mathcal{P}(\emptyset), \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

0074

4. *Symmetric Lax Monoidality With Respect to Products I.* The powerset functor \mathcal{P}_* of [Item 1](#) of [Proposition 2.4.3.5](#) has a symmetric lax monoidal structure

$$\left(\mathcal{P}_*, \mathcal{P}_*^{\otimes}, \mathcal{P}_*^{\otimes}\right): (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}, \times, \text{pt})$$

being equipped with morphisms

$$\begin{aligned} \mathcal{P}_*^{\otimes} : \mathcal{P}(X) \times \mathcal{P}(Y) &\rightarrow \mathcal{P}(X \times Y), \\ \mathcal{P}_*^{\otimes} : \text{pt} &\rightarrow \mathcal{P}(\text{pt}), \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$, where

- The map \mathcal{P}_*^{\otimes} is given by

$$\mathcal{P}_*^{\otimes}(U, V) \stackrel{\text{def}}{=} U \times V$$

for each $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(Y)$,

- The map \mathcal{P}_*^{\otimes} is given by

$$\mathcal{P}_*^{\otimes}(\star) = \text{pt}.$$

0075

5. *Symmetric Lax Monoidality With Respect to Products II.* The powerset functor \mathcal{P}^{-1} of [Item 2](#) of [Proposition 2.4.3.5](#) has a symmetric lax monoidal structure

$$\left(\mathcal{P}^{-1}, \mathcal{P}^{-1|\otimes}, \mathcal{P}^{-1|\otimes}\right): (\text{Sets}^{\text{op}}, \times^{\text{op}}, \text{pt}) \rightarrow (\text{Sets}, \times, \text{pt})$$

being equipped with morphisms

$$\begin{aligned}\mathcal{P}_{X,Y}^{-1|\times} : \mathcal{P}(X) \times \mathcal{P}(Y) &\rightarrow \mathcal{P}(X \times Y), \\ \mathcal{P}_{\mathbb{1}}^{\times} : \text{pt} &\rightarrow \mathcal{P}(\emptyset),\end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$, defined as in [Item 4](#).

0076

6. *Symmetric Lax Monoidality With Respect to Products III*. The powerset functor $\mathcal{P}_!$ of [Item 3](#) of [Proposition 2.4.3.5](#) has a symmetric lax monoidal structure

$$\left(\mathcal{P}_!, \mathcal{P}_!^{\otimes}, \mathcal{P}_{\mathbb{1}\mathbb{1}}^{\otimes}\right) : (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}, \times, \text{pt})$$

being equipped with morphisms

$$\begin{aligned}\mathcal{P}_{!X,Y}^{\times} : \mathcal{P}(X) \times \mathcal{P}(Y) &\rightarrow \mathcal{P}(X \times Y), \\ \mathcal{P}_{\mathbb{1}\mathbb{1}}^{\times} : \text{pt} &\rightarrow \mathcal{P}(\emptyset),\end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$, defined as in [Item 4](#).

PROOF 2.4.3.8 ► PROOF OF PROPOSITION 2.4.3.7

Item 1: Symmetric Strong Monoidality With Respect to Coproducts I

The isomorphism

$$\mathcal{P}_{*|X,Y}^{\coprod} : \mathcal{P}(X) \times \mathcal{P}(Y) \rightarrow \mathcal{P}(X \coprod Y)$$

is given by sending $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(Y)$ to $U \coprod V$, with inverse given by sending a subset S of $X \coprod Y$ to the pair $(S_X, S_Y) \in \mathcal{P}(X) \times \mathcal{P}(Y)$ with

$$\begin{aligned}S_X &\stackrel{\text{def}}{=} \{x \in X \mid (0, x) \in S\} \\ S_Y &\stackrel{\text{def}}{=} \{y \in Y \mid (1, y) \in S\}.\end{aligned}$$

The isomorphism $\text{pt} \cong \mathcal{P}(\emptyset)$ is given by $\star \mapsto \emptyset \in \mathcal{P}(\emptyset)$.

Naturality for the isomorphism $\mathcal{P}_{*|X,Y}^{\amalg}$ is the statement that, given maps of sets $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$, the diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(Y) & \xrightarrow{f_* \times g_*} & \mathcal{P}(X') \times \mathcal{P}(Y') \\ \downarrow \wr & & \downarrow \wr \\ \mathcal{P}(X \amalg Y) & \xrightarrow{(f \amalg g)_*} & \mathcal{P}(X' \amalg Y') \end{array}$$

commutes, which is clear, as it acts on elements as

$$\begin{array}{ccc} (U, V) & \longmapsto & (f_*(U), g_*(V)) \\ \downarrow & & \downarrow \\ U \amalg V & \longmapsto & (f \amalg g)_*(U \amalg V) = f_*(U) \amalg g_*(V), \end{array}$$

where we are using [Item 7](#) of [Proposition 2.4.4.4](#).

Finally, monoidality, unity, and symmetry of \mathcal{P}_* as a monoidal functor follow by checking the commutativity of the relevant diagrams on elements.

Item 2: Symmetric Strong Monoidality With Respect to Coproducts II

The proof is similar to [Item 1](#), and is hence omitted.

Item 3: Symmetric Strong Monoidality With Respect to Coproducts III

The proof is similar to [Item 1](#), and is hence omitted.

Item 4: Symmetric Lax Monoidality With Respect to Products I

Naturality for the morphism $\mathcal{P}_{*|X,Y}^{\times}$ is the statement that, given maps of sets $f: X \rightarrow X'$ and $g: Y \rightarrow Y'$, the diagram

$$\begin{array}{ccc} \mathcal{P}(X) \times \mathcal{P}(Y) & \xrightarrow{f_* \times g_*} & \mathcal{P}(X') \times \mathcal{P}(Y') \\ \downarrow \wr & & \downarrow \wr \\ \mathcal{P}(X \times Y) & \xrightarrow{(f \times g)_*} & \mathcal{P}(X' \times Y') \end{array}$$

commutes, which is clear, as it acts on elements as

$$\begin{array}{ccc} (U, V) & \longmapsto & (f_*(U), g_*(V)) \\ \downarrow & & \downarrow \\ U \times V & \longmapsto & (f \times g)_*(U \times V) = f_*(U) \times g_*(V), \end{array}$$

where we are using **Item 8** of **Proposition 2.4.4.4**.

Finally, monoidality, unity, and symmetry of \mathcal{P}_* as a monoidal functor follow by checking the commutativity of the relevant diagrams on elements.

Item 5: Symmetric Lax Monoidality With Respect to Products II

The proof is similar to **Item 4**, and is hence omitted.

Item 6: Symmetric Lax Monoidality With Respect to Products III

The proof is similar to **Item 4**, and is hence omitted. 

PROPOSITION 2.4.3.9 ► PROPERTIES OF POWERSETS: AS SETS OF FUNCTIONS/RELATIONS

0077

Let X be a set.

0078

1. *Powersets as Sets of Functions I.* The assignment $U \mapsto \chi_U$ defines a bijection

$$\chi_{(-)}: \mathcal{P}(X) \xrightarrow{\cong} \text{Sets}(X, \{\text{t}, \text{f}\}),$$

for each $X \in \text{Obj}(\text{Sets})$.

0079

2. *Powersets as Sets of Functions II.* The bijection

$$\mathcal{P}(X) \cong \text{Sets}(X, \{\text{t}, \text{f}\})$$

of **Item 1** is natural in $X \in \text{Obj}(\text{Sets})$, refining to a natural isomorphism of functors

$$\mathcal{P}^{-1} \cong \text{Sets}(-, \{\text{t}, \text{f}\}).$$

007A

3. *Powersets as Sets of Relations.* We have bijections

$$\mathcal{P}(X) \cong \text{Rel}(\text{pt}, X),$$

$$\mathcal{P}(X) \cong \text{Rel}(X, \text{pt}),$$

natural in $X \in \text{Obj}(\text{Sets})$.

PROOF 2.4.3.10 ► PROOF OF PROPOSITION 2.4.3.9

Item 1: Powersets as Sets of Functions I

Indeed, the inverse of $\chi_{(-)}$ is given by sending a function $f: X \rightarrow \{\text{t}, \text{f}\}$ to the subset U_f of $\mathcal{P}(X)$ defined by

$$U_f \stackrel{\text{def}}{=} \{x \in X \mid f(x) = \text{true}\},$$

i.e. by $U_f = f^{-1}(\text{true})$. That $\chi_{(-)}$ and $f \mapsto U_f$ are inverses is then straightforward to check.

Item 2: Powersets as Sets of Functions II

We need to check that, given a function $f: X \rightarrow Y$, the diagram

$$\begin{array}{ccc} \mathcal{P}(Y) & \xrightarrow{f^{-1}} & \mathcal{P}(X) \\ \chi_{(-)} \downarrow \wr & & \wr \downarrow \chi_{(-)} \\ \text{Sets}(Y, \{\text{t}, \text{f}\}) & \xrightarrow{f^*} & \text{Sets}(X, \{\text{t}, \text{f}\}) \end{array}$$

commutes, i.e. that for each $V \in \mathcal{P}(Y)$, we have

$$\chi_V \circ f = \chi_{f^{-1}(V)}.$$

And indeed, we have

$$\begin{aligned} [\chi_V \circ f](v) &\stackrel{\text{def}}{=} \chi_V(f(v)) \\ &= \begin{cases} \text{true} & \text{if } f(v) \in V, \\ \text{false} & \text{otherwise} \end{cases} \\ &= \begin{cases} \text{true} & \text{if } v \in f^{-1}(V), \\ \text{false} & \text{otherwise} \end{cases} \\ &\stackrel{\text{def}}{=} \chi_{f^{-1}(V)}(v) \end{aligned}$$

for each $v \in V$.

Item 3: Powersets as Sets of Relations

Indeed, we have

$$\begin{aligned} \text{Rel}(\text{pt}, X) &\stackrel{\text{def}}{=} \mathcal{P}(\text{pt} \times X) \\ &\cong \mathcal{P}(X) \end{aligned}$$

and

$$\begin{aligned} \text{Rel}(X, \text{pt}) &\stackrel{\text{def}}{=} \mathcal{P}(X \times \text{pt}) \\ &\cong \mathcal{P}(X), \end{aligned}$$

where we have used [Item 4](#) of [Proposition 2.1.3.3](#). 

007B

REMARK 2.4.3.11 ► POWERSETS AS SETS OF FUNCTIONS AND UN/STRAIGHTENING

The bijection

$$\mathcal{P}(X) \cong \text{Sets}(X, \{\text{t}, \text{f}\})$$

of Item 1 of Proposition 2.4.3.9, which

- Takes a subset $U \hookrightarrow X$ of X and *straightens* it to a function $\chi_U: X \rightarrow \{\text{true}, \text{false}\}$;
- Takes a function $f: X \rightarrow \{\text{true}, \text{false}\}$ and *unstraightens* it to a subset $f^{-1}(\text{true}) \hookrightarrow X$ of X ;

may be viewed as the (-1) -categorical version of the un/straightening isomorphism for indexed and fibred sets

$$\underbrace{\text{FibSets}(X)}_{\stackrel{\text{def}}{=} \text{Sets}_{/X}} \cong \underbrace{\text{ISets}(X)}_{\stackrel{\text{def}}{=} \text{Fun}(X_{\text{disc}}, \text{Sets})}$$

of ??, where we view:

- Subsets $U \hookrightarrow X$ as analogous to X -fibred sets $\phi_X: A \rightarrow X$.
- Functions $f: X \rightarrow \{\text{t}, \text{f}\}$ as analogous to X -indexed sets $A: X_{\text{disc}} \rightarrow \text{Sets}$.

007C

PROPOSITION 2.4.3.12 ► PROPERTIES OF POWERSETS: AS FREE COCOMPLETIONS

Let X be a set.

007D

1. *Universal Property.* The pair $(\mathcal{P}(X), \chi_{(-)})$ consisting of

- The powerset $\mathcal{P}(X)$ of X ;
- The characteristic embedding $\chi_{(-)}: X \hookrightarrow \mathcal{P}(X)$ of X into $\mathcal{P}(X)$;


satisfies the following universal property:

- (★) Given another pair (Y, f) consisting of
- A cocomplete poset (Y, \preceq) ;
 - A function $f: X \rightarrow Y$;

there exists a unique cocontinuous morphism of posets

$$(\mathcal{P}(X), \subset) \xrightarrow{\exists!} (Y, \preceq)$$

making the diagram



commute.

007E

2. *Adjointness.* We have an adjunction¹

$$(\mathcal{P} \dashv \exists): \text{Sets} \begin{array}{c} \xrightarrow{\mathcal{P}} \\ \perp \\ \xleftarrow{\exists} \end{array} \text{Pos}^{\text{cocomp.}},$$

witnessed by a bijection

$$\text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y, \preceq)) \cong \text{Sets}(X, Y),$$

natural in $X \in \text{Obj}(\text{Sets})$ and $(Y, \preceq) \in \text{Obj}(\text{Pos}^{\text{cocomp.}})$, where the maps witnessing this bijection are given by

· The map

$$\chi_X^*: \text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y, \preceq)) \rightarrow \text{Sets}(X, Y)$$

defined by

$$\chi_X^*(f) \stackrel{\text{def}}{=} f \circ \chi_X,$$

i.e. by sending a cocontinuous morphism of posets $f: \mathcal{P}(X) \rightarrow Y$ to the composition

$$X \xrightarrow{\chi_X} \mathcal{P}(X) \xrightarrow{f} Y.$$

· The map

$$\text{Lan}_{\chi_X}: \text{Sets}(X, Y) \rightarrow \text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y, \preceq))$$

is given by sending a function $f: X \rightarrow Y$ to its left Kan extension along χ_X ,

$$\text{Lan}_{\chi_X}(f): \mathcal{P}(X) \rightarrow Y,$$

Moreover, $\text{Lan}_{\chi_X}(f)$ can be explicitly computed by

$$\begin{aligned} [\text{Lan}_{\chi_X}(f)](U) &\cong \int^{x \in X} \chi_{\mathcal{P}(X)}(\chi_x, U) \odot f(x) \\ &\cong \int^{x \in X} \chi_U(x) \odot f(x) \quad (\text{by Proposition 2.4.2.1}) \\ &\cong \bigvee_{x \in X} (\chi_U(x) \odot f(x)) \end{aligned}$$

for each $U \in \mathcal{P}(X)$, where:

- \bigvee is the join in (Y, \preceq) .
- We have

$$\begin{aligned} \text{true} \odot f(x) &\stackrel{\text{def}}{=} f(x), \\ \text{false} \odot f(x) &\stackrel{\text{def}}{=} \emptyset_Y, \end{aligned}$$

where \emptyset_Y is the minimal element of (Y, \preceq) .

¹In this sense, $\mathcal{P}(A)$ is the free cocompletion of A . (Note that, despite its name, however, this is not an idempotent operation, as we have $\mathcal{P}(\mathcal{P}(A)) \neq \mathcal{P}(A)$.)

PROOF 2.4.3.13 ► PROOF OF PROPOSITION 2.4.3.7

Item 1: Universal Property

This is a rephrasing of **Item 2**.

Item 2: Adjointness

We claim we have adjunction $\mathcal{P} \dashv \overline{\text{S}}$, witnessed by a bijection

$$\text{Pos}^{\text{cocomp}}((\mathcal{P}(X), \subset), (Y, \preceq)) \cong \text{Sets}(X, Y),$$

natural in $X \in \text{Obj}(\text{Sets})$ and $(Y, \preceq) \in \text{Obj}(\text{Pos}^{\text{cocomp}})$.

- *Map I*. We define a map

$$\Phi_{X,Y}: \text{Pos}^{\text{cocomp}}((\mathcal{P}(X), \subset), (Y, \preceq)) \rightarrow \text{Sets}(X, Y)$$

as in the statement, by


$$\Phi_{X,Y}(f) \stackrel{\text{def}}{=} f \circ \chi_X$$

for each $f \in \text{Pos}^{\text{cocomp}}((\mathcal{P}(X), \subset), (Y, \preceq))$.

• *Map II.* We define a map

$$\Psi_{X,Y}: \text{Sets}(X, Y) \rightarrow \text{Pos}^{\text{cocomp}}((\mathcal{P}(X), \subset), (Y, \leq))$$

as in the statement, by

$$\Psi_{X,Y}(f) \stackrel{\text{def}}{=} \text{Lan}_{\chi_X}(f),$$


for each $f \in \text{Sets}(X, Y)$.

• *Invertibility I.* We claim that

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{Pos}^{\text{cocomp}}((\mathcal{P}(X), \subset), (Y, \leq))}.$$

Indeed, given a cocontinuous morphism of posets

$$\xi: (\mathcal{P}(X), \subset) \rightarrow (Y, \leq),$$

we have

$$\begin{aligned} [\Psi_{X,Y} \circ \Phi_{X,Y}](\xi) &\stackrel{\text{def}}{=} \Psi_{X,Y}(\Phi_{X,Y}(\xi)) \\ &\stackrel{\text{def}}{=} \Psi_{X,Y}(\xi \circ \chi_X) \\ &\stackrel{\text{def}}{=} \text{Lan}_{\chi_X}(\xi \circ \chi_X) \\ &\cong \bigvee_{x \in X} \chi_{(-)}(x) \odot \xi(\chi_X(x)) \\ &\stackrel{\text{dlim}}{=} \xi, \end{aligned}$$

where indeed

$$\begin{aligned} \left[\bigvee_{x \in X} \chi_{(-)}(x) \odot \xi(\chi_X(x)) \right](U) &\stackrel{\text{def}}{=} \bigvee_{x \in X} \chi_U(x) \odot \xi(\chi_X(x)) \\ &= \left(\bigvee_{x \in U} \chi_U(x) \odot \xi(\chi_X(x)) \right) \vee \left(\bigvee_{x \in X \setminus U} \chi_U(x) \odot \xi(\chi_X(x)) \right) \\ &= \left(\bigvee_{x \in U} \xi(\chi_X(x)) \right) \vee \left(\bigvee_{x \in X \setminus U} \emptyset_Y \right) \\ &= \bigvee_{x \in U} \xi(\chi_X(x)) \\ &\stackrel{(*)}{=} \xi \left(\bigvee_{x \in U} \chi_X(x) \right) \\ &= \xi(U) \end{aligned}$$

for each $U \in \mathcal{P}(X)$, where we have used that ξ is cocontinuous for the equality $\stackrel{(\dagger)}{=}$.

· *Invertibility II.* We claim that

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Sets}(X,Y)}.$$

Indeed, given a function $f: X \rightarrow Y$, we have

$$\begin{aligned} [\Phi_{X,Y} \circ \Psi_{X,Y}](f) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\Psi_{X,Y}(f)) \\ &\stackrel{\text{def}}{=} \Phi_{X,Y}(\text{Lan}_{\chi_X}(f)) \\ &\stackrel{\text{def}}{=} \text{Lan}_{\chi_X}(f) \circ \chi_X \\ &\stackrel{\text{clm}}{=} f, \end{aligned}$$

where indeed

$$\begin{aligned} [\text{Lan}_{\chi_X}(f) \circ \chi_X](x) &\stackrel{\text{def}}{=} \bigvee_{y \in X} \chi_{\{x\}}(y) \odot f(y) \\ &= (\chi_{\{x\}}(x) \odot f(x)) \vee \left(\bigvee_{y \in X \setminus \{x\}} \chi_{\{x\}}(y) \odot f(y) \right) \\ &= f(x) \vee \left(\bigvee_{y \in X \setminus \{x\}} \emptyset_Y \right) \\ &= f(x) \vee \emptyset_Y \\ &= f(x) \end{aligned}$$

for each $x \in X$.

· *Naturality for Φ , Part I.* We need to show that, given a function $f: X \rightarrow X'$, the diagram

$$\begin{array}{ccc} \text{Pos}^{\text{cocomp}}((\mathcal{P}(X'), \subset), (Y, \preceq)) & \xrightarrow{\Phi_{X',Y}} & \text{Sets}(X', Y) \\ \mathcal{P}_*(f)^* \downarrow & & \downarrow f^* \\ \text{Pos}^{\text{cocomp}}((\mathcal{P}(X), \subset), (Y, \preceq)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \end{array}$$

commutes. Indeed, given a cocontinuous morphism of posets

$$\xi: (\mathcal{P}(X'), \subset) \rightarrow (Y, \preceq),$$

we have

$$\begin{aligned}
[\Phi_{X,Y} \circ \mathcal{P}_*(f)^*](\xi) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\mathcal{P}_*(f)^*(\xi)) \\
&\stackrel{\text{def}}{=} \Phi_{X,Y}(\xi \circ f_*) \\
&\stackrel{\text{def}}{=} (\xi \circ f_*) \circ \chi_X \\
&= \xi \circ (f_* \circ \chi_X) \\
&\stackrel{(\dagger)}{=} \xi \circ (\chi_{X'} \circ f) \\
&= (\xi \circ \chi_{X'}) \circ f \\
&\stackrel{\text{def}}{=} \Phi_{X',Y}(\xi) \circ f \\
&\stackrel{\text{def}}{=} f^*(\Phi_{X',Y}(\xi)) \\
&\stackrel{\text{def}}{=} [f^* \circ \Phi_{X',Y}](\xi),
\end{aligned}$$

where we have used **Item 9** of **Proposition 2.4.1.3** for the equality $\stackrel{(\dagger)}{=}$ above.

- *Naturality for Φ , Part II.* We need to show that, given a cocontinuous morphism of posets

$$g: (Y, \preceq_Y) \rightarrow (Y', \preceq_{Y'}),$$

the diagram

$$\begin{array}{ccc}
\text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y, \preceq)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \\
\downarrow g_* & & \downarrow g_* \\
\text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y', \preceq)) & \xrightarrow{\Phi_{X,Y'}} & \text{Sets}(X, Y')
\end{array}$$


commutes. Indeed, given a cocontinuous morphism of posets

$$\xi: (\mathcal{P}(X), \subset) \rightarrow (Y, \preceq),$$

we have

$$\begin{aligned}
[\Phi_{X,Y'} \circ g_*](\xi) &\stackrel{\text{def}}{=} \Phi_{X,Y'}(g_*(\xi)) \\
&\stackrel{\text{def}}{=} \Phi_{X,Y'}(g \circ \xi) \\
&\stackrel{\text{def}}{=} (g \circ \xi) \circ \chi_X \\
&= g \circ (\xi \circ \chi_X) \\
&\stackrel{\text{def}}{=} g \circ (\Phi_{X,Y}(\xi)) \\
&\stackrel{\text{def}}{=} g_*(\Phi_{X,Y}(\xi)) \\
&\stackrel{\text{def}}{=} [g_* \circ \Phi_{X,Y}](\xi).
\end{aligned}$$

- *Naturality for Ψ* . Since Φ is natural in each argument and Φ is a componentwise inverse to Ψ in each argument, it follows from [Item 2 of Proposition 8.8.6.2](#) that Ψ is also natural in each argument.

This finishes the proof. 

007F 2.4.4 Direct Images

Let A and B be sets and let $f: A \rightarrow B$ be a function.

007G DEFINITION 2.4.4.1 ► DIRECT IMAGES

The **direct image function associated to f** is the function

$$f_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by^{1,2}

$$\begin{aligned} f_*(U) &\stackrel{\text{def}}{=} f(U) \\ &\stackrel{\text{def}}{=} \left\{ b \in B \mid \begin{array}{l} \text{there exists some } a \in U \\ \text{such that } b = f(a) \end{array} \right\} \\ &= \{f(a) \in B \mid a \in U\} \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

¹*Further Terminology:* The set $f(U)$ is called the **direct image of U by f** .

²We also have

$$f_*(U) = B \setminus f_*(A \setminus U);$$

see [Item 9 of Proposition 2.4.4.4](#).

007H NOTATION 2.4.4.2 ► FURTHER NOTATION FOR DIRECT IMAGES

Sometimes one finds the notation

$$\exists_f: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

for f_* . This notation comes from the fact that the following statements are equivalent, where $b \in B$ and $U \in \mathcal{P}(A)$:

- We have $b \in \exists_f(U)$.
- There exists some $a \in U$ such that $f(a) = b$.

007J

REMARK 2.4.4.3 ► UNWINDING DEFINITION 2.4.4.1

Identifying subsets of A with functions from A to $\{\text{true}, \text{false}\}$ via [Items 1 and 2](#) of [Proposition 2.4.3.9](#), we see that the direct image function associated to f is equivalently the function

$$f_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\begin{aligned} f_*(\chi_U) &\stackrel{\text{def}}{=} \text{Lan}_f(\chi_U) \\ &= \text{colim} \left(\left(f \overset{\rightarrow}{\times} \underline{(-1)} \right) \overset{\text{pr}}{\twoheadrightarrow} A \xrightarrow{\chi_U} \{\text{t}, \text{f}\} \right) \\ &= \text{colim}_{\substack{a \in A \\ f(a) = -1}} (\chi_U(a)) \\ &= \bigvee_{\substack{a \in A \\ f(a) = -1}} (\chi_U(a)), \end{aligned}$$

where we have used ?? for the second equality. In other words, we have

$$\begin{aligned} [f_*(\chi_U)](b) &= \bigvee_{\substack{a \in A \\ f(a) = b}} (\chi_U(a)) \\ &= \begin{cases} \text{true} & \text{if there exists some } a \in A \text{ such} \\ & \text{that } f(a) = b \text{ and } a \in U, \\ \text{false} & \text{otherwise} \end{cases} \\ &= \begin{cases} \text{true} & \text{if there exists some } a \in U \\ & \text{such that } f(a) = b, \\ \text{false} & \text{otherwise} \end{cases} \end{aligned}$$

for each $b \in B$.

007K

PROPOSITION 2.4.4.4 ► PROPERTIES OF DIRECT IMAGES I

Let $f: A \rightarrow B$ be a function.

007L

1. *Functoriality.* The assignment $U \mapsto f_*(U)$ defines a functor

$$f_*: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

where

007M

· *Action on Objects.* For each $U \in \mathcal{P}(A)$, we have

$$[f_*](U) \stackrel{\text{def}}{=} f_*(U).$$

· *Action on Morphisms.* For each $U, V \in \mathcal{P}(A)$:

(★) If $U \subset V$, then $f_*(U) \subset f_*(V)$.

2. *Triple Adjointness.* We have a triple adjunction

$$(f_* \dashv f^{-1} \dashv f_!): \quad \mathcal{P}(A) \begin{array}{c} \xrightarrow{f_*} \\ \perp \\ \xleftarrow{f^{-1}} \\ \perp \\ \xrightarrow{f_!} \end{array} \mathcal{P}(B),$$

witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(B)}(f_*(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(A)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f_!(V)), \end{aligned}$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$ and (respectively) $V \in \mathcal{P}(A)$ and $U \in \mathcal{P}(B)$, i.e. where:

(a) The following conditions are equivalent:

- i. We have $f_*(U) \subset V$.
- ii. We have $U \subset f^{-1}(V)$.

(b) The following conditions are equivalent:

- i. We have $f^{-1}(U) \subset V$.
- ii. We have $U \subset f_!(V)$.

007N

3. *Preservation of Colimits.* We have an equality of sets

$$f_*\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} f_*(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f_*(U) \cup f_*(V) &= f_*(U \cup V), \\ f_*(\emptyset) &= \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

007P

4. *Oplax Preservation of Limits.* We have an inclusion of sets

$$f_*\left(\bigcap_{i \in I} U_i\right) \subset \bigcap_{i \in I} f_*(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} f_*(U \cap V) &\subset f_*(U) \cap f_*(V), \\ f_*(A) &\subset B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

007Q

5. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$(f_*, f_*^\otimes, f_{*|\mathbb{1}}^\otimes): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} f_{*|U,V}^\otimes: f_*(U) \cup f_*(V) &\xrightarrow{=} f_*(U \cup V), \\ f_{*|\mathbb{1}}^\otimes: \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

007R

6. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric oplax monoidal structure

$$(f_*, f_*^\otimes, f_{*|\mathbb{1}}^\otimes): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with inclusions

$$\begin{aligned} f_{*|U,V}^\otimes: f_*(U \cap V) &\hookrightarrow f_*(U) \cap f_*(V), \\ f_{*|\mathbb{1}}^\otimes: f_*(A) &\hookrightarrow B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

007S

7. *Interaction With Coproducts.* Let $f: A \rightarrow A'$ and $g: B \rightarrow B'$ be maps of sets. We have

$$(f \amalg g)_*(U \amalg V) = f_*(U) \amalg g_*(V)$$

for each $U \in \mathcal{P}(A)$ and each $V \in \mathcal{P}(B)$.

007T

8. *Interaction With Products.* Let $f: A \rightarrow A'$ and $g: B \rightarrow B'$ be maps of sets. We have

$$(f \times g)_*(U \times V) = f_*(U) \times g_*(V)$$

for each $U \in \mathcal{P}(A)$ and each $V \in \mathcal{P}(B)$.

007U

9. *Relation to Direct Images With Compact Support.* We have

$$f_*(U) = B \setminus f_!(A \setminus U)$$

for each $U \in \mathcal{P}(A)$.

PROOF 2.4.4.5 ► PROOF OF PROPOSITION 2.4.4.4

Item 1: Functoriality

Clear.

Item 2: Triple Adjointness

This follows from [Remark 2.4.4.3](#), [Remark 2.4.5.2](#), [Remark 2.4.6.3](#), and ?? of ??.

Item 3: Preservation of Colimits

This follows from [Item 2](#) and ?? of ??.¹

Item 4: Oplax Preservation of Limits

The inclusion $f_*(A) \subset B$ is clear. See [[Pro24s](#)] for the other inclusions.

Item 5: Symmetric Strict Monoidality With Respect to Unions

This follows from [Item 3](#).

Item 6: Symmetric Oplax Monoidality With Respect to Intersections

This follows from [Item 4](#).

Item 7: Interaction With Coproducts

Clear.

Item 8: Interaction With Products

Clear.


Item 9: Relation to Direct Images With Compact Support

Applying [Item 9](#) of [Proposition 2.4.6.6](#) to $A \setminus U$, we have

$$\begin{aligned} f_!(A \setminus U) &= B \setminus f_*(A \setminus (A \setminus U)) \\ &= B \setminus f_*(U). \end{aligned}$$

Taking complements, we then obtain

$$\begin{aligned} f_*(U) &= B \setminus (B \setminus f_*(U)), \\ &= B \setminus f_!(A \setminus U), \end{aligned}$$

which finishes the proof. 

¹See also [Pro24t].

007V PROPOSITION 2.4.4.6 ► PROPERTIES OF DIRECT IMAGES II

Let $f: A \rightarrow B$ be a function.

007W 1. *Functionality I.* The assignment $f \mapsto f_*$ defines a function

$$(-)_{*|A,B}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

007X 2. *Functionality II.* The assignment $f \mapsto f_*$ defines a function

$$(-)_{*|A,B}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

007Y 3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$(\text{id}_A)_* = \text{id}_{\mathcal{P}(A)}.$$

007Z 4. *Interaction With Composition.* For each pair of composable functions $f: A \rightarrow B$ and $g: B \rightarrow C$, we have

$$(g \circ f)_* = g_* \circ f_*$$

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{f_*} & \mathcal{P}(B) \\ & \searrow (g \circ f)_* & \downarrow g_* \\ & & \mathcal{P}(C). \end{array}$$

PROOF 2.4.4.7 ► PROOF OF PROPOSITION 2.4.4.6

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

This follows from [Remark 2.4.4.3](#) and ?? of ??.

Item 4: Interaction With Composition

This follows from [Remark 2.4.4.3](#) and ?? of ??.



0080 2.4.5 Inverse Images

Let A and B be sets and let $f: A \rightarrow B$ be a function.

0081 DEFINITION 2.4.5.1 ► INVERSE IMAGES

The **inverse image function associated to f** is the function¹

$$f^{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by²

$$f^{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid \text{we have } f(a) \in V\}$$

for each $V \in \mathcal{P}(B)$.

¹Further Notation: Also written $f^*: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$.

²Further Terminology: The set $f^{-1}(V)$ is called the **inverse image of V by f** .

0082 REMARK 2.4.5.2 ► UNWINDING DEFINITION 2.4.5.1

Identifying subsets of B with functions from B to $\{\text{true}, \text{false}\}$ via [Items 1 and 2](#) of [Proposition 2.4.3.9](#), we see that the inverse image function associated to f is equivalently the function

$$f^*: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by

$$f^*(\chi_V) \stackrel{\text{def}}{=} \chi_V \circ f$$

for each $\chi_V \in \mathcal{P}(B)$, where $\chi_V \circ f$ is the composition

$$A \xrightarrow{f} B \xrightarrow{\chi_V} \{\text{true}, \text{false}\}$$

in Sets.

0083 PROPOSITION 2.4.5.3 ► PROPERTIES OF INVERSE IMAGES I

Let $f: A \rightarrow B$ be a function.

0084 1. *Functoriality.* The assignment $V \mapsto f^{-1}(V)$ defines a functor

$$f^{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

where

• *Action on Objects.* For each $V \in \mathcal{P}(B)$, we have

$$[f^{-1}](V) \stackrel{\text{def}}{=} f^{-1}(V).$$

• *Action on Morphisms.* For each $U, V \in \mathcal{P}(B)$:

(★) If $U \subset V$, then $f^{-1}(U) \subset f^{-1}(V)$.

0085 2. *Triple Adjointness.* We have a triple adjunction

$$(f_* \dashv f^{-1} \dashv f_!): \mathcal{P}(A) \leftarrow f^{-1} \text{ --- } \mathcal{P}(B),$$

witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(B)}(f_*(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(A)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f_!(V)), \end{aligned}$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$ and (respectively) $V \in \mathcal{P}(A)$ and $U \in \mathcal{P}(B)$, i.e. where:

(a) The following conditions are equivalent:

- i. We have $f_*(U) \subset V$;
- ii. We have $U \subset f^{-1}(V)$;

(b) The following conditions are equivalent:

- i. We have $f^{-1}(U) \subset V$.
- ii. We have $U \subset f_!(V)$.

0086 3. *Preservation of Colimits.* We have an equality of sets

$$f^{-1}\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} f^{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f^{-1}(U) \cup f^{-1}(V) &= f^{-1}(U \cup V), \\ f^{-1}(\emptyset) &= \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

0087 4. *Preservation of Limits.* We have an equality of sets

$$f^{-1}\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} f^{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f^{-1}(U) \cap f^{-1}(V) &= f^{-1}(U \cap V), \\ f^{-1}(B) &= A, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

0088 5. *Symmetric Strict Monoidality With Respect to Unions.* The inverse image function of **Item 1** has a symmetric strict monoidal structure

$$\left(f^{-1}, f^{-1, \otimes}, f_{\mathbb{1}}^{-1, \otimes}\right): (\mathcal{P}(B), \cup, \emptyset) \rightarrow (\mathcal{P}(A), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} f_{U, V}^{-1, \otimes}: f^{-1}(U) \cup f^{-1}(V) &\xrightarrow{=} f^{-1}(U \cup V), \\ f_{\mathbb{1}}^{-1, \otimes}: \emptyset &\xrightarrow{=} f^{-1}(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

0089 6. *Symmetric Strict Monoidality With Respect to Intersections.* The inverse image function of **Item 1** has a symmetric strict monoidal structure

$$\left(f^{-1}, f^{-1, \otimes}, f_{\mathbb{1}}^{-1, \otimes}\right): (\mathcal{P}(B), \cap, B) \rightarrow (\mathcal{P}(A), \cap, A),$$

being equipped with equalities

$$\begin{aligned} f_{U, V}^{-1, \otimes}: f^{-1}(U) \cap f^{-1}(V) &\xrightarrow{=} f^{-1}(U \cap V), \\ f_{\mathbb{1}}^{-1, \otimes}: A &\xrightarrow{=} f^{-1}(B), \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

008A

7. *Interaction With Coproducts.* Let $f: A \rightarrow A'$ and $g: B \rightarrow B'$ be maps of sets. We have

$$(f \amalg g)^{-1}(U' \amalg V') = f^{-1}(U') \amalg g^{-1}(V')$$

for each $U' \in \mathcal{P}(A')$ and each $V' \in \mathcal{P}(B')$.

008B

8. *Interaction With Products.* Let $f: A \rightarrow A'$ and $g: B \rightarrow B'$ be maps of sets. We have

$$(f \times g)^{-1}(U' \times V') = f^{-1}(U') \times g^{-1}(V')$$

for each $U' \in \mathcal{P}(A')$ and each $V' \in \mathcal{P}(B')$.

PROOF 2.4.5.4 ► PROOF OF PROPOSITION 2.4.5.3

Item 1: Functoriality

Clear.

Item 2: Triple Adjointness

This follows from [Remark 2.4.4.3](#), [Remark 2.4.5.2](#), [Remark 2.4.6.3](#), and ?? of ??.

Item 3: Preservation of Colimits

This follows from [Item 2](#) and ?? of ??.¹

Item 4: Preservation of Limits

This follows from [Item 2](#) and ?? of ??.²

Item 5: Symmetric Strict Monoidality With Respect to Unions

This follows from [Item 3](#).

Item 6: Symmetric Strict Monoidality With Respect to Intersections

This follows from [Item 4](#).

Item 7: Interaction With Coproducts

Clear.

Item 8: Interaction With Products

Clear. 

¹See also [\[Pro24ac\]](#).

²See also [\[Pro24ab\]](#).

008C **PROPOSITION 2.4.5.5 ► PROPERTIES OF INVERSE IMAGES II**

Let $f: A \rightarrow B$ be a function.

008D 1. *Functionality I.* The assignment $f \mapsto f^{-1}$ defines a function

$$(-)_{A,B}^{-1}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(B), \mathcal{P}(A)).$$

008E 2. *Functionality II.* The assignment $f \mapsto f^{-1}$ defines a function

$$(-)_{A,B}^{-1}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(B), \subset), (\mathcal{P}(A), \subset)).$$

008F 3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$\text{id}_A^{-1} = \text{id}_{\mathcal{P}(A)}.$$

008G 4. *Interaction With Composition.* For each pair of composable functions $f: A \rightarrow B$ and $g: B \rightarrow C$, we have

$$(g \circ f)^{-1} = f^{-1} \circ g^{-1},$$

$$\begin{array}{ccc} \mathcal{P}(C) & \xrightarrow{g^{-1}} & \mathcal{P}(B) \\ & \searrow (g \circ f)^{-1} & \downarrow f^{-1} \\ & & \mathcal{P}(A). \end{array}$$

PROOF 2.4.5.6 ► PROOF OF PROPOSITION 2.4.5.5

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

This follows from [Remark 2.4.5.2](#) and [Item 5 of Proposition 8.1.6.2](#).

Item 4: Interaction With Composition

This follows from [Remark 2.4.5.2](#) and [Item 2 of Proposition 8.1.6.2](#). 

008H 2.4.6 Direct Images With Compact Support

Let A and B be sets and let $f: A \rightarrow B$ be a function.

008J

DEFINITION 2.4.6.1 ► DIRECT IMAGES WITH COMPACT SUPPORT

The **direct image with compact support function associated to** f is the function

$$f_! : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by^{1,2}

$$\begin{aligned} f_!(U) &\stackrel{\text{def}}{=} \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ if we have} \\ f(a) = b, \text{ then } a \in U \end{array} \right\} \\ &= \{ b \in B \mid \text{we have } f^{-1}(b) \subset U \} \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

¹*Further Terminology:* The set $f_!(U)$ is called the **direct image with compact support of** U **by** f .

²We also have

$$f_!(U) = B \setminus f_*(A \setminus U);$$

see [Item 9 of Proposition 2.4.6.6](#).

008K

NOTATION 2.4.6.2 ► FURTHER NOTATION FOR DIRECT IMAGES WITH COMPACT SUPPORT

Sometimes one finds the notation

$$\forall_f : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

for f_* . This notation comes from the fact that the following statements are equivalent, where $b \in B$ and $U \in \mathcal{P}(A)$:

- We have $b \in \forall_f(U)$.
- For each $a \in A$, if $b = f(a)$, then $a \in U$.

008L

REMARK 2.4.6.3 ► UNWINDING DEFINITION 2.4.6.1

Identifying subsets of A with functions from A to $\{\text{true}, \text{false}\}$ via [Items 1 and 2](#) of [Proposition 2.4.3.9](#), we see that the direct image with compact support function associated to f is equivalently the function

$$f_! : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\begin{aligned}
 f_i(\chi_U) &\stackrel{\text{def}}{=} \text{Ran}_f(\chi_U) \\
 &= \lim \left(\left(\underline{(-1)} \vec{\times} f \right) \xrightarrow{\text{pr}} A \xrightarrow{\chi_U} \{\text{true}, \text{false}\} \right) \\
 &= \lim_{\substack{a \in A \\ f(a) = -1}} (\chi_U(a)) \\
 &= \bigwedge_{\substack{a \in A \\ f(a) = -1}} (\chi_U(a)).
 \end{aligned}$$

where we have used ?? for the second equality. In other words, we have

$$\begin{aligned}
 [f_i(\chi_U)](b) &= \bigwedge_{\substack{a \in A \\ f(a) = b}} (\chi_U(a)) \\
 &= \begin{cases} \text{true} & \text{if, for each } a \in A \text{ such that} \\ & f(a) = b, \text{ we have } a \in U, \\ \text{false} & \text{otherwise} \end{cases} \\
 &= \begin{cases} \text{true} & \text{if } f^{-1}(b) \subset U \\ \text{false} & \text{otherwise} \end{cases}
 \end{aligned}$$

for each $b \in B$.

008M

DEFINITION 2.4.6.4 ► THE IMAGE AND COMPLEMENT PARTS OF f_i

Let U be a subset of A .^{1,2}

008N

1. The **image part of the direct image with compact support** $f_i(U)$ of U is the set $f_{i,\text{im}}(U)$ defined by

$$\begin{aligned}
 f_{i,\text{im}}(U) &\stackrel{\text{def}}{=} f_i(U) \cap \text{Im}(f) \\
 &= \left\{ b \in B \mid \begin{array}{l} \text{we have } f^{-1}(b) \subset U \\ \text{and } f^{-1}(b) \neq \emptyset \end{array} \right\}.
 \end{aligned}$$

2. The **complement part of the direct image with compact support** $f_i(U)$

008P

of U is the set $f_{!,\text{cp}}(U)$ defined by

$$\begin{aligned} f_{!,\text{cp}}(U) &\stackrel{\text{def}}{=} f_!(U) \cap (B \setminus \text{Im}(f)) \\ &= B \setminus \text{Im}(f) \\ &= \left\{ b \in B \mid \begin{array}{l} \text{we have } f^{-1}(b) \subset U \\ \text{and } f^{-1}(b) = \emptyset \end{array} \right\} \\ &= \{ b \in B \mid f^{-1}(b) = \emptyset \}. \end{aligned}$$

¹Note that we have

$$f_!(U) = f_{!,\text{im}}(U) \cup f_{!,\text{cp}}(U),$$

as

$$\begin{aligned} f_!(U) &= f_!(U) \cap B \\ &= f_!(U) \cap (\text{Im}(f) \cup (B \setminus \text{Im}(f))) \\ &= (f_!(U) \cap \text{Im}(f)) \cup (f_!(U) \cap (B \setminus \text{Im}(f))) \\ &\stackrel{\text{def}}{=} f_{!,\text{im}}(U) \cup f_{!,\text{cp}}(U). \end{aligned}$$

²In terms of the meet computation of $f_!(U)$ of [Remark 2.4.6.3](#), namely

$$f_!(\chi_U) = \bigwedge_{\substack{a \in A \\ f(a) = -1}} (\chi_U(a)),$$

we see that $f_{!,\text{im}}$ corresponds to meets indexed over nonempty sets, while $f_{!,\text{cp}}$ corresponds to meets indexed over the empty set.

008Q

EXAMPLE 2.4.6.5 ► EXAMPLES OF DIRECT IMAGES WITH COMPACT SUPPORT

Here are some examples of direct images with compact support.

1. *The Multiplication by Two Map on the Natural Numbers.* Consider the function $f: \mathbb{N} \rightarrow \mathbb{N}$ given by

$$f(n) \stackrel{\text{def}}{=} 2n$$

for each $n \in \mathbb{N}$. Since f is injective, we have

$$\begin{aligned} f_{!,\text{im}}(U) &= f_*(U) \\ f_{!,\text{cp}}(U) &= \{\text{odd natural numbers}\} \end{aligned}$$

for any $U \subset \mathbb{N}$.

2. *Parabolas.* Consider the function $f: \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) \stackrel{\text{def}}{=} x^2$$

for each $x \in \mathbb{R}$. We have

$$f_{!,\text{cp}}(U) = \mathbb{R}_{<0}$$

for any $U \subset \mathbb{R}$. Moreover, since $f^{-1}(x) = \{-\sqrt{x}, \sqrt{x}\}$, we have e.g.:

$$\begin{aligned} f_{!,\text{im}}([0, 1]) &= \{0\}, \\ f_{!,\text{im}}([-1, 1]) &= [0, 1], \\ f_{!,\text{im}}([1, 2]) &= \emptyset, \\ f_{!,\text{im}}([-2, -1] \cup [1, 2]) &= [1, 4]. \end{aligned}$$

3. *Circles.* Consider the function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ given by

$$f(x, y) \stackrel{\text{def}}{=} x^2 + y^2$$

for each $(x, y) \in \mathbb{R}^2$. We have

$$f_{!,\text{cp}}(U) = \mathbb{R}_{<0}$$

for any $U \subset \mathbb{R}^2$, and since

$$f^{-1}(r) = \begin{cases} \text{a circle of radius } r \text{ about the origin} & \text{if } r > 0, \\ \{(0, 0)\} & \text{if } r = 0, \\ \emptyset & \text{if } r < 0, \end{cases}$$

we have e.g.:

$$\begin{aligned} f_{!,\text{im}}([-1, 1] \times [-1, 1]) &= [0, 1], \\ f_{!,\text{im}}([-1, 1] \times [-1, 1] \setminus [-1, 1] \times \{0\}) &= \emptyset. \end{aligned}$$

008R PROPOSITION 2.4.6.6 ► PROPERTIES OF DIRECT IMAGES WITH COMPACT SUPPORT I

Let $f: A \rightarrow B$ be a function.

008S 1. *Functoriality.* The assignment $U \mapsto f_!(U)$ defines a functor

$$f_!: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

where

• *Action on Objects.* For each $U \in \mathcal{P}(A)$, we have

$$[f_!](U) \stackrel{\text{def}}{=} f_!(U).$$

008T

· *Action on Morphisms.* For each $U, V \in \mathcal{P}(A)$:

(★) If $U \subset V$, then $f_!(U) \subset f_!(V)$.

2. *Triple Adjointness.* We have a triple adjunction

$$(f_* \dashv f^{-1} \dashv f_!): \quad \mathcal{P}(A) \begin{array}{c} \xrightarrow{f_*} \\ \perp \\ \xleftarrow{f^{-1}} \\ \perp \\ \xrightarrow{f_!} \end{array} \mathcal{P}(B),$$

witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(B)}(f_*(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(A)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f_!(V)), \end{aligned}$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$ and (respectively) $V \in \mathcal{P}(A)$ and $U \in \mathcal{P}(B)$, i.e. where:

(a) The following conditions are equivalent:

- i. We have $f_*(U) \subset V$.
- ii. We have $U \subset f^{-1}(V)$.

(b) The following conditions are equivalent:

- i. We have $f^{-1}(U) \subset V$.
- ii. We have $U \subset f_!(V)$.

008U

3. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} f_!(U_i) \subset f_!\left(\bigcup_{i \in I} U_i\right),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} f_!(U) \cup f_!(V) &\hookrightarrow f_!(U \cup V), \\ \emptyset &\hookrightarrow f_!(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

008V

4. *Preservation of Limits.* We have an equality of sets

$$f_!\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} f_!(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f^{-1}(U \cap V) &= f_!(U) \cap f^{-1}(V), \\ f_!(A) &= B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

008W

5. *Symmetric Lax Monoidality With Respect to Unions.* The direct image with compact support function of **Item 1** has a symmetric lax monoidal structure

$$\left(f_!, f_!^\otimes, f_{!1}^\otimes\right): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with inclusions

$$\begin{aligned} f_{!|U,V}^\otimes: f_!(U) \cup f_!(V) &\hookrightarrow f_!(U \cup V), \\ f_{!1}^\otimes: \emptyset &\hookrightarrow f_!(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

008X

6. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$\left(f_!, f_!^\otimes, f_{!1}^\otimes\right): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with equalities

$$\begin{aligned} f_{!|U,V}^\otimes: f_!(U \cap V) &\xrightarrow{=} f_!(U) \cap f_!(V), \\ f_{!1}^\otimes: f_!(A) &\xrightarrow{=} B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

008Y

7. *Interaction With Coproducts.* Let $f: A \rightarrow A'$ and $g: B \rightarrow B'$ be maps of sets. We have

$$(f \amalg g)_!(U \amalg V) = f_!(U) \amalg g_!(V)$$

for each $U \in \mathcal{P}(A)$ and each $V \in \mathcal{P}(B)$.

008Z

8. *Interaction With Products.* Let $f: A \rightarrow A'$ and $g: B \rightarrow B'$ be maps of sets. We have

$$(f \times g)_!(U \times V) = f_!(U) \times g_!(V)$$

for each $U \in \mathcal{P}(A)$ and each $V \in \mathcal{P}(B)$.

0090 9. *Relation to Direct Images.* We have

$$f_! (U) = B \setminus f_* (A \setminus U)$$

for each $U \in \mathcal{P}(A)$.

0091 10. *Interaction With Injections.* If f is injective, then we have

$$\begin{aligned} f_{!,\text{im}}(U) &= f_*(U), \\ f_{!,\text{cp}}(U) &= B \setminus \text{Im}(f), \\ f_!(U) &= f_{!,\text{im}}(U) \cup f_{!,\text{cp}}(U) \\ &= f_*(U) \cup (B \setminus \text{Im}(f)) \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

0092 11. *Interaction With Surjections.* If f is surjective, then we have

$$\begin{aligned} f_{!,\text{im}}(U) &\subset f_*(U), \\ f_{!,\text{cp}}(U) &= \emptyset, \\ f_!(U) &\subset f_*(U) \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

PROOF 2.4.6.7 ► PROOF OF PROPOSITION 2.4.6.6

Item 1: Functoriality

Clear.

Item 2: Triple Adjointness

This follows from [Remark 2.4.4.3](#), [Remark 2.4.5.2](#), [Remark 2.4.6.3](#), and ?? of ??.

Item 3: Lax Preservation of Colimits

Omitted.

Item 4: Preservation of Limits

This follows from [Item 2](#) and ?? of ??.

Item 5: Symmetric Lax Monoidality With Respect to Unions

This follows from [Item 3](#).

Item 6: Symmetric Strict Monoidality With Respect to Intersections

This follows from [Item 4](#).

Item 7: Interaction With Coproducts

Clear.

Item 8: Interaction With Products

Clear.

Item 9: Relation to Direct Images

We claim that $f_!(U) = B \setminus f_*(A \setminus U)$.

- *The First Implication.* We claim that

$$f_!(U) \subset B \setminus f_*(A \setminus U).$$

Let $b \in f_!(U)$. We need to show that $b \notin f_*(A \setminus U)$, i.e. that there is no $a \in A \setminus U$ such that $f(a) = b$.

This is indeed the case, as otherwise we would have $a \in f^{-1}(b)$ and $a \notin U$, contradicting $f^{-1}(b) \subset U$ (which holds since $b \in f_!(U)$).

Thus $b \in B \setminus f_*(A \setminus U)$.

- *The Second Implication.* We claim that

$$B \setminus f_*(A \setminus U) \subset f_!(U).$$

Let $b \in B \setminus f_*(A \setminus U)$. We need to show that $b \in f_!(U)$, i.e. that $f^{-1}(b) \subset U$.

Since $b \notin f_*(A \setminus U)$, there exists no $a \in A \setminus U$ such that $b = f(a)$, and hence $f^{-1}(b) \subset U$.

Thus $b \in f_!(U)$.

This finishes the proof of **Item 9**.

Item 10: Interaction With Injections

Clear.

Item 11: Interaction With Surjections

Clear. 

0093

PROPOSITION 2.4.6.8 ► PROPERTIES OF DIRECT IMAGES WITH COMPACT SUPPORT II

Let $f: A \rightarrow B$ be a function.

0094

1. *Functionality I.* The assignment $f \mapsto f_!$ defines a function

$$(-)_{!|A,B}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

0095

2. *Functionality II.* The assignment $f \mapsto f_!$ defines a function

$$(-)_{!|A,B}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

0096

3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$(\text{id}_A)_! = \text{id}_{\mathcal{P}(A)}.$$

0097

4. *Interaction With Composition.* For each pair of composable functions $f: A \rightarrow B$ and $g: B \rightarrow C$, we have

$$(g \circ f)_! = g_! \circ f_!$$

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{f_!} & \mathcal{P}(B) \\ & \searrow (g \circ f)_! & \downarrow g_! \\ & & \mathcal{P}(C). \end{array}$$

PROOF 2.4.6.9 ► PROOF OF PROPOSITION 2.4.6.8

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

This follows from [Remark 2.4.6.3](#) and ?? of ??.

Item 4: Interaction With Composition

This follows from [Remark 2.4.6.3](#) and ?? of ??.



Appendices

2.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets

Relations

5. Relations

6. Constructions With Relations

7. Equivalence Relations and Apartness Relations

Category Theory

8. Categories

Bicategories

9. Types of Morphisms in Bicategories

Chapter 3

Pointed Sets

0098 This chapter contains some foundational material on pointed sets.

Contents

3.1	Pointed Sets	120
3.1.1	Foundations	120
3.1.2	Morphisms of Pointed Sets	122
3.1.3	The Category of Pointed Sets	122
3.1.4	Elementary Properties of Pointed Sets	123
3.2	Limits of Pointed Sets	126
3.2.1	The Terminal Pointed Set	126
3.2.2	Products of Families of Pointed Sets	127
3.2.3	Products	129
3.2.4	Pullbacks	131
3.2.5	Equalisers	136
3.3	Colimits of Pointed Sets	139
3.3.1	The Initial Pointed Set	139
3.3.2	Coproducts of Families of Pointed Sets	140
3.3.3	Coproducts	142
3.3.4	Pushouts	146
3.3.5	Coequalisers	151
3.4	Constructions With Pointed Sets	153
3.4.1	Free Pointed Sets	153
3.A	Other Chapters	159

0099 **3.1 Pointed Sets**

009A **3.1.1 Foundations**

009B

DEFINITION 3.1.1.1 ► POINTED SETS

A **pointed set**¹ is equivalently:

- An \mathbb{E}_0 -monoid in $(\mathbf{N}_\bullet(\mathbf{Sets}), \text{pt})$.
- A pointed object in $(\mathbf{Sets}, \text{pt})$.

¹*Further Terminology:* In the context of monoids with zero as models for \mathbb{F}_1 -algebras, pointed sets are viewed as \mathbb{F}_1 -**modules**.

009C

REMARK 3.1.1.2 ► UNWINDING DEFINITION 3.1.1.1

In detail, a **pointed set** is a pair (X, x_0) consisting of:

- *The Underlying Set.* A set X , called the **underlying set of** (X, x_0) .
- *The Basepoint.* A morphism

$$[x_0]: \text{pt} \rightarrow X$$

in \mathbf{Sets} , determining an element $x_0 \in X$, called the **basepoint of** X .

009D

EXAMPLE 3.1.1.3 ► THE ZERO SPHERE

The **0-sphere**¹ is the pointed set $(S^0, 0)$ ² consisting of:

- *The Underlying Set.* The set S^0 defined by

$$S^0 \stackrel{\text{def}}{=} \{0, 1\}.$$

- *The Basepoint.* The element 0 of S^0 .

¹*Further Terminology:* In the context of monoids with zero as models for \mathbb{F}_1 -algebras, the 0-sphere is viewed as the **underlying pointed set of the field with one element**.

²*Further Notation:* In the context of monoids with zero as models for \mathbb{F}_1 -algebras, S^0 is also denoted $(\mathbb{F}_1, 0)$.

009E

EXAMPLE 3.1.1.4 ► THE TRIVIAL POINTED SET

The **trivial pointed set** is the pointed set (pt, \star) consisting of:

- *The Underlying Set.* The punctual set $\text{pt} \stackrel{\text{def}}{=} \{\star\}$.
- *The Basepoint.* The element \star of pt .

009F EXAMPLE 3.1.1.5 ► THE UNDERLYING POINTED SET OF A SEMIMODULE

The **underlying pointed set** of a semimodule (M, α_M) is the pointed set $(M, 0_M)$.

009G EXAMPLE 3.1.1.6 ► THE UNDERLYING POINTED SET OF A MODULE

The **underlying pointed set** of a module (M, α_M) is the pointed set $(M, 0_M)$.

009H 3.1.2 Morphisms of Pointed Sets

009J DEFINITION 3.1.2.1 ► MORPHISMS OF POINTED SETS

A **morphism of pointed sets**^{1,2} is equivalently:

- A morphism of \mathbb{E}_0 -monoids in $(\mathbf{N}_\bullet(\mathbf{Sets}), \text{pt})$.
- A morphism of pointed objects in $(\mathbf{Sets}, \text{pt})$.

¹Further Terminology: Also called a **pointed function**.

²Further Terminology: In the context of monoids with zero as models for \mathbb{F}_1 -algebras, morphisms of pointed sets are also called **morphism of \mathbb{F}_1 -modules**.

009K REMARK 3.1.2.2 ► UNWINDING DEFINITION 3.1.2.1

In detail, a **morphism of pointed sets** $f: (X, x_0) \rightarrow (Y, y_0)$ is a morphism of sets $f: X \rightarrow Y$ such that the diagram

$$\begin{array}{ccc} & \text{pt} & \\ [x_0] \swarrow & & \searrow [y_0] \\ X & \xrightarrow{f} & Y \end{array}$$

commutes, i.e. such that

$$f(x_0) = y_0.$$

009L 3.1.3 The Category of Pointed Sets

009M DEFINITION 3.1.3.1 ► THE CATEGORY OF POINTED SETS

The **category of pointed sets** is the category \mathbf{Sets}_* defined equivalently as

- The homotopy category of the ∞ -category $\text{Mon}_{\mathbb{E}_0}(\mathbf{N}_\bullet(\mathbf{Sets}), \text{pt})$ of ??;

· The category Sets_* of ??.

009N

REMARK 3.1.3.2 ► UNWINDING DEFINITION 3.1.3.1

In detail, the **category of pointed sets** is the category Sets_* where

- *Objects.* The objects of Sets_* are pointed sets;
- *Morphisms.* The morphisms of Sets_* are morphisms of pointed sets;
- *Identities.* For each $(X, x_0) \in \text{Obj}(\text{Sets}_*)$, the unit map

$$\mathbb{1}_{(X, x_0)}^{\text{Sets}_*} : \text{pt} \rightarrow \text{Sets}_*((X, x_0), (X, x_0))$$

of Sets_* at (X, x_0) is defined by¹

$$\text{id}_{(X, x_0)}^{\text{Sets}_*} \stackrel{\text{def}}{=} \text{id}_X;$$

- *Composition.* For each $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$, the composition map

$$\circ_{(X, x_0), (Y, y_0), (Z, z_0)}^{\text{Sets}_*} : \text{Sets}_*((Y, y_0), (Z, z_0)) \times \text{Sets}_*((X, x_0), (Y, y_0)) \rightarrow \text{Sets}_*((X, x_0), (Z, z_0))$$

of Sets_* at $((X, x_0), (Y, y_0), (Z, z_0))$ is defined by²


$$g \circ_{(X, x_0), (Y, y_0), (Z, z_0)}^{\text{Sets}_*} f \stackrel{\text{def}}{=} g \circ f.$$

¹Note that id_X is indeed a morphism of pointed sets, as we have $\text{id}_X(x_0) = x_0$.

²Note that the composition of two morphisms of pointed sets is indeed a morphism of pointed sets, as we have

$$\begin{aligned} g(f(x_0)) &= g(y_0) \\ &= z_0, \end{aligned}$$

or



in terms of diagrams.

009P 3.1.4 Elementary Properties of Pointed Sets

009Q **PROPOSITION 3.1.4.1 ► ELEMENTARY PROPERTIES OF POINTED SETS**

Let (X, x_0) be a pointed set.

- 009R 1. *Completeness.* The category \mathbf{Sets}_* of pointed sets and morphisms between them is complete, having in particular:
- 009S (a) Products, described as in [Definition 3.2.3.1](#);
- 009T (b) Pullbacks, described as in [Definition 3.2.4.1](#);
- 009U (c) Equalisers, described as in [Definition 3.2.5.1](#).
- 009V 2. *Cocompleteness.* The category \mathbf{Sets}_* of pointed sets and morphisms between them is cocomplete, having in particular:
- 009W (a) Coproducts, described as in [Definition 3.3.3.1](#);
- 009X (b) Pushouts, described as in [Definition 3.3.4.1](#);
- 009Y (c) Coequalisers, described as in [Definition 3.3.5.1](#).
- 009Z 3. *Failure To Be Cartesian Closed.* The category \mathbf{Sets}_* is not Cartesian closed.¹
- 00A0 4. *Morphisms From the Monoidal Unit.* We have a bijection of sets²

$$\mathbf{Sets}_*(S^0, X) \cong X,$$

natural in $(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)$, internalising also to an isomorphism of pointed sets

$$\mathbf{Sets}_*(S^0, X) \cong (X, x_0),$$

again natural in $(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)$.

- 00A1 5. *Relation to Partial Functions.* We have an equivalence of categories³

$$\mathbf{Sets}_* \stackrel{\text{eq.}}{\cong} \mathbf{Sets}^{\text{part.}}$$

between the category of pointed sets and pointed functions between them and the category of sets and partial functions between them, where:

- (a) *From Pointed Sets to Sets With Partial Functions.* The equivalence

$$\zeta: \mathbf{Sets}_* \xrightarrow{\cong} \mathbf{Sets}^{\text{part.}}$$

sends:

- i. A pointed set (X, x_0) to X .
- ii. A pointed function

$$f: (X, x_0) \rightarrow (Y, y_0)$$

to the partial function

$$\xi_f: X \rightarrow Y$$

defined on $f^{-1}(Y \setminus y_0)$ and given by

$$\xi_f(x) \stackrel{\text{def}}{=} f(x)$$

for each $x \in f^{-1}(Y \setminus y_0)$.

(b) *From Sets With Partial Functions to Pointed Sets.* The equivalence

$$\xi^{-1}: \text{Sets}^{\text{part.}} \xrightarrow{\cong} \text{Sets}_*$$

sends:

- i. A set X is to the pointed set (X, \star) with \star an element that is not in X .
- ii. A partial function

$$f: X \rightarrow Y$$

defined on $U \subset X$ to the pointed function

$$\xi_f^{-1}: (X, x_0) \rightarrow (Y, y_0)$$

defined by

$$\xi_f^{-1}(x) \stackrel{\text{def}}{=} \begin{cases} f(x) & \text{if } x \in U, \\ y_0 & \text{otherwise.} \end{cases}$$


for each $x \in X$.

¹The category Sets_* does admit monoidal closed structures however; see [Tensor Products of Pointed Sets](#).

²In other words, the forgetful functor

$$\text{忘}: \text{Sets}_* \rightarrow \text{Sets}$$

defined on objects by sending a pointed set to its underlying set is corepresentable by S^0 .

³ *Warning:* This is not an isomorphism of categories, only an equivalence.

END TEXTDBEND

PROOF 3.1.4.2 ► PROOF OF PROPOSITION 3.1.4.1

Item 1: Completeness

This follows from (the proofs) of [Definitions 3.2.3.1, 3.2.4.1 and 3.2.5.1](#) and ??.

Item 2: Cocompleteness

This follows from (the proofs) of [Definitions 3.3.3.1, 3.3.4.1 and 3.3.5.1](#) and ??.

Item 3: Failure To Be Cartesian Closed

See [[MSE 2855868](#)].

Item 4: Morphisms From the Monoidal Unit

Since a morphism from S^0 to a pointed set (X, x_0) sends $0 \in S^0$ to x_0 and then can send $1 \in S^0$ to any element of X , we obtain a bijection between pointed maps $S^0 \rightarrow X$ and the elements of X .

The isomorphism then

$$\mathbf{Sets}_*(S^0, X) \cong (X, x_0)$$

follows by noting that $\Delta_{x_0} : S^0 \rightarrow X$, the basepoint of $\mathbf{Sets}_*(S^0, X)$, corresponds to the pointed map $S^0 \rightarrow X$ picking the element x_0 of X , and thus we see that the bijection between pointed maps $S^0 \rightarrow X$ and elements of X is compatible with basepoints, lifting to an isomorphism of pointed sets.

Item 5: Relation to Partial Functions

See [[MSE 884460](#)].



00A2 3.2 Limits of Pointed Sets

00A3 3.2.1 The Terminal Pointed Set

00A4 DEFINITION 3.2.1.1 ► THE TERMINAL POINTED SET

The **terminal pointed set** is the pair $\left((\text{pt}, \star), \{!_X\}_{(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)} \right)$ consisting of:

- *The Limit.* The pointed set (pt, \star) .
- *The Cone.* The collection of morphisms of pointed sets

$$\{!_X : (X, x_0) \rightarrow (\text{pt}, \star)\}_{(X, x_0) \in \text{Obj}(\mathbf{Sets})}$$

defined by

$$!_X(x) \stackrel{\text{def}}{=} \star$$

for each $x \in X$ and each $(X, x_0) \in \text{Obj}(\text{Sets})$.

PROOF 3.2.1.2 ► PROOF OF DEFINITION 3.2.1.1

We claim that (pt, \star) is the terminal object of Sets_* . Indeed, suppose we have a diagram of the form


$$(X, x_0) \quad (\text{pt}, \star)$$

in Sets_* . Then there exists a unique morphism of pointed sets

$$\phi: (X, x_0) \rightarrow (\text{pt}, \star)$$

making the diagram

$$(X, x_0) \xrightarrow[\exists!]{\phi} (\text{pt}, \star)$$

commute, namely $!_X$. 

00A5 3.2.2 Products of Families of Pointed Sets

Let $\{(X_i, x_0^i)\}_{i \in I}$ be a family of pointed sets.

00A6 DEFINITION 3.2.2.1 ► THE PRODUCT OF A FAMILY OF POINTED SETS

The **product** of $\{(X_i, x_0^i)\}_{i \in I}$ is the pair $((\prod_{i \in I} X_i, (x_0^i)_{i \in I}), \{\text{pr}_i\}_{i \in I})$ consisting of:

- *The Limit.* The pointed set $(\prod_{i \in I} X_i, (x_0^i)_{i \in I})$.
- *The Cone.* The collection

$$\left\{ \text{pr}_i : \left(\prod_{i \in I} X_i, (x_0^i)_{i \in I} \right) \rightarrow (X_i, x_0^i) \right\}_{i \in I}$$

of maps given by

$$\text{pr}_i \left((x_j)_{j \in I} \right) \stackrel{\text{def}}{=} x_i$$

for each $(x_j)_{j \in I} \in \prod_{i \in I} X_i$ and each $i \in I$.

PROOF 3.2.2.2 ► PROOF OF DEFINITION 3.2.2.1

We claim that $(\prod_{i \in I} X_i, (x_0^i)_{i \in I})$ is the categorical product of $\{(X_i, x_0^i)\}_{i \in I}$ in \mathbf{Sets}_* . Indeed, suppose we have, for each $i \in I$, a diagram of the form

$$\begin{array}{ccc} (P, *) & & \\ & \searrow p_i & \\ (\prod_{i \in I} X_i, (x_0^i)_{i \in I}) & \xrightarrow{\text{pr}_i} & (X_i, x_0^i) \end{array}$$

in \mathbf{Sets}_* . Then there exists a unique morphism of pointed sets

$$\phi: (P, *) \rightarrow \left(\prod_{i \in I} X_i, (x_0^i)_{i \in I} \right)$$

making the diagram

$$\begin{array}{ccc} (P, *) & & \\ \downarrow \phi \exists! & \searrow p_i & \\ (\prod_{i \in I} X_i, (x_0^i)_{i \in I}) & \xrightarrow{\text{pr}_i} & (X_i, x_0^i) \end{array}$$

commute, being uniquely determined by the condition $\text{pr}_i \circ \phi = p_i$ for each $i \in I$ via

$$\phi(x) = (p_i(x))_{i \in I}$$

for each $x \in P$. Note that this is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \phi(*) &= (p_i(*))_{i \in I} \\ &= (x_0^i)_{i \in I}, \end{aligned}$$

where we have used that p_i is a morphism of pointed sets for each $i \in I$. \square

00A7

PROPOSITION 3.2.2.3 ► PROPERTIES OF PRODUCTS OF FAMILIES OF POINTED SETS

Let $\{(X_i, x_0^i)\}_{i \in I}$ be a family of pointed sets.

00A8

1. *Functoriality.* The assignment $\{(X_i, x_0^i)\}_{i \in I} \mapsto (\prod_{i \in I} X_i, (x_0^i)_{i \in I})$ de-

defines a functor

$$\prod_{i \in I}: \text{Fun}(I_{\text{disc}}, \text{Sets}_*) \rightarrow \text{Sets}_*.$$

PROOF 3.2.2.4 ► PROOF OF PROPOSITION 3.2.2.3

Item 1: Functoriality

This follows from ?? of ??.

00A9 **3.2.3 Products**

Let (X, x_0) and (Y, y_0) be pointed sets.

00AA DEFINITION 3.2.3.1 ► PRODUCTS OF POINTED SETS

The **product** of (X, x_0) and (Y, y_0) is the pair consisting of:

- *The Limit.* The pointed set $(X \times Y, (x_0, y_0))$.
- *The Cone.* The morphisms of pointed sets

$$\text{pr}_1: (X \times Y, (x_0, y_0)) \rightarrow (X, x_0),$$

$$\text{pr}_2: (X \times Y, (x_0, y_0)) \rightarrow (Y, y_0)$$

defined by

$$\text{pr}_1(x, y) \stackrel{\text{def}}{=} x,$$

$$\text{pr}_2(x, y) \stackrel{\text{def}}{=} y$$

for each $(x, y) \in X \times Y$.

PROOF 3.2.3.2 ► PROOF OF DEFINITION 3.2.3.1

We claim that $(X \times Y, (x_0, y_0))$ is the categorical product of (X, x_0) and (Y, y_0) in Sets_* . Indeed, suppose we have a diagram of the form

$$\begin{array}{ccc} & (P, *) & \\ p_1 \swarrow & & \searrow p_2 \\ (X, x_0) & \xleftarrow{\text{pr}_1} (X \times Y, (x_0, y_0)) \xrightarrow{\text{pr}_2} & (Y, y_0) \end{array}$$

in \mathbf{Sets}_* . Then there exists a unique morphism of pointed sets

$$\phi: (P, *) \rightarrow (X \times Y, (x_0, y_0))$$

making the diagram

$$\begin{array}{ccccc} & & (P, *) & & \\ & \swarrow p_1 & \downarrow \phi \exists! & \searrow p_2 & \\ (X, x_0) & \xleftarrow{\text{pr}_1} & (X \times Y, (x_0, y_0)) & \xrightarrow{\text{pr}_2} & (Y, y_0) \end{array}$$

commute, being uniquely determined by the conditions

$$\text{pr}_1 \circ \phi = p_1,$$


$$\text{pr}_2 \circ \phi = p_2$$

via

$$\phi(x) = (p_1(x), p_2(x))$$

for each $x \in P$. Note that this is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \phi(*) &= (p_1(*), p_2(*)) \\ &= (x_0, y_0), \end{aligned}$$

where we have used that p_1 and p_2 are morphisms of pointed sets. 

00AB

PROPOSITION 3.2.3.3 ► PROPERTIES OF PRODUCTS OF POINTED SETS

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets.

00AC

1. *Functoriality.* The assignments

$$(X, x_0), (Y, y_0), ((X, x_0), (Y, y_0)) \mapsto (X \times Y, (x_0, y_0))$$

define functors

$$X \times -: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

$$- \times Y: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

$$-_1 \times -_2: \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

defined in the same way as the functors of [Item 1 of Proposition 2.1.3.3](#).

00AD

2. *Associativity.* We have an isomorphism of pointed sets

$$((X \times Y) \times Z, ((x_0, y_0), z_0)) \cong (X \times (Y \times Z), (x_0, (y_0, z_0)))$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$.

00AE

3. *Unitality.* We have isomorphisms of pointed sets

$$(\text{pt}, \star) \times (X, x_0) \cong (X, x_0),$$

$$(X, x_0) \times (\text{pt}, \star) \cong (X, x_0),$$

natural in $(X, x_0) \in \text{Obj}(\text{Sets}_*)$.

00AF

4. *Commutativity.* We have an isomorphism of pointed sets

$$(X \times Y, (x_0, y_0)) \cong (Y \times X, (y_0, x_0)),$$

natural in $(X, x_0), (Y, y_0) \in \text{Obj}(\text{Sets}_*)$.

00AG

5. *Symmetric Monoidality.* The triple $(\text{Sets}_*, \times, (\text{pt}, \star))$ is a symmetric monoidal category.

PROOF 3.2.3.4 ► PROOF OF PROPOSITION 3.2.3.3

Item 1: Functoriality

This is a special case of functoriality of limits, ?? of ??.

Item 2: Associativity

This follows from **Item 3** of **Proposition 2.1.3.3**.

Item 3: Unitality

This follows from **Item 4** of **Proposition 2.1.3.3**.

Item 4: Commutativity

This follows from **Item 5** of **Proposition 2.1.3.3**.

Item 5: Symmetric Monoidality

This follows from **Item 12** of **Proposition 2.1.3.3**. 

00AH 3.2.4 Pullbacks

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets and let $f: (X, x_0) \rightarrow (Z, z_0)$ and $g: (Y, y_0) \rightarrow (Z, z_0)$ be morphisms of pointed sets.

00AJ

DEFINITION 3.2.4.1 ► PULLBACKS OF POINTED SETS

The **pullback of** (X, x_0) **and** (Y, y_0) **over** (Z, z_0) **along** (f, g) is the pair consisting of:

- *The Limit.* The pointed set $(X \times_Z Y, (x_0, y_0))$.
- *The Cone.* The morphisms of pointed sets

$$\text{pr}_1: (X \times_Z Y, (x_0, y_0)) \rightarrow (X, x_0),$$

$$\text{pr}_2: (X \times_Z Y, (x_0, y_0)) \rightarrow (Y, y_0)$$

defined by

$$\text{pr}_1(x, y) \stackrel{\text{def}}{=} x,$$

$$\text{pr}_2(x, y) \stackrel{\text{def}}{=} y$$

for each $(x, y) \in X \times_Z Y$.

PROOF 3.2.4.2 ► PROOF OF DEFINITION 3.2.4.1

We claim that $X \times_Z Y$ is the categorical pullback of (X, x_0) and (Y, y_0) over (Z, z_0) with respect to (f, g) in Sets_* . First we need to check that the relevant pullback diagram commutes, i.e. that we have


$$f \circ \text{pr}_1 = g \circ \text{pr}_2, \quad \begin{array}{ccc} (X \times_Z Y, (x_0, y_0)) & \xrightarrow{\text{pr}_2} & (Y, y_0) \\ \text{pr}_1 \downarrow & & \downarrow g \\ (X, x_0) & \xrightarrow{f} & (Z, z_0). \end{array}$$

Indeed, given $(x, y) \in X \times_Z Y$, we have

$$\begin{aligned} [f \circ \text{pr}_1](x, y) &= f(\text{pr}_1(x, y)) \\ &= f(x) \\ &= g(y) \\ &= g(\text{pr}_2(x, y)) \\ &= [g \circ \text{pr}_2](x, y), \end{aligned}$$

where $f(x) = g(y)$ since $(x, y) \in X \times_Z Y$. Next, we prove that $X \times_Z Y$ satisfies the universal property of the pullback. Suppose we have a diagram


of the form



in Sets_* . Then there exists a unique morphism of pointed sets

$$\phi: (P, *) \rightarrow (X \times_Z Y, (x_0, y_0))$$

making the diagram



commute, being uniquely determined by the conditions

$$\begin{aligned} \text{pr}_1 \circ \phi &= p_1, \\ \text{pr}_2 \circ \phi &= p_2 \end{aligned}$$

via

$$\phi(x) = (p_1(x), p_2(x))$$

for each $x \in P$, where we note that $(p_1(x), p_2(x)) \in X \times Y$ indeed lies in $X \times_Z Y$ by the condition

$$f \circ p_1 = g \circ p_2,$$

which gives

$$f(p_1(x)) = g(p_2(x))$$

for each $x \in P$, so that $(p_1(x), p_2(x)) \in X \times_Z Y$. Lastly, we note that ϕ is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \phi(*) &= (p_1(*), p_2(*)) \\ &= (x_0, y_0), \end{aligned}$$

where we have used that p_1 and p_2 are morphisms of pointed sets. ▢

00AK

PROPOSITION 3.2.4.3 ► PROPERTIES OF PULLBACKS OF POINTED SETS


Let $(X, x_0), (Y, y_0), (Z, z_0)$, and (A, a_0) be pointed sets.

00AL


1. *Functoriality.* The assignment $(X, Y, Z, f, g) \mapsto X \times_{f,Z,g} Y$ defines a functor

$$-_1 \times_{-3} -_1 : \text{Fun}(\mathcal{P}, \text{Sets}_*) \rightarrow \text{Sets}_*,$$

where \mathcal{P} is the category that looks like this:



In particular, the action on morphisms of $-_1 \times_{-3} -_1$ is given by sending a morphism



in $\text{Fun}(\mathcal{P}, \text{Sets}_*)$ to the morphism of pointed sets


$$\xi : (X \times_Z Y, (x_0, y_0)) \xrightarrow{\exists!} (X' \times_{Z'} Y', (x'_0, y'_0))$$

given by

$$\xi(x, y) \stackrel{\text{def}}{=} (\phi(x), \psi(y))$$

for each $(x, y) \in X \times_Z Y$, which is the unique morphism of pointed


sets making the diagram



commute.

00AM


2. *Associativity.* Given a diagram



in \mathbf{Sets}_* , we have isomorphisms of pointed sets

$$(X \times_W Y) \times_V Z \cong (X \times_W Y) \times_Y (Y \times_V Z) \cong X \times_W (Y \times_V Z),$$

where these pullbacks are built as in the diagrams



00AN

3. *Unitality.* We have isomorphisms of pointed sets

$$\begin{array}{ccc} A & \xlongequal{\quad} & A \\ \downarrow f & \lrcorner & \downarrow f \\ X & \xlongequal{\quad} & X \end{array} \quad \begin{array}{l} X \times_X A \cong A, \\ A \times_X X \cong A, \end{array} \quad \begin{array}{ccc} A & \xrightarrow{f} & X \\ \parallel & \lrcorner & \parallel \\ X & \xrightarrow{f} & X. \end{array}$$

00AP

4. *Commutativity.* We have an isomorphism of pointed sets

$$\begin{array}{ccc}
 A \times_X B & \longrightarrow & B \\
 \downarrow \lrcorner & & \downarrow g \\
 A & \xrightarrow{f} & X,
 \end{array}
 \quad A \times_X B \cong B \times_X A
 \quad
 \begin{array}{ccc}
 B \times_X A & \longrightarrow & A \\
 \downarrow \lrcorner & & \downarrow f \\
 B & \xrightarrow{g} & X.
 \end{array}$$

00AQ

5. *Interaction With Products.* We have an isomorphism of pointed sets

$$\begin{array}{ccc}
 X \times Y & \longrightarrow & Y \\
 \downarrow \lrcorner & & \downarrow !_Y \\
 X & \xrightarrow{!_X} & \text{pt.}
 \end{array}
 \quad X \times_{\text{pt}} Y \cong X \times Y,$$

00AR

6. *Symmetric Monoidality.* The triple $(\text{Sets}_*, \times_X, X)$ is a symmetric monoidal category.

PROOF 3.2.4.4 ► PROOF OF PROPOSITION 3.2.4.3

Item 1: Functoriality

This is a special case of functoriality of co/limits, ?? of ??, with the explicit expression for ξ following from the commutativity of the cube pullback diagram.

Item 2: Associativity

This follows from **Item 2** of **Proposition 3.2.4.3**.

Item 3: Unitality

This follows from **Item 3** of **Proposition 2.1.4.5**.

Item 4: Commutativity

This follows from **Item 4** of **Proposition 2.1.4.5**.

Item 5: Interaction With Products

This follows from **Item 6** of **Proposition 2.1.4.5**.

Item 6: Symmetric Monoidality

This follows from **Item 7** of **Proposition 2.1.4.5**. 

00AS 3.2.5 Equalisers

Let $f, g: (X, x_0) \rightrightarrows (Y, y_0)$ be morphisms of pointed sets.

00AT

DEFINITION 3.2.5.1 ► EQUALISERS OF POINTED SETS

The **equaliser of** (f, g) is the pair consisting of:

- *The Limit.* The pointed set $(\text{Eq}(f, g), x_0)$.
- *The Cone.* The morphism of pointed sets

$$\text{eq}(f, g): (\text{Eq}(f, g), x_0) \hookrightarrow (X, x_0)$$

given by the canonical inclusion $\text{eq}(f, g) \hookrightarrow \text{Eq}(f, g) \hookrightarrow X$.

PROOF 3.2.5.2 ► PROOF OF DEFINITION 3.2.5.1

We claim that $(\text{Eq}(f, g), x_0)$ is the categorical equaliser of f and g in Sets_* . First we need to check that the relevant equaliser diagram commutes, i.e. that we have

$$f \circ \text{eq}(f, g) = g \circ \text{eq}(f, g),$$

which indeed holds by the definition of the set $\text{Eq}(f, g)$. Next, we prove that $\text{Eq}(f, g)$ satisfies the universal property of the equaliser. Suppose we have a diagram of the form

$$\begin{array}{ccc} (\text{Eq}(f, g), x_0) & \xrightarrow{\text{eq}(f, g)} & (X, x_0) \xrightarrow[f]{g} (Y, y_0) \\ & \nearrow e & \\ (E, *) & & \end{array}$$

in Sets_* . Then there exists a unique morphism of pointed sets

$$\phi: (E, *) \rightarrow (\text{Eq}(f, g), x_0)$$

making the diagram

$$\begin{array}{ccc} (\text{Eq}(f, g), x_0) & \xrightarrow{\text{eq}(f, g)} & (X, x_0) \xrightarrow[f]{g} (Y, y_0) \\ \uparrow \phi \exists! & \nearrow e & \\ (E, *) & & \end{array}$$

commute, being uniquely determined by the condition

$$\text{eq}(f, g) \circ \phi = e$$

via

$$\phi(x) = e(x)$$

for each $x \in E$, where we note that $e(x) \in A$ indeed lies in $\text{Eq}(f, g)$ by the condition


$$f \circ e = g \circ e,$$

which gives

$$f(e(x)) = g(e(x))$$

for each $x \in E$, so that $e(x) \in \text{Eq}(f, g)$. Lastly, we note that ϕ is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \phi(*) &= e(*) \\ &= x_0, \end{aligned}$$

where we have used that e is a morphism of pointed sets. 

00AU PROPOSITION 3.2.5.3 ► PROPERTIES OF EQUALISERS OF POINTED SETS

Let (X, x_0) and (Y, y_0) be pointed sets and let $f, g, h: (X, x_0) \rightarrow (Y, y_0)$ be morphisms of pointed sets.

00AV 1. *Associativity.* We have isomorphisms of pointed sets

$$\underbrace{\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h))}_{=\text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))} \cong \text{Eq}(f, g, h) \cong \underbrace{\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g))}_{=\text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))},$$

where $\text{Eq}(f, g, h)$ is the limit of the diagram

$$(X, x_0) \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{-g} \\ \xrightarrow{h} \end{array} (Y, y_0)$$

in Sets_* , being explicitly given by

$$\text{Eq}(f, g, h) \cong \{a \in A \mid f(a) = g(a) = h(a)\}.$$

00AW 2. *Unitality.* We have an isomorphism of pointed sets

$$\text{Eq}(f, f) \cong X.$$

00AX 3. *Commutativity.* We have an isomorphism of pointed sets

$$\text{Eq}(f, g) \cong \text{Eq}(g, f).$$

PROOF 3.2.5.4 ► PROOF OF PROPOSITION 3.2.5.3

Item 1: Associativity

This follows from **Item 1** of **Proposition 2.1.5.3**.

Item 2: Unitality

This follows from **Item 2** of **Proposition 2.1.5.3**.

Item 3: Commutativity

This follows from **Item 3** of **Proposition 2.1.5.3**. 00AY **3.3 Colimits of Pointed Sets**00AZ **3.3.1 The Initial Pointed Set**00B0 **DEFINITION 3.3.1.1 ► THE INITIAL POINTED SET**

The **initial pointed set** is the pair $\left((pt, \star), \{ \iota_X \}_{(X, x_0) \in \text{Obj}(\text{Sets}_*)} \right)$ consisting of:

- *The Limit*. The pointed set (pt, \star) .
- *The Cone*. The collection of morphisms of pointed sets

$$\{ \iota_X : (pt, \star) \rightarrow (X, x_0) \}_{(X, x_0) \in \text{Obj}(\text{Sets}_*)}$$

defined by

$$\iota_X(\star) \stackrel{\text{def}}{=} x_0.$$

PROOF 3.3.1.2 ► PROOF OF DEFINITION 3.3.1.1

We claim that (pt, \star) is the initial object of Sets_* . Indeed, suppose we have a diagram of the form


$$(pt, \star) \quad (X, x_0)$$

in Sets_* . Then there exists a unique morphism of pointed sets

$$\phi : (pt, \star) \rightarrow (X, x_0)$$

making the diagram

$$(pt, \star) \xrightarrow[\exists!]{\phi} (X, x_0)$$

commute, namely ι_X . 

00B1 **3.3.2 Coproducts of Families of Pointed Sets**

Let $\{(X_i, x_0^i)\}_{i \in I}$ be a family of pointed sets.

00B2 **DEFINITION 3.3.2.1 ► COPRODUCTS OF FAMILIES OF POINTED SETS**

The **coproduct of the family** $\{(X_i, x_0^i)\}_{i \in I}$, also called their **wedge sum**, is the pair consisting of:

• *The Colimit.* The pointed set $(\bigvee_{i \in I} X_i, p_0)$ consisting of:

– *The Underlying Set.* The set $\bigvee_{i \in I} X_i$ defined by

$$\bigvee_{i \in I} X_i \stackrel{\text{def}}{=} \left(\coprod_{i \in I} X_i \right) / \sim,$$

where \sim is the equivalence relation on $\coprod_{i \in I} X_i$ given by declaring

$$(i, x_0^i) \sim (j, x_0^j)$$

for each $i, j \in I$.

– *The Basepoint.* The element p_0 of $\bigvee_{i \in I} X_i$ defined by

$$\begin{aligned} p_0 &\stackrel{\text{def}}{=} [(i, x_0^i)] \\ &= [(j, x_0^j)] \end{aligned}$$

for any $i, j \in I$.

• *The Cocone.* The collection

$$\left\{ \text{inj}_i: (X_i, x_0^i) \rightarrow \left(\bigvee_{i \in I} X_i, p_0 \right) \right\}_{i \in I}$$

of morphism of pointed sets given by

$$\text{inj}_i(x) \stackrel{\text{def}}{=} (i, x)$$

for each $x \in X_i$ and each $i \in I$.

PROOF 3.3.2.2 ► PROOF OF DEFINITION 3.3.2.1

We claim that $(\bigvee_{i \in I} X_i, p_0)$ is the categorical coproduct of $\{(X_i, x_0^i)\}_{i \in I}$ in \mathbf{Sets}_* . Indeed, suppose we have, for each $i \in I$, a diagram of the form

$$\begin{array}{ccc} & & (C, *) \\ & \nearrow \iota_i & \\ (X_i, x_0^i) & \xrightarrow{\text{inj}_i} & \left(\bigvee_{i \in I} X_i, p_0 \right) \end{array}$$

in \mathbf{Sets}_* . Then there exists a unique morphism of pointed sets

$$\phi: \left(\bigvee_{i \in I} X_i, p_0 \right) \rightarrow (C, *)$$

making the diagram

$$\begin{array}{ccc} & & (C, *) \\ & \nearrow \iota_i & \uparrow \exists! \phi \\ (X_i, x_0^i) & \xrightarrow{\text{inj}_i} & \left(\bigvee_{i \in I} X_i, p_0 \right) \end{array}$$

commute, being uniquely determined by the condition $\phi \circ \text{inj}_i = \iota_i$ for each $i \in I$ via

$$\phi([(i, x)]) = \iota_i(x)$$

for each $[(i, x)] \in \bigvee_{i \in I} X_i$, where we note that ϕ is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \phi(p_0) &= \iota_i([(i, x_0^i)]) \\ &= *, \end{aligned}$$

as ι_i is a morphism of pointed sets. ▢

00B3 PROPOSITION 3.3.2.3 ► PROPERTIES OF COPRODUCTS OF FAMILIES OF POINTED SETS

Let $\{(X_i, x_0^i)\}_{i \in I}$ be a family of pointed sets.

00B4 1. *Functoriality.* The assignment $\{(X_i, x_0^i)\}_{i \in I} \mapsto (\bigvee_{i \in I} X_i, p_0)$ defines a

functor

$$\bigvee_{i \in I} : \text{Fun}(I_{\text{disc}}, \text{Sets}_*) \rightarrow \text{Sets}_*.$$

PROOF 3.3.2.4 ► PROOF OF PROPOSITION 3.3.2.3

Item 1: Functoriality

This follows from ?? of ??.

00B5 3.3.3 Coproducts

Let (X, x_0) and (Y, y_0) be pointed sets.

00B6 DEFINITION 3.3.3.1 ► COPRODUCTS OF POINTED SETS

The **coproduct** of (X, x_0) and (Y, y_0) , also called their **wedge sum**, is the pair consisting of:

- *The Colimit.* The pointed set $(X \vee Y, p_0)$ consisting of:

- *The Underlying Set.* The set $X \vee Y$ defined by

$$\begin{aligned} (X \vee Y, p_0) &\stackrel{\text{def}}{=} (X, x_0) \amalg (Y, y_0) & \begin{array}{ccc} X \vee Y & \longleftarrow & Y \\ \uparrow \lrcorner & & \uparrow [y_0] \\ X & \xleftarrow{[x_0]} & \text{pt} \end{array} \\ &\cong (X \amalg_{\text{pt}} Y, p_0) \\ &\cong (X \amalg Y / \sim, p_0), \end{aligned}$$

where \sim is the equivalence relation on $X \amalg Y$ obtained by declaring $(0, x_0) \sim (1, y_0)$.

- *The Basepoint.* The element p_0 of $X \vee Y$ defined by

$$\begin{aligned} p_0 &\stackrel{\text{def}}{=} [(0, x_0)] \\ &= [(1, y_0)]. \end{aligned}$$

- *The Cocone.* The morphisms of pointed sets

$$\begin{aligned} \text{inj}_1 &: (X, x_0) \rightarrow (X \vee Y, p_0), \\ \text{inj}_2 &: (Y, y_0) \rightarrow (X \vee Y, p_0), \end{aligned}$$

given by

$$\begin{aligned} \text{inj}_1(x) &\stackrel{\text{def}}{=} [(0, x)], \\ \text{inj}_2(y) &\stackrel{\text{def}}{=} [(1, y)], \end{aligned}$$

for each $x \in X$ and each $y \in Y$.

PROOF 3.3.3.2 ► PROOF OF DEFINITION 3.3.3.1

We claim that $(X \vee Y, p_0)$ is the categorical coproduct of (X, x_0) and (Y, y_0) in \mathbf{Sets}_* . Indeed, suppose we have a diagram of the form

$$\begin{array}{ccccc} & & (C, *) & & \\ & \nearrow \iota_X & & \nwarrow \iota_Y & \\ (X, x_0) & \xrightarrow{\text{inj}_X} & (X \vee Y, p_0) & \xleftarrow{\text{inj}_Y} & (Y, y_0) \end{array}$$

in \mathbf{Sets} . Then there exists a unique morphism of pointed sets

$$\phi: (X \vee Y, p_0) \rightarrow (C, *)$$

making the diagram

$$\begin{array}{ccccc} & & (C, *) & & \\ & \nearrow \iota_X & \uparrow \phi \exists! & \nwarrow \iota_Y & \\ (X, x_0) & \xrightarrow{\text{inj}_X} & (X \vee Y, p_0) & \xleftarrow{\text{inj}_Y} & (Y, y_0) \end{array}$$

commute, being uniquely determined by the conditions

$$\phi \circ \text{inj}_X = \iota_X,$$

$$\phi \circ \text{inj}_Y = \iota_Y$$

via

$$\phi(z) = \begin{cases} \iota_X(x) & \text{if } z = [(0, x)] \text{ with } x \in X, \\ \iota_Y(y) & \text{if } z = [(1, y)] \text{ with } y \in Y \end{cases}$$

for each $z \in X \vee Y$, where we note that ϕ is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \phi(p_0) &= \iota_X([(0, x_0)]) \\ &= \iota_Y([(1, y_0)]) \\ &= *, \end{aligned}$$

as ι_X and ι_Y are morphisms of pointed sets. ▢

00B7 **PROPOSITION 3.3.3.3 ► PROPERTIES OF WEDGE SUMS OF POINTED SETS**

Let (X, x_0) and (Y, y_0) be pointed sets.

00B8 1. *Functoriality.* The assignments

$$(X, x_0), (Y, y_0), ((X, x_0), (Y, y_0)) \mapsto (X \vee Y, p_0)$$

define functors

$$\begin{aligned} X \vee - &: \text{Sets}_* \rightarrow \text{Sets}_*, \\ - \vee Y &: \text{Sets}_* \rightarrow \text{Sets}_*, \\ -_1 \vee -_2 &: \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*. \end{aligned}$$

00B9 2. *Associativity.* We have an isomorphism of pointed sets

$$(X \vee Y) \vee Z \cong X \vee (Y \vee Z),$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Sets}_*$.

00BA 3. *Unitality.* We have isomorphisms of pointed sets

$$\begin{aligned} (\text{pt}, *) \vee (X, x_0) &\cong (X, x_0), \\ (X, x_0) \vee (\text{pt}, *) &\cong (X, x_0), \end{aligned}$$

natural in $(X, x_0) \in \text{Sets}_*$.

00BB 4. *Commutativity.* We have an isomorphism of pointed sets

$$X \vee Y \cong Y \vee X,$$

natural in $(X, x_0), (Y, y_0) \in \text{Sets}_*$.

00BC 5. *Symmetric Monoidality.* The triple $(\text{Sets}_*, \vee, \text{pt})$ is a symmetric monoidal category.

00BD 6. *The Fold Map.* We have a natural transformation

$$\nabla: \vee \circ \Delta_{\text{Sets}_*}^{\text{Cats}} \Longrightarrow \text{id}_{\text{Sets}_*},$$

called the **fold map**, whose component

$$\nabla_X: X \vee X \rightarrow X$$

at X is given by

$$\nabla_X(p) \stackrel{\text{def}}{=} \begin{cases} x & \text{if } p = [(0, x)], \\ x & \text{if } p = [(1, x)] \end{cases}$$

for each $p \in X \vee X$.

PROOF 3.3.3.4 ► PROOF OF PROPOSITION 3.3.3.3

Item 1: Functoriality

This follows from ?? of ??.

Item 2: Associativity

Clear.

Item 3: Unitality

Clear.

Item 4: Commutativity

Clear.

Item 5: Symmetric Monoidality

Omitted.

Item 6: The Fold Map

Naturality for the transformation ∇ is the statement that, given a morphism of pointed sets $f: (X, x_0) \rightarrow (Y, y_0)$, we have

$$\nabla_Y \circ (f \vee f) = f \circ \nabla_X,$$

$$\begin{array}{ccc} X \vee X & \xrightarrow{\nabla_X} & X \\ f \vee f \downarrow & & \downarrow f \\ Y \vee Y & \xrightarrow{\nabla_Y} & Y. \end{array}$$

Indeed, we have

$$\begin{aligned} [\nabla_Y \circ (f \vee f)]([(i, x)]) &= \nabla_Y([(i, f(x))]) \\ &= f(x) \\ &= f(\nabla_X([(i, x)])) \\ &= [f \circ \nabla_X]([(i, x)]) \end{aligned}$$

for each $[(i, x)] \in X \vee X$, and thus ∇ is indeed a natural transformation. \square

00BE 3.3.4 Pushouts

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets and let $f: (Z, z_0) \rightarrow (X, x_0)$ and $g: (Z, z_0) \rightarrow (Y, y_0)$ be morphisms of pointed sets.

00BF DEFINITION 3.3.4.1 ► PUSHOUTS OF POINTED SETS

The **pushout of (X, x_0) and (Y, y_0) over (Z, z_0) along (f, g)** is the pair consisting of:

- *The Colimit.* The pointed set $(X \amalg_{f,Z,g} Y, p_0)$, where:
 - The set $X \amalg_{f,Z,g} Y$ is the pushout (of unpointed sets) of X and Y over Z with respect to f and g ;
 - We have $p_0 = [x_0] = [y_0]$.
- *The Cocone.* The morphisms of pointed sets

$$\text{inj}_1: (X, x_0) \rightarrow (X \amalg_Z Y, p_0),$$

$$\text{inj}_2: (Y, y_0) \rightarrow (X \amalg_Z Y, p_0)$$

given by

$$\text{inj}_1(x) \stackrel{\text{def}}{=} [(0, x)]$$

$$\text{inj}_2(y) \stackrel{\text{def}}{=} [(1, y)]$$

for each $x \in X$ and each $y \in Y$.

PROOF 3.3.4.2 ► PROOF OF DEFINITION 3.3.4.1

Firstly, we note that indeed $[x_0] = [y_0]$, as we have

$$\begin{aligned} x_0 &= f(z_0), \\ y_0 &= g(z_0) \end{aligned}$$

since f and g are morphisms of pointed sets, with the relation \sim on $X \amalg_Z Y$ then identifying $x_0 = f(z_0) \sim g(z_0) = y_0$.

We now claim that $(X \amalg_Z Y, p_0)$ is the categorical pushout of (X, x_0) and (Y, y_0) over (Z, z_0) with respect to (f, g) in \mathbf{Sets}_* . First we need to check that the relevant pushout diagram commutes, i.e. that we have


$$\text{inj}_1 \circ f = \text{inj}_2 \circ g,$$

$$\begin{array}{ccc} (X \amalg_Z Y, p_0) & \xleftarrow{\text{inj}_2} & (Y, y_0) \\ \text{inj}_1 \uparrow & & \uparrow g \\ (X, x_0) & \xleftarrow{f} & (Z, z_0). \end{array}$$

Indeed, given $z \in Z$, we have

$$\begin{aligned} [\text{inj}_1 \circ f](z) &= \text{inj}_1(f(z)) \\ &= [(0, f(z))] \\ &= [(1, g(z))] \\ &= \text{inj}_2(g(z)) \\ &= [\text{inj}_2 \circ g](z), \end{aligned}$$

where $[(0, f(z))] = [(1, g(z))]$ by the definition of the relation \sim on $X \amalg Y$ (the coproduct of unpointed sets of X and Y). Next, we prove that $X \amalg_Z Y$ satisfies the universal property of the pushout. Suppose we have a diagram of the form



in \mathbf{Sets}_* . Then there exists a unique morphism of pointed sets

$$\phi: (X \amalg_Z Y, p_0) \rightarrow (P, *)$$

making the diagram

$$\begin{array}{ccc}
 (P, *) & & \\
 \uparrow \phi & \xleftarrow{\exists!} & \\
 (X \amalg_Z Y, p_0) & \xleftarrow{\text{inj}_2} & (Y, y_0) \\
 \uparrow \text{inj}_1 & \lrcorner & \uparrow g \\
 (X, x_0) & \xleftarrow{f} & (Z, z_0)
 \end{array}$$

ι_1 (curved arrow from (X, x_0) to $(P, *)$)
 ι_2 (curved arrow from (Y, y_0) to $(P, *)$)

commute, being uniquely determined by the conditions

$$\phi \circ \text{inj}_1 = \iota_1,$$

$$\phi \circ \text{inj}_2 = \iota_2$$

via

$$\phi(p) = \begin{cases} \iota_1(x) & \text{if } x = [(0, x)], \\ \iota_2(y) & \text{if } x = [(1, y)] \end{cases}$$

for each $p \in X \amalg_Z Y$, where the well-definedness of ϕ is proven in the same way as in the proof of [Definition 2.2.4.1](#). Finally, we show that ϕ is indeed a morphism of pointed sets, as we have

$$\begin{aligned}
 \phi(p_0) &= \phi([(0, x_0)]) \\
 &= \iota_1(x_0) \\
 &= *,
 \end{aligned}$$

or alternatively

$$\begin{aligned}
 \phi(p_0) &= \phi([(1, y_0)]) \\
 &= \iota_2(y_0) \\
 &= *,
 \end{aligned}$$

where we use that ι_1 (resp. ι_2) is a morphism of pointed sets. ▢

00BG

PROPOSITION 3.3.4.3 ► PROPERTIES OF PUSHOUTS OF POINTED SETS


Let $(X, x_0), (Y, y_0), (Z, z_0),$ and (A, a_0) be pointed sets.

00BH


1. *Functoriality.* The assignment $(X, Y, Z, f, g) \mapsto X \coprod_{f,Z,g} Y$ defines a functor

$$-_1 \coprod_{-3} -_1 : \text{Fun}(\mathcal{P}, \text{Sets}) \rightarrow \text{Sets}_*$$

where \mathcal{P} is the category that looks like this:



In particular, the action on morphisms of ${}_1 \coprod_{-3} -_1$ is given by sending a morphism



in $\text{Fun}(\mathcal{P}, \text{Sets}_*)$ to the morphism of pointed sets


$$\xi : (X \coprod_Z Y, p_0) \xrightarrow{\exists!} (X' \coprod_{Z'} Y', p'_0)$$

given by

$$\xi(p) \stackrel{\text{def}}{=} \begin{cases} \phi(x) & \text{if } p = [(0, x)], \\ \psi(y) & \text{if } p = [(1, y)] \end{cases}$$

for each $p \in X \coprod_Z Y$, which is the unique morphism of pointed sets


making the diagram



commute.

00BJ


2. *Associativity.* Given a diagram



in Sets, we have isomorphisms of pointed sets

$$(X \amalg_W Y) \amalg_V Z \cong (X \amalg_W Y) \amalg_Y (Y \amalg_V Z) \cong X \amalg_W (Y \amalg_V Z),$$

where these pullbacks are built as in the diagrams



00BK

3. *Unitality.* We have isomorphisms of sets

$$\begin{array}{ccc} A & \xlongequal{\quad} & A \\ \uparrow f & \lrcorner & \uparrow f \\ X & \xlongequal{\quad} & X \end{array} \quad \begin{array}{l} X \amalg_X A \cong A, \\ A \amalg_X X \cong A, \end{array} \quad \begin{array}{ccc} A & \xleftarrow{f} & X \\ \parallel & \lrcorner & \parallel \\ X & \xleftarrow{f} & X. \end{array}$$

00BL

4. *Commutativity.* We have an isomorphism of sets

$$\begin{array}{ccc}
 X \amalg_Z Y \longleftarrow Y & & Y \amalg_Z X \longleftarrow X \\
 \uparrow \ulcorner & \uparrow g & \uparrow \ulcorner \\
 X \xleftarrow{f} Z & & Y \xleftarrow{g} Z.
 \end{array}
 \quad X \amalg_Z Y \cong Y \amalg_Z X$$

00BM

5. *Interaction With Coproducts.* We have

$$\begin{array}{ccc}
 X \vee Y \longleftarrow Y & & \\
 \uparrow \ulcorner & \uparrow [y_0] & \\
 X \xrightarrow{[x_0]} \text{pt.} & &
 \end{array}
 \quad X \amalg_{\text{pt.}} Y \cong X \vee Y,$$

00BN

6. *Symmetric Monoidality.* The triple $(\text{Sets}_*, \amalg_X, (X, x_0))$ is a symmetric monoidal category.

PROOF 3.3.4.4 ► PROOF OF PROPOSITION 3.3.4.3

Item 1: Functoriality

This is a special case of functoriality of co/limits, ?? of ??, with the explicit expression for ξ following from the commutativity of the cube pushout diagram.

Item 2: Associativity

This follows from [Item 2 of Proposition 2.2.4.6](#).

Item 3: Unitality

This follows from [Item 3 of Proposition 2.2.4.6](#).

Item 4: Commutativity

This follows from [Item 4 of Proposition 2.2.4.6](#).

Item 5: Interaction With Coproducts

Clear.

Item 6: Symmetric Monoidality

Omitted. 

00BP 3.3.5 Coequalisers

Let $f, g: (X, x_0) \rightrightarrows (Y, y_0)$ be morphisms of pointed sets.

00BQ

DEFINITION 3.3.5.1 ► COEQUALISERS OF POINTED SETS

The **coequaliser of** (f, g) is the pointed set $(\text{CoEq}(f, g), [y_0])$.

PROOF 3.3.5.2 ► PROOF OF DEFINITION 3.3.5.1

We claim that $(\text{CoEq}(f, g), [y_0])$ is the categorical coequaliser of f and g in Sets_* . First we need to check that the relevant coequaliser diagram commutes, i.e. that we have

$$\text{coeq}(f, g) \circ f = \text{coeq}(f, g) \circ g.$$

Indeed, we have

$$\begin{aligned} [\text{coeq}(f, g) \circ f](x) &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](f(x)) \\ &\stackrel{\text{def}}{=} [f(x)] \\ &= [g(x)] \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](g(x)) \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g) \circ g](x) \end{aligned}$$

for each $x \in X$. Next, we prove that $\text{CoEq}(f, g)$ satisfies the universal property of the coequaliser. Suppose we have a diagram of the form


$$\begin{array}{ccc} (X, x_0) & \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} & (Y, y_0) & \xrightarrow{\text{coeq}(f, g)} & (\text{CoEq}(f, g), [y_0]) \\ & & & \searrow c & \\ & & & & (C, *) \end{array}$$

in Sets . Then, since $c(f(a)) = c(g(a))$ for each $a \in A$, it follows from **Items 4 and 5 of Proposition 7.5.2.3** that there exists a unique map $\phi: \text{CoEq}(f, g) \xrightarrow{\exists!} C$ making the diagram

$$\begin{array}{ccc} (X, x_0) & \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} & (Y, y_0) & \xrightarrow{\text{coeq}(f, g)} & (\text{CoEq}(f, g), [y_0]) \\ & & & \searrow c & \downarrow \phi \exists! \\ & & & & (C, *) \end{array}$$

commute, where we note that ϕ is indeed a morphism of pointed sets since

$$\begin{aligned} \phi([y_0]) &= [\phi \circ \text{coeq}(f, g)]([y_0]) \\ &= c([y_0]) \\ &= *, \end{aligned}$$

where we have used that c is a morphism of pointed sets. 

00BR PROPOSITION 3.3.5.3 ► PROPERTIES OF COEQUALISERS OF POINTED SETS

Let (X, x_0) and (Y, y_0) be pointed sets and let $f, g, h: (X, x_0) \rightarrow (Y, y_0)$ be morphisms of pointed sets.

00BS 1. *Associativity.* We have isomorphisms of pointed sets

$$\underbrace{\text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h)}_{=\text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ h)} \cong \text{CoEq}(f, g, h) \cong \underbrace{\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ g)}_{=\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ h)}$$

where $\text{CoEq}(f, g, h)$ is the colimit of the diagram

$$(X, x_0) \begin{array}{c} \xrightarrow{f} \\ \xrightarrow[-g]{\rightarrow} \\ \xrightarrow{h} \end{array} (Y, y_0)$$

in Sets_* .

00BT 2. *Unitality.* We have an isomorphism of pointed sets

$$\text{CoEq}(f, f) \cong B.$$

00BU 3. *Commutativity.* We have an isomorphism of pointed sets

$$\text{CoEq}(f, g) \cong \text{CoEq}(g, f).$$

PROOF 3.3.5.4 ► PROOF OF PROPOSITION 3.3.5.3

Item 1: Associativity

This follows from **Item 1** of [Proposition 2.2.5.6](#).

Item 2: Unitality

This follows from **Item 2** of [Proposition 2.2.5.6](#).

Item 3: Commutativity

This follows from **Item 3** of [Proposition 2.2.5.6](#). 

00BV 3.4 Constructions With Pointed Sets

00BW 3.4.1 Free Pointed Sets

Let X be a set.

00BX

DEFINITION 3.4.1.1 ► FREE POINTED SETS

The **free pointed set on X** is the pointed set X^+ consisting of:

- *The Underlying Set.* The set X^+ defined by¹

$$\begin{aligned} X^+ &\stackrel{\text{def}}{=} X \amalg \text{pt} \\ &\stackrel{\text{def}}{=} X \amalg \{\star\}. \end{aligned}$$

- *The Basepoint.* The element \star of X^+ .

¹*Further Notation:* We sometimes write \star_X for the basepoint of X^+ for clarity when there are multiple free pointed sets involved in the current discussion.

00BY

PROPOSITION 3.4.1.2 ► PROPERTIES OF FREE POINTED SETS

Let X be a set.

00BZ

1. *Functoriality.* The assignment $X \mapsto X^+$ defines a functor

$$(-)^+ : \text{Sets} \rightarrow \text{Sets}_*,$$

where

- *Action on Objects.* For each $X \in \text{Obj}(\text{Sets})$, we have

$$[(-)^+](X) \stackrel{\text{def}}{=} X^+,$$

where X^+ is the pointed set of [Definition 3.4.1.1](#);

- *Action on Morphisms.* For each morphism $f: X \rightarrow Y$ of Sets , the image

$$f^+ : X^+ \rightarrow Y^+$$

of f by $(-)^+$ is the map of pointed sets defined by

$$f^+(x) \stackrel{\text{def}}{=} \begin{cases} f(x) & \text{if } x \in X, \\ \star_Y & \text{if } x = \star_X. \end{cases}$$

00C0

2. *Adjointness.* We have an adjunction

$$((-)^+ \dashv \text{忘}) : \text{Sets} \begin{array}{c} \xrightarrow{(-)^+} \\ \perp \\ \xleftarrow{\text{忘}} \end{array} \text{Sets}_*,$$

witnessed by a bijection of sets

$$\mathbf{Sets}_*((X^+, \star_X), (Y, y_0)) \cong \mathbf{Sets}(X, Y),$$

natural in $X \in \mathbf{Obj}(\mathbf{Sets})$ and $(Y, y_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$.

00C1

3. *Symmetric Strong Monoidality With Respect to Wedge Sums.* The free pointed set functor of **Item 1** has a symmetric strong monoidal structure

$$\left((-)^+, (-)^+, \amalg, (-)^+, \amalg \right) : (\mathbf{Sets}, \amalg, \emptyset) \rightarrow (\mathbf{Sets}_*, \vee, \text{pt}),$$

being equipped with isomorphisms of pointed sets

$$\begin{aligned} (-)_{X,Y}^+, \amalg : X^+ \vee Y^+ &\xrightarrow{\cong} (X \amalg Y)^+, \\ (-)_{\mathbb{1}}^+, \amalg : \text{pt} &\xrightarrow{\cong} \emptyset^+, \end{aligned}$$

natural in $X, Y \in \mathbf{Obj}(\mathbf{Sets})$.

00C2

4. *Symmetric Strong Monoidality With Respect to Smash Products.* The free pointed set functor of **Item 1** has a symmetric strong monoidal structure

$$\left((-)^+, (-)^+, \times, (-)^+, \times \right) : (\mathbf{Sets}, \times, \text{pt}) \rightarrow (\mathbf{Sets}_*, \wedge, S^0),$$

being equipped with isomorphisms of pointed sets

$$\begin{aligned} (-)_{X,Y}^+, \times : X^+ \wedge Y^+ &\xrightarrow{\cong} (X \times Y)^+, \\ (-)_{\mathbb{1}}^+, \times : S^0 &\xrightarrow{\cong} \text{pt}^+, \end{aligned}$$

natural in $X, Y \in \mathbf{Obj}(\mathbf{Sets})$.

PROOF 3.4.1.3 ► PROOF OF PROPOSITION 3.4.1.2

Item 1: Functoriality

Clear.

Item 2: Adjointness

We claim there's an adjunction $(-)^+ \dashv \overline{\mathbf{Sets}}$, witnessed by a bijection of sets

$$\mathbf{Sets}_*((X^+, \star_X), (Y, y_0)) \cong \mathbf{Sets}(X, Y),$$

natural in $X \in \mathbf{Obj}(\mathbf{Sets})$ and $(Y, y_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$.

- *Map I.* We define a map

$$\Phi_{X,Y}: \text{Sets}_*((X^+, \star_X), (Y, y_0)) \rightarrow \text{Sets}(X, Y)$$

by sending a pointed function

$$\xi: (X^+, \star_X) \rightarrow (Y, y_0)$$

to the function

$$\xi^\dagger: X \rightarrow Y$$

given by

$$\xi^\dagger(x) \stackrel{\text{def}}{=} \xi(x)$$

for each $x \in X$.

- *Map II.* We define a map

$$\Psi_{X,Y}: \text{Sets}(X, Y) \rightarrow \text{Sets}_*((X^+, \star_X), (Y, y_0))$$

given by sending a function $\xi: X \rightarrow Y$ to the pointed function

$$\xi^\dagger: (X^+, \star_X) \rightarrow (Y, y_0)$$

defined by

$$\xi^\dagger(x) \stackrel{\text{def}}{=} \begin{cases} \xi(x) & \text{if } x \in X, \\ y_0 & \text{if } x = \star_X \end{cases}$$

for each $x \in X^+$.

- *Invertibility I.* We claim that

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{Sets}_*((X^+, \star_X), (Y, y_0))},$$

which is clear.

- *Invertibility II.* We claim that

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Sets}(X, Y)},$$

which is clear.

- *Naturality for Φ , Part I.* We need to show that, given a pointed function $g: (Y, y_0) \rightarrow (Y', y'_0)$, the diagram

$$\begin{array}{ccc} \text{Sets}_*((X^+, \star_X), (Y, y_0)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \\ g_* \downarrow & & \downarrow g_* \\ \text{Sets}_*((X^+, \star_X), (Y', y'_0)) & \xrightarrow{\Phi_{X,Y'}} & \text{Sets}(X, Y') \end{array}$$

commutes. Indeed, given a pointed function

$$\xi^\dagger: (X^+, \star_X) \rightarrow (Y, y_0)$$

we have

$$\begin{aligned} [\Phi_{X,Y'} \circ g_*](\xi) &= \Phi_{X,Y'}(g_*(\xi)) \\ &= \Phi_{X,Y'}(g \circ \xi) \\ &= g \circ \xi \\ &= g \circ \Phi_{X,Y'}(\xi) \\ &= g_*(\Phi_{X,Y'}(\xi)) \\ &= [g_* \circ \Phi_{X,Y'}](\xi). \end{aligned}$$

- *Naturality for Φ , Part II.* We need to show that, given a pointed function $f: (X, x_0) \rightarrow (X', x'_0)$, the diagram

$$\begin{array}{ccc} \text{Sets}_*((X'^+, \star_{X'}), (Y, y_0)) & \xrightarrow{\Phi_{X',Y}} & \text{Sets}(X', Y) \\ f_* \downarrow & & \downarrow f_* \\ \text{Sets}_*((X^+, \star_X), (Y, y_0)) & \xrightarrow{\Phi_{X,Y}} & \text{Sets}(X, Y) \end{array}$$

commutes. Indeed, given a function

$$\xi: X' \rightarrow Y,$$

we have

$$\begin{aligned}
 [\Phi_{X,Y} \circ f^*](\xi) &= \Phi_{X,Y}(f^*(\xi)) \\
 &= \Phi_{X,Y}(\xi \circ f) \\
 &= \xi \circ f \\
 &= \Phi_{X',Y}(\xi) \circ f \\
 &= f^*(\Phi_{X',Y}(\xi)) \\
 &= f^*(\Phi_{X',Y}(\xi)) \\
 &= [f^* \circ \Phi_{X',Y}](\xi).
 \end{aligned}$$

- *Naturality for Ψ .* Since Φ is natural in each argument and Φ is a componentwise inverse to Ψ in each argument, it follows from **Item 2 of Proposition 8.8.6.2** that Ψ is also natural in each argument.

Item 3: Symmetric Strong Monoidality With Respect to Wedge Sums

The isomorphism

$$\phi: X^+ \vee Y^+ \xrightarrow{\cong} (X \amalg Y)^+$$

is given by

$$\phi(z) = \begin{cases} x & \text{if } z = [(0, x)] \text{ with } x \in X, \\ y & \text{if } z = [(1, y)] \text{ with } y \in Y, \\ \star_X \amalg Y & \text{if } z = [(0, \star_X)], \\ \star_X \amalg Y & \text{if } z = [(1, \star_Y)] \end{cases}$$

for each $z \in X^+ \vee Y^+$, with inverse

$$\phi^{-1}: (X \amalg Y)^+ \xrightarrow{\cong} X^+ \vee Y^+$$

given by

$$\phi^{-1}(z) \stackrel{\text{def}}{=} \begin{cases} [(0, x)] & \text{if } z = [(0, x)], \\ [(1, y)] & \text{if } z = [(1, y)], \\ p_0 & \text{if } z = \star_X \amalg Y \end{cases}$$

for each $z \in (X \amalg Y)^+$.

Meanwhile, the isomorphism $\text{pt} \cong \emptyset^+$ is given by sending \star_X to \star_\emptyset .

That these isomorphisms satisfy the coherence conditions making the functor $(-)^+$ symmetric strong monoidal can be directly checked element by element.

Item 4: Symmetric Strong Monoidality With Respect to Smash Products

The isomorphism

$$\phi: X^+ \wedge Y^+ \xrightarrow{\cong} (X \times Y)^+$$

is given by

$$\phi(x \wedge y) = \begin{cases} (x, y) & \text{if } x \neq \star_X \text{ and } y \neq \star_Y \\ \star_{X \times Y} & \text{otherwise} \end{cases}$$

for each $x \wedge y \in X^+ \wedge Y^+$, with inverse

$$\phi^{-1}: (X \times Y)^+ \xrightarrow{\cong} X^+ \wedge Y^+$$

given by

$$\phi^{-1}(z) \stackrel{\text{def}}{=} \begin{cases} x \wedge y & \text{if } z = (x, y) \text{ with } (x, y) \in X \times Y, \\ \star_X \wedge \star_Y & \text{if } z = \star_{X \times Y}, \end{cases}$$

for each $z \in (X \amalg Y)^+$.

Meanwhile, the isomorphism $S^0 \cong \text{pt}^+$ is given by sending \star to $1 \in S^0 = \{0, 1\}$ and \star_{pt} to $0 \in S^0$.

That these isomorphisms satisfy the coherence conditions making the functor $(-)^+$ symmetric strong monoidal can be directly checked element by element.



Appendices

3.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets

Relations

5. Relations

6. Constructions With Relations

7. Equivalence Relations and Apartness Relations

Category Theory

8. Categories

Bicategories

9. Types of Morphisms in Bicategories

Chapter 4

Tensor Products of Pointed Sets

00C3 In this chapter we introduce, construct, and study tensor products of pointed sets. The most well-known among these is the *smash product of pointed sets*

$$\wedge : \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

introduced in [Section 4.5.1](#), defined via a universal property as inducing a bijection between the following data:

- Pointed maps $f : X \wedge Y \rightarrow Z$.
- Maps of sets $f : X \times Y \rightarrow Z$ satisfying

$$f(x_0, y) = z_0,$$

$$f(x, y_0) = z_0$$

for each $x \in X$ and each $y \in Y$.

As it turns out, however, dropping either of the *bilinearity* conditions

$$f(x_0, y) = z_0,$$

$$f(x, y_0) = z_0$$

while retaining the other leads to two other tensor products of pointed sets,

$$\triangleleft : \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

$$\triangleright : \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

called the *left* and *right tensor products of pointed sets*. In contrast to \wedge , which turns out to endow \mathbf{Sets}_* with a monoidal category structure ([Proposition 4.5.9.1](#)), these do not admit invertible associators and unitors, but do endow \mathbf{Sets}_* with the structure of a skew monoidal category, however ([Propositions 4.3.8.1](#) and [4.4.8.1](#)).

Finally, in addition to the tensor products \triangleleft , \triangleright , and \wedge , we also have a “tensor product” of the form

$$\odot : \mathbf{Sets} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

called the *tensor* of sets with pointed sets. All in all, these tensor products assemble into a family of functors of the form

$$\begin{aligned}\otimes_{k,\ell} &: \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \times \text{Mon}_{\mathbb{E}_\ell}(\text{Sets}) \rightarrow \text{Mon}_{\mathbb{E}_{k+\ell}}(\text{Sets}), \\ \triangleleft_{i,k} &: \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \times \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \rightarrow \text{Mon}_{\mathbb{E}_k}(\text{Sets}), \\ \triangleright_{i,k} &: \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \times \text{Mon}_{\mathbb{E}_k}(\text{Sets}) \rightarrow \text{Mon}_{\mathbb{E}_k}(\text{Sets}),\end{aligned}$$

where $k, \ell, i \in \mathbb{N}$ with $i \leq k - 1$. Together with the Cartesian product \times of Sets, the tensor products studied in this chapter form the cases:

- $(k, \ell) = (-1, -1)$ for the Cartesian product of Sets;
- $(k, \ell) = (0, -1)$ and $(-1, 0)$ for the tensor of sets with pointed sets of [Definition 4.2.1.1](#);
- $(i, k) = (-1, 0)$ for the left and right tensor products of pointed sets of [Sections 4.3](#) and [4.4](#);
- $(k, \ell) = (-1, -1)$ for the smash product of pointed sets of [Section 4.5](#).

In this chapter, we will carefully define and study bilinearity for pointed sets, as well as all the tensor products described above. Then, in [??](#), we will extend these to tensor products involving also monoids and commutative monoids, which will end up covering all cases up to $k, \ell \leq 2$, and hence *all* cases since \mathbb{E}_k -monoids on Sets are the same as \mathbb{E}_2 -monoids on Sets when $k \geq 2$.

Contents

4.1	Bilinear Morphisms of Pointed Sets	162
4.1.1	Left Bilinear Morphisms of Pointed Sets	162
4.1.2	Right Bilinear Morphisms of Pointed Sets	163
4.1.3	Bilinear Morphisms of Pointed Sets	164
4.2	Tensors and Cotensors of Pointed Sets by Sets	166
4.2.1	Tensors of Pointed Sets by Sets	166
4.2.2	Cotensors of Pointed Sets by Sets	174
4.3	The Left Tensor Product of Pointed Sets	182
4.3.1	Foundations	182
4.3.2	The Left Internal Hom of Pointed Sets	187
4.3.3	The Left Skew Unit	189
4.3.4	The Left Skew Associator	189
4.3.5	The Left Skew Left Unitor	192
4.3.6	The Left Skew Right Unitor	195
4.3.7	The Diagonal	197
4.3.8	The Left Skew Monoidal Structure on Pointed Sets	
	Associated to \triangleleft	199

4.3.9	Monoids With Respect to the Left Tensor Product of Pointed Sets	203
4.4	The Right Tensor Product of Pointed Sets	208
4.4.1	Foundations	208
4.4.2	The Right Internal Hom of Pointed Sets	212
4.4.3	The Right Skew Unit	215
4.4.4	The Right Skew Associator	215
4.4.5	The Right Skew Left Unitor	218
4.4.6	The Right Skew Right Unitor	220
4.4.7	The Diagonal	223
4.4.8	The Right Skew Monoidal Structure on Pointed Sets Associated to \triangleright	224
4.4.9	Monoids With Respect to the Right Tensor Product of Pointed Sets	229
4.5	The Smash Product of Pointed Sets	233
4.5.1	Foundations	233
4.5.2	The Internal Hom of Pointed Sets	243
4.5.3	The Monoidal Unit	247
4.5.4	The Associator	247
4.5.5	The Left Unitor	250
4.5.6	The Right Unitor	253
4.5.7	The Symmetry	256
4.5.8	The Diagonal	258
4.5.9	The Monoidal Structure on Pointed Sets Associated to \wedge	262
4.5.10	Universal Properties of the Smash Product of Pointed Sets I	267
4.5.11	Universal Properties of the Smash Product of Pointed Sets II	268
4.5.12	Monoids With Respect to the Smash Product of Pointed Sets	269
4.5.13	Comonoids With Respect to the Smash Product of Pointed Sets	269
4.6	Miscellany	270
4.6.1	The Smash Product of a Family of Pointed Sets	270
4.A	Other Chapters	270

00C4 4.1 Bilinear Morphisms of Pointed Sets

00C5 4.1.1 Left Bilinear Morphisms of Pointed Sets

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets.

00C6


DEFINITION 4.1.1.1 ▶ LEFT BILINEAR MORPHISMS OF POINTED SETS

A **left bilinear morphism of pointed sets from $(X \times Y, (x_0, y_0))$ to (Z, z_0)** is a map of sets

$$f: X \times Y \rightarrow Z$$

satisfying the following condition:^{1,2}

(★) *Left Unital Bilinearity.* The diagram



commutes, i.e. for each $y \in Y$, we have

$$f(x_0, y) = z_0.$$

¹Slogan: The map f is left bilinear if it preserves basepoints in its first argument.

²Succinctly, f is bilinear if we have

$$f(x_0, y) = z_0$$

for each $y \in Y$.

00C7

DEFINITION 4.1.1.2 ▶ THE SET OF LEFT BILINEAR MORPHISMS OF POINTED SETS

The **set of left bilinear morphisms of pointed sets from $(X \times Y, (x_0, y_0))$ to (Z, z_0)** is the set $\text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z)$ defined by

$$\text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \mid f \text{ is left bilinear}\}.$$

00C8 4.1.2 Right Bilinear Morphisms of Pointed Sets

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets.

00C9

DEFINITION 4.1.2.1 ▶ RIGHT BILINEAR MORPHISMS OF POINTED SETS

A **right bilinear morphism of pointed sets from $(X \times Y, (x_0, y_0))$ to (Z, z_0)** is a map of sets

$$f: X \times Y \rightarrow Z$$

satisfying the following condition:^{1,2}

(★) *Right Unital Bilinearity.* The diagram

$$\begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \epsilon_X \times \text{id}_{\text{pt}} \nearrow & & \dashrightarrow \sim \\
 X \times \text{pt} & & \text{pt} \\
 \text{id}_X \times [y_0] \searrow & & \downarrow [z_0] \\
 X \times Y & \xrightarrow{f} & Z
 \end{array}$$

commutes, i.e. for each $x \in X$, we have

$$f(x, y_0) = z_0.$$

¹*Slogan:* The map f is right bilinear if it preserves basepoints in its second argument.

²Succinctly, f is bilinear if we have

$$f(x, y_0) = z_0$$

for each $x \in X$.

00CA

DEFINITION 4.1.2.2 ► THE SET OF RIGHT BILINEAR MORPHISMS OF POINTED SETS

The **set of right bilinear morphisms of pointed sets from $(X \times Y, (x_0, y_0))$ to (Z, z_0)** is the set $\text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z)$ defined by

$$\text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \mid f \text{ is right bilinear}\}.$$

00CB 4.1.3 Bilinear Morphisms of Pointed Sets

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets.

00CC

DEFINITION 4.1.3.1 ► BILINEAR MORPHISMS OF POINTED SETS

A **bilinear morphism of pointed sets from $(X \times Y, (x_0, y_0))$ to (Z, z_0)** is a map of sets

$$f: X \times Y \rightarrow Z$$

that is both left bilinear and right bilinear.

∅∅CD


REMARK 4.1.3.2 ► UNWINDING DEFINITION 4.1.3.1

In detail, a **bilinear morphism of pointed sets from $(X \times Y, (x_0, y_0))$ to (Z, z_0)** is a map of sets

$$f: (X \times Y, (x_0, y_0)) \rightarrow (Z, z_0)$$

satisfying the following conditions:^{1,2}


1. *Left Unital Bilinearity.* The diagram



commutes, i.e. for each $y \in Y$, we have

$$f(x_0, y) = z_0.$$

2. *Right Unital Bilinearity.* The diagram



commutes, i.e. for each $x \in X$, we have

$$f(x, y_0) = z_0.$$

¹Slogan: The map f is bilinear if it preserves basepoints in each argument.

²Succinctly, f is bilinear if we have

$$f(x_0, y) = z_0,$$

$$f(x, y_0) = z_0$$

for each $x \in X$ and each $y \in Y$.

00CE

DEFINITION 4.1.3.3 ► THE SET OF BILINEAR MORPHISMS OF POINTED SETS

The **set of bilinear morphisms of pointed sets from** $(X \times Y, (x_0, y_0))$ **to** (Z, z_0) is the set $\text{Hom}_{\text{Sets}_*}^{\otimes}(X \times Y, Z)$ defined by

$$\text{Hom}_{\text{Sets}_*}^{\otimes}(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \mid f \text{ is bilinear}\}.$$

00CF

4.2 Tensors and Cotensors of Pointed Sets by Sets

00CG

4.2.1 Tensors of Pointed Sets by Sets

Let (X, x_0) be a pointed set and let A be a set.

00CH

DEFINITION 4.2.1.1 ► TENSORS OF POINTED SETS BY SETS

The **tensor of** (X, x_0) **by** A^1 is the pointed set² $A \odot (X, x_0)$ satisfying the following universal property:

(UP) We have a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}(A, \text{Sets}_*(X, K)),$$

natural in $(K, k_0) \in \text{Obj}(\text{Sets}_*)$.

¹Further Terminology: Also called the **copower of** (X, x_0) **by** A .

²Further Notation: Often written $A \odot X$ for simplicity.

00CJ

REMARK 4.2.1.2 ► UNWINDING DEFINITION 4.2.1.1

The universal property in Definition 4.2.1.1 is equivalent to the following one:

(UP) We have a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K),$$

natural in $(K, k_0) \in \text{Obj}(\text{Sets}_*)$, where $\text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K)$ is the set defined by

$$\text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K) \stackrel{\text{def}}{=} \left\{ f \in \text{Sets}(A \times X, K) \left| \begin{array}{l} \text{for each } a \in A, \text{ we} \\ \text{have } f(a, x_0) = k_0 \end{array} \right. \right\}.$$

PROOF 4.2.1.3 ► PROOF OF REMARK 4.2.1.2

We claim we have a bijection

$$\text{Sets}(A, \text{Sets}_*(X, K)) \cong \text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K)$$

natural in $(K, k_0) \in \text{Obj}(\text{Sets}_*)$. Indeed, this bijection is a restriction of the bijection

$$\text{Sets}(A, \text{Sets}(X, K)) \cong \text{Sets}(A \times X, K)$$

of [Item 2 of Proposition 2.1.3.3](#):

· A map

$$\begin{aligned} \xi: A &\longrightarrow \text{Sets}_*(X, K), \\ a &\longmapsto (\xi_a: X \rightarrow K), \end{aligned}$$

in $\text{Sets}(A, \text{Sets}_*(X, K))$ gets sent to the map

$$\xi^\dagger: A \times X \rightarrow K$$

defined by

$$\xi^\dagger(a, x) \stackrel{\text{def}}{=} \xi_a(x)$$

for each $(a, x) \in A \times X$, which indeed lies in $\text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K)$, as we have

$$\begin{aligned} \xi^\dagger(a, x_0) &\stackrel{\text{def}}{=} \xi_a(x_0) \\ &\stackrel{\text{def}}{=} k_0 \end{aligned}$$

for each $a \in A$, where we have used that $\xi_a \in \text{Sets}_*(X, K)$ is a morphism of pointed sets.

· Conversely, a map

$$\xi: A \times X \rightarrow K$$

in $\text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K)$ gets sent to the map

$$\begin{aligned} \xi^\dagger: A &\longrightarrow \text{Sets}_*(X, K), \\ a &\longmapsto (\xi_a^\dagger: X \rightarrow K), \end{aligned}$$

where


$$\xi_a^\dagger: X \rightarrow K$$

is the map defined by

$$\xi_a^\dagger(x) \stackrel{\text{def}}{=} \xi(a, x)$$

for each $x \in X$, and indeed lies in $\text{Sets}_*(X, K)$, as we have

$$\begin{aligned} \xi_a^\dagger(x_0) &\stackrel{\text{def}}{=} \xi(a, x_0) \\ &\stackrel{\text{def}}{=} k_0. \end{aligned}$$

This finishes the proof. 

00CK

CONSTRUCTION 4.2.1.4 ► CONSTRUCTION OF TENSORS OF POINTED SETS BY SETS

Concretely, the **tensor of** (X, x_0) **by** A is the pointed set $A \odot (X, x_0)$ consisting of:


- *The Underlying Set.* The set $A \odot X$ given by

$$A \odot X \cong \bigvee_{a \in A} (X, x_0),$$

where $\bigvee_{a \in A} (X, x_0)$ is the wedge product of the A -indexed family $((X, x_0))_{a \in A}$ of [Definition 3.3.2.1](#).

- *The Basepoint.* The point $[(a, x_0)] = [(a', x_0)]$ of $\bigvee_{a \in A} (X, x_0)$.

PROOF 4.2.1.5 ► PROOF OF CONSTRUCTION 4.2.1.4

(Proven below in a bit.) 

00CL

NOTATION 4.2.1.6 ► ELEMENTS OF TENSORS OF POINTED SETS BY SETS

We write $a \odot x$ for the element $[(a, x)]$ of

$$\begin{aligned} A \odot X &\cong \bigvee_{a \in A} (X, x_0) \\ &\stackrel{\text{def}}{=} \left(\prod_{i \in I} X_i \right) / \sim. \end{aligned}$$

00CM

REMARK 4.2.1.7 ► BASEPOINTS OF TENSORS OF POINTED SETS BY SETS

Taking the tensor of any element of A with the basepoint x_0 of X leads to the same element in $A \odot X$, i.e. we have

$$a \odot x_0 = a' \odot x_0,$$

for each $a, a' \in A$. This is due to the equivalence relation \sim on

$$\bigsqcup_{a \in A} (X, x_0) \stackrel{\text{def}}{=} \bigsqcup_{a \in A} X / \sim$$

identifying (a, x_0) with (a', x_0) , so that the equivalence class $a \odot x_0$ is independent from the choice of $a \in A$.

PROOF 4.2.1.8 ► PROOF OF CONSTRUCTION 4.2.1.4

We claim we have a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}(A, \text{Sets}_*(X, K))$$

natural in $(K, k_0) \in \text{Obj}(\text{Sets}_*)$.

• *Map I.* We define a map

$$\Phi_K : \text{Sets}_*(A \odot X, K) \rightarrow \text{Sets}(A, \text{Sets}_*(X, K))$$

by sending a morphism of pointed sets

$$\xi : (A \odot X, a \odot x_0) \rightarrow (K, k_0)$$

to the map of sets

$$\begin{aligned} \xi^\dagger : A &\rightarrow \text{Sets}_*(X, K), \\ a &\mapsto (\xi_a : X \rightarrow K), \end{aligned}$$

where

$$\xi_a : (X, x_0) \rightarrow (K, k_0)$$

is the morphism of pointed sets defined by

$$\xi_a(x) \stackrel{\text{def}}{=} \xi(a \odot x)$$

for each $x \in X$. Note that we have

$$\begin{aligned} \xi_a(x_0) &\stackrel{\text{def}}{=} \xi(a \odot x_0) \\ &= k_0, \end{aligned}$$

so that ξ_a is indeed a morphism of pointed sets, where we have used that ξ is a morphism of pointed sets.

• *Map II.* We define a map

$$\Psi_K : \text{Sets}(A, \text{Sets}_*(X, K)) \rightarrow \text{Sets}_*(A \odot X, K)$$

given by sending a map

$$\begin{aligned} \xi : A &\rightarrow \text{Sets}_*(X, K), \\ a &\mapsto (\xi_a : X \rightarrow K), \end{aligned}$$

to the morphism of pointed sets

$$\xi^\dagger : (A \odot X, a \odot x_0) \rightarrow (K, k_0)$$

defined by

$$\xi^\dagger(a \odot x) \stackrel{\text{def}}{=} \xi_a(x)$$

for each $a \odot x \in A \odot X$. Note that ξ^\dagger is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \xi^\dagger(a \odot x_0) &\stackrel{\text{def}}{=} \xi_a(x_0) \\ &= k_0, \end{aligned}$$

where we have used that $\xi(a) \in \text{Sets}_*(X, K)$ is a morphism of pointed sets.

• *Invertibility I.* We claim that

$$\Psi_K \circ \Phi_K = \text{id}_{\text{Sets}_*(A \odot X, K)}.$$

Indeed, given a morphism of pointed sets

$$\xi : (A \odot X, a \odot x_0) \rightarrow (K, k_0),$$

we have

$$\begin{aligned} [\Psi_K \circ \Phi_K](\xi) &= \Psi_K(\Phi_K(\xi)) \\ &= \Psi_K(\llbracket a \mapsto \llbracket x \mapsto \xi(a \odot x) \rrbracket \rrbracket) \\ &= \Psi_K(\llbracket a' \mapsto \llbracket x' \mapsto \xi(a' \odot x') \rrbracket \rrbracket) \\ &= \llbracket a \odot x \mapsto \text{ev}_x(\text{ev}_a(\llbracket a' \mapsto \llbracket x' \mapsto \xi(a' \odot x') \rrbracket \rrbracket)) \rrbracket \\ &= \llbracket a \odot x \mapsto \text{ev}_x(\llbracket x' \mapsto \xi(a \odot x') \rrbracket) \rrbracket \\ &= \llbracket a \odot x \mapsto \xi(a \odot x) \rrbracket \\ &= \xi. \end{aligned}$$

- *Invertibility II.* We claim that

$$\Phi_K \circ \Psi_K = \text{id}_{\text{Sets}(A, \text{Sets}_*(X, K))}.$$

Indeed, given a morphism $\xi: A \rightarrow \text{Sets}_*(X, K)$, we have

$$\begin{aligned} [\Phi_K \circ \Psi_K](\xi) &= \Phi_K(\Psi_K(\xi)) \\ &= \Phi_K(\llbracket a \odot x \mapsto \xi_a(x) \rrbracket) \\ &= \llbracket a \mapsto \llbracket x \mapsto \xi_a(x) \rrbracket \rrbracket \\ &= \llbracket a \mapsto \xi(a) \rrbracket \\ &= \xi. \end{aligned}$$

- *Naturality of Φ .* We need to show that, given a morphism of pointed sets

$$\phi: (K, k_0) \rightarrow (K', k'_0),$$

the diagram

$$\begin{array}{ccc} \text{Sets}_*(A \odot X, K) & \xrightarrow{\Phi_K} & \text{Sets}(A, \text{Sets}_*(X, K)) \\ \phi_* \downarrow & & \downarrow (\phi_*)_* \\ \text{Sets}_*(A \odot X, K') & \xrightarrow{\Phi_{K'}} & \text{Sets}(A, \text{Sets}_*(X, K')) \end{array}$$


commutes. Indeed, given a morphism of pointed sets

$$\xi: (A \odot X, a \odot x_0) \rightarrow (K, k_0),$$

we have

$$\begin{aligned} [\Phi_{K'} \circ \phi_*](\xi) &= \Phi_{K'}(\phi_*(\xi)) \\ &= \Phi_{K'}(\phi \circ \xi) \\ &= (\phi \circ \xi)^\dagger \\ &= \llbracket a \mapsto \phi \circ \xi(a \odot -) \rrbracket \\ &= \llbracket a \mapsto \phi_*(\xi(a \odot -)) \rrbracket \\ &= (\phi_*)_* (\llbracket a \mapsto \xi(a \odot -) \rrbracket) \\ &= (\phi_*)_* (\Phi_K(\xi)) \\ &= [(\phi_*)_* \circ \Phi_K](\xi). \end{aligned}$$

- *Naturality of Ψ .* Since Φ is natural and Φ is a componentwise inverse to Ψ , it follows from [Item 2 of Proposition 8.8.6.2](#) that Ψ is also natural.

This finishes the proof. 

00CN

PROPOSITION 4.2.1.9 ► PROPERTIES OF TENSORS OF POINTED SETS BY SETS

Let (X, x_0) be a pointed set and let A be a set.

00CP

1. *Functoriality.* The assignments $A, (X, x_0), (A, (X, x_0))$ define functors

$$\begin{aligned} A \odot - &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ - \odot X &: \mathbf{Sets} \rightarrow \mathbf{Sets}_*, \\ -_1 \odot -_2 &: \mathbf{Sets} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given:

- A map of sets $f: A \rightarrow B$;
- A pointed map $\phi: (X, x_0) \rightarrow (Y, y_0)$;

the induced map

$$f \odot \phi: A \odot X \rightarrow B \odot Y$$

is given by

$$[f \odot \phi](a \odot x) \stackrel{\text{def}}{=} f(a) \odot \phi(x)$$

for each $a \odot x \in A \odot X$.

00CQ

2. *Adjointness I.* We have an adjunction

$$(- \odot X \dashv \mathbf{Sets}_*(X, -)): \mathbf{Sets} \begin{array}{c} \xrightarrow{- \odot X} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*$$

witnessed by a bijection

$$\mathbf{Sets}_*(A \odot X, K) \cong \mathbf{Sets}(A, \mathbf{Sets}_*(X, K)),$$

natural in $A \in \text{Obj}(\mathbf{Sets})$ and $X, Y \in \text{Obj}(\mathbf{Sets}_*)$.

00CR

3. *Adjointness II.* We have an adjunctions

$$(A \odot - \dashv A \pitchfork -): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{A \odot -} \\ \perp \\ \xleftarrow{A \pitchfork -} \end{array} \mathbf{Sets}_*$$

witnessed by a bijection

$$\text{Hom}_{\mathbf{Sets}_*}(A \odot X, Y) \cong \text{Hom}_{\mathbf{Sets}_*}(X, A \pitchfork Y),$$

natural in $A \in \text{Obj}(\mathbf{Sets})$ and $X, Y \in \text{Obj}(\mathbf{Sets}_*)$.

00CS

4. *As a Weighted Colimit.* We have

$$A \odot X \cong \operatorname{colim}^{[A]}(X),$$

where in the right hand side we write:

- A for the functor $A: \mathbf{pt} \rightarrow \mathbf{Sets}$ picking $A \in \operatorname{Obj}(\mathbf{Sets})$;
- X for the functor $X: \mathbf{pt} \rightarrow \mathbf{Sets}_*$ picking $(X, x_0) \in \operatorname{Obj}(\mathbf{Sets}_*)$.

00CT

5. *Iterated Tensors.* We have an isomorphism of pointed sets

$$A \odot (B \odot X) \cong (A \times B) \odot X,$$

natural in $A, B \in \operatorname{Obj}(\mathbf{Sets})$ and $(X, x_0) \in \operatorname{Obj}(\mathbf{Sets}_*)$.

00CU

6. *Interaction With Homs.* We have a natural isomorphism

$$\mathbf{Sets}_*(A \odot X, -) \cong A \pitchfork \mathbf{Sets}_*(X, -).$$

00CV

7. *The Tensor Evaluation Map.* For each $X, Y \in \operatorname{Obj}(\mathbf{Sets}_*)$, we have a map

$$\operatorname{ev}_{X,Y}^\odot: \mathbf{Sets}_*(X, Y) \odot X \rightarrow Y,$$

natural in $X, Y \in \operatorname{Obj}(\mathbf{Sets}_*)$, and given by

$$\operatorname{ev}_{X,Y}^\odot(f \odot x) \stackrel{\text{def}}{=} f(x)$$

for each $f \odot x \in \mathbf{Sets}_*(X, Y) \odot X$.

00CW

8. *The Tensor Coevaluation Map.* For each $A \in \operatorname{Obj}(\mathbf{Sets})$ and each $X \in \operatorname{Obj}(\mathbf{Sets}_*)$, we have a map

$$\operatorname{coev}_{A,X}^\odot: A \rightarrow \mathbf{Sets}_*(X, A \odot X),$$

natural in $A \in \operatorname{Obj}(\mathbf{Sets})$ and $X \in \operatorname{Obj}(\mathbf{Sets}_*)$, and given by

$$\operatorname{coev}_{A,X}^\odot(a) \stackrel{\text{def}}{=} \llbracket x \mapsto a \odot x \rrbracket$$

for each $a \in A$.

PROOF 4.2.1.10 ► PROOF OF PROPOSITION 4.2.1.9

Item 1: Functoriality

This is the special case of ?? of ?? for when $C = \mathbf{Sets}_*$.

Item 2: Adjointness I

This is simply a rephrasing of [Definition 4.2.1.1](#).

Item 3: Adjointness II

This is the special case of ?? of ?? for when $C = \mathbf{Sets}_*$.

Item 4: As a Weighted Colimit

This is the special case of ?? of ?? for when $C = \mathbf{Sets}_*$.

Item 5: Iterated Tensors

This is the special case of ?? of ?? for when $C = \mathbf{Sets}_*$.


Item 6: Interaction With Homs

This is the special case of ?? of ?? for when $C = \mathbf{Sets}_*$.

Item 7: The Tensor Evaluation Map

This is the special case of ?? of ?? for when $C = \mathbf{Sets}_*$.

Item 8: The Tensor Coevaluation Map

This is the special case of ?? of ?? for when $C = \mathbf{Sets}_*$. 

00CX 4.2.2 Cotensors of Pointed Sets by Sets

Let (X, x_0) be a pointed set and let A be a set.

00CY DEFINITION 4.2.2.1 ► COTENSORS OF POINTED SETS BY SETS

The **cotensor of (X, x_0) by A** ¹ is the pointed set² $A \pitchfork (X, x_0)$ satisfying the following universal property:

(UP) We have a bijection

$$\mathbf{Sets}_*(K, A \pitchfork X) \cong \mathbf{Sets}(A, \mathbf{Sets}_*(K, X)),$$

natural in $(K, k_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$.

¹ *Further Terminology:* Also called the **power of (X, x_0) by A** .

² *Further Notation:* Often written $A \pitchfork X$ for simplicity.

00CZ

REMARK 4.2.2.2 ► UNWINDING DEFINITION 4.2.2.1

The universal property of [Definition 4.2.2.1](#) is equivalent to the following one:

(UP) We have a bijection

$$\text{Sets}_*(K, A \pitchfork X) \cong \text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times K, X),$$

natural in $(K, k_0) \in \text{Obj}(\text{Sets}_*)$, where $\text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times K, X)$ is the set defined by

$$\text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times K, X) \stackrel{\text{def}}{=} \left\{ f \in \text{Sets}(A \times K, X) \mid \begin{array}{l} \text{for each } a \in A, \text{ we} \\ \text{have } f(a, k_0) = x_0 \end{array} \right\}.$$

PROOF 4.2.2.3 ► PROOF OF REMARK 4.2.2.2

This follows from the bijection

$$\text{Sets}(A, \text{Sets}_*(K, X)) \cong \text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times K, X),$$

natural in $(K, k_0) \in \text{Obj}(\text{Sets}_*)$ constructed in the proof of [Remark 4.2.1.2](#). \square

00D0

CONSTRUCTION 4.2.2.4 ► CONSTRUCTION OF COTENSORS OF POINTED SETS BY SETS

Concretely, the **cotensor of (X, x_0) by A** is the pointed set $A \pitchfork (X, x_0)$ consisting of:

- *The Underlying Set.* The set $A \pitchfork X$ given by

$$A \pitchfork X \cong \bigwedge_{a \in A} (X, x_0),$$

where $\bigwedge_{a \in A} (X, x_0)$ is the smash product of the A -indexed family $((X, x_0))_{a \in A}$ of [Definition 4.6.1.1](#).

- *The Basepoint.* The point $[(x_0)_{a \in A}] = [(x_0, x_0, x_0, \dots)]$ of $\bigwedge_{a \in A} (X, x_0)$.

PROOF 4.2.2.5 ► PROOF OF CONSTRUCTION 4.2.2.4

We claim we have a bijection

$$\text{Sets}_*(K, A \pitchfork X) \cong \text{Sets}(A, \text{Sets}_*(K, X)),$$

natural in $(K, k_0) \in \text{Obj}(\text{Sets}_*)$.

• *Map 1.* We define a map

$$\Phi_K: \text{Sets}_*(K, A \pitchfork X) \rightarrow \text{Sets}(A, \text{Sets}_*(K, X)),$$

by sending a morphism of pointed sets

$$\xi: (K, k_0) \rightarrow (A \pitchfork X, [(x_0)_{a \in A}])$$

to the map of sets

$$\begin{aligned} \xi^\dagger: A &\longrightarrow \text{Sets}_*(K, X), \\ a &\longmapsto (\xi_a: K \rightarrow X), \end{aligned}$$

where

$$\xi_a: (K, k_0) \rightarrow (X, x_0)$$

is the morphism of pointed sets defined by

$$\xi_a(k) = \begin{cases} x_a^k & \text{if } \xi(k) \neq [(x_0)_{a \in A}], \\ x_0 & \text{if } \xi(k) = [(x_0)_{a \in A}] \end{cases}$$

for each $k \in K$, where x_a^k is the a th component of $\xi(k) = [(x_a^k)_{a \in A}]$. Note that:

1. The definition of $\xi_a(k)$ is independent of the choice of equivalence class. Indeed, suppose we have

$$\begin{aligned} \xi(k) &= \left[\left(x_a^k \right)_{a \in A} \right] \\ &= \left[\left(y_a^k \right)_{a \in A} \right] \end{aligned}$$

with $x_a^k \neq y_a^k$ for some $a \in A$. Then there exist $a_x, a_y \in A$ such that $x_{a_x}^k = y_{a_y}^k = x_0$. The equivalence relation \sim on $\prod_{a \in A} X$ then forces

$$\begin{aligned} \left[\left(x_a^k \right)_{a \in A} \right] &= [(x_0)_{a \in A}], \\ \left[\left(y_a^k \right)_{a \in A} \right] &= [(x_0)_{a \in A}], \end{aligned}$$

however, and $\xi_a(k)$ is defined to be x_0 in this case.

2. The map ξ_a is indeed a morphism of pointed sets, as we have

$$\xi_a(k_0) = x_0$$

since $\xi(k_0) = [(x_0)_{a \in A}]$ as ξ is a morphism of pointed sets and $\xi_a(k_0)$, defined to be the a th component of $[(x_0)_{a \in A}]$, is equal to x_0 .

• *Map II.* We define a map

$$\Psi_K: \text{Sets}(A, \text{Sets}_*(K, X)) \rightarrow \text{Sets}_*(K, A \pitchfork X),$$

given by sending a map

$$\begin{aligned} \xi: A &\rightarrow \text{Sets}_*(K, X), \\ a &\mapsto (\xi_a: K \rightarrow X), \end{aligned}$$

to the morphism of pointed sets

$$\xi^\dagger: (K, k_0) \rightarrow (A \pitchfork X, [(x_0)_{a \in A}])$$

defined by

$$\xi^\dagger(k) \stackrel{\text{def}}{=} [(\xi_a(k))_{a \in A}]$$

for each $k \in K$. Note that ξ^\dagger is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \xi^\dagger(k_0) &\stackrel{\text{def}}{=} [(\xi_a(k_0))_{a \in A}] \\ &= x_0, \end{aligned}$$

where we have used that $\xi_a \in \text{Sets}_*(K, X)$ is a morphism of pointed sets for each $a \in A$.

• *Naturality of Ψ .* We need to show that, given a morphism of pointed sets

$$\phi: (K, k_0) \rightarrow (K', k'_0),$$

the diagram

$$\begin{array}{ccc} \text{Sets}(A, \text{Sets}_*(K', X)) & \xrightarrow{\Psi_{K'}} & \text{Sets}_*(K', A \pitchfork X) \\ (\phi^*)_* \downarrow & & \downarrow \phi^* \\ \text{Sets}(A, \text{Sets}_*(K, X)) & \xrightarrow{\Psi_K} & \text{Sets}_*(K, A \pitchfork X) \end{array}$$

commutes. Indeed, given a map of sets

$$\begin{aligned}\xi: A &\longrightarrow \text{Sets}_*(K', X), \\ a &\longmapsto (\xi_a: K' \rightarrow X),\end{aligned}$$

we have

$$\begin{aligned}[\Psi_K \circ (\phi^*)_*](\xi) &= \Psi_K((\phi^*)_*(\xi)) \\ &= \Psi_K((\phi^*)_*(\llbracket a \mapsto \xi_a \rrbracket)) \\ &= \Psi_K(\llbracket a \mapsto \phi^*(\xi_a) \rrbracket) \\ &= \Psi_K(\llbracket a \mapsto \llbracket k \mapsto \xi_a(\phi(k)) \rrbracket \rrbracket) \\ &= \llbracket k \mapsto [(\xi_a(\phi(k)))_{a \in A}] \rrbracket \\ &= \phi^*(\llbracket k' \mapsto [(\xi_a(k'))_{a \in A}] \rrbracket) \\ &= \phi^*(\Psi_{K'}(\xi)) \\ &= [\phi^* \circ \Psi_{K'}](\xi).\end{aligned}$$

- *Naturality of Φ .* Since Ψ is natural and Ψ is a componentwise inverse to Φ , it follows from [Item 2 of Proposition 8.8.6.2](#) that Φ is also natural.
- *Invertibility I.* We claim that

$$\Psi_K \circ \Phi_K = \text{id}_{\text{Sets}_*(K, A \hat{\curvearrowright} X)}.$$

Indeed, given a morphism of pointed sets

$$\xi: (K, k_0) \rightarrow (A \hat{\curvearrowright} X, [(x_0)_{a \in A}])$$

we have

$$\begin{aligned}[\Psi_K \circ \Phi_K](\xi) &= \Psi_K(\Phi_K(\xi)) \\ &= \Psi_K(\llbracket a \mapsto \xi_a \rrbracket) \\ &= \Psi_K(\llbracket a' \mapsto \xi_{a'} \rrbracket) \\ &= \llbracket k \mapsto [(\text{ev}_a(\llbracket a' \mapsto \xi_{a'}(k) \rrbracket))_{a \in A}] \rrbracket \\ &= \llbracket k \mapsto [(\xi_a(k))_{a \in A}] \rrbracket.\end{aligned}$$

Now, we have two cases:

1. If $\xi(k) = [(x_0)_{a \in A}]$, we have

$$\begin{aligned} [\Psi_K \circ \Phi_K](\xi) &= \dots \\ &= \llbracket k \mapsto [(\xi_a(k))_{a \in A}] \rrbracket \\ &= \llbracket k \mapsto [(x_0)_{a \in A}] \rrbracket \\ &= \llbracket k \mapsto \xi(k) \rrbracket \\ &= \xi. \end{aligned}$$

2. If $\xi(k) \neq [(x_0)_{a \in A}]$ and $\xi(k) = [(x_a^k)_{a \in A}]$ instead, we have

$$\begin{aligned} [\Psi_K \circ \Phi_K](\xi) &= \dots \\ &= \llbracket k \mapsto [(\xi_a(k))_{a \in A}] \rrbracket \\ &= \llbracket k \mapsto [(x_a^k)_{a \in A}] \rrbracket \\ &= \llbracket k \mapsto \xi(k) \rrbracket \\ &= \xi. \end{aligned}$$


In both cases, we have $[\Psi_K \circ \Phi_K](\xi) = \xi$, and thus we are done.

· *Invertibility II.* We claim that

$$\Phi_K \circ \Psi_K = \text{id}_{\text{Sets}(A, \text{Sets}_*(K, X))}.$$

Indeed, given a morphism $\xi: A \rightarrow \text{Sets}_*(K, X)$, we have

$$\begin{aligned} [\Phi_K \circ \Psi_K](\xi) &= \Phi_K(\Psi_K(\xi)) \\ &= \Phi_K(\llbracket k \mapsto [(\xi_a(k))_{a \in A}] \rrbracket) \\ &= \llbracket a \mapsto \llbracket k \mapsto \xi_a(k) \rrbracket \rrbracket \\ &= \xi \end{aligned}$$

This finishes the proof. 

00D1

PROPOSITION 4.2.2.6 ► PROPERTIES OF COTENSORS OF POINTED SETS BY SETS

Let (X, x_0) be a pointed set and let A be a set.

00D2

1. *Functoriality.* The assignments $A, (X, x_0), (A, (X, x_0))$ define functors

$$\begin{aligned} A \pitchfork - &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ - \pitchfork X &: \mathbf{Sets}^{\text{op}} \rightarrow \mathbf{Sets}_*, \\ -_1 \pitchfork -_2 &: \mathbf{Sets}^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given:

- A map of sets $f: A \rightarrow B$;
- A pointed map $\phi: (X, x_0) \rightarrow (Y, y_0)$;

the induced map

$$f \odot \phi: A \pitchfork X \rightarrow B \pitchfork Y$$

is given by

$$[f \odot \phi]([(x_a)_{a \in A}]) \stackrel{\text{def}}{=} [(\phi(x_{f(a)}))_{a \in A}]$$

for each $[(x_a)_{a \in A}] \in A \pitchfork X$.

00D3

2. *Adjointness I.* We have an adjunction

$$(- \pitchfork X \dashv \mathbf{Sets}_*(-, X)): \mathbf{Sets}^{\text{op}} \begin{array}{c} \xrightarrow{- \pitchfork X} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(-, X)} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection

$$\mathbf{Sets}_*^{\text{op}}(A \pitchfork X, K) \cong \mathbf{Sets}(A, \mathbf{Sets}_*(K, X)),$$

i.e. by a bijection

$$\mathbf{Sets}_*(K, A \pitchfork X) \cong \mathbf{Sets}(A, \mathbf{Sets}_*(K, X)),$$

natural in $A \in \text{Obj}(\mathbf{Sets})$ and $X, Y \in \text{Obj}(\mathbf{Sets}_*)$.

00D4

3. *Adjointness II.* We have an adjunctions

$$(A \odot - \dashv A \pitchfork -): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{A \odot -} \\ \perp \\ \xleftarrow{A \pitchfork -} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection

$$\text{Hom}_{\mathbf{Sets}_*}(A \odot X, Y) \cong \text{Hom}_{\mathbf{Sets}_*}(X, A \pitchfork Y),$$

natural in $A \in \text{Obj}(\mathbf{Sets})$ and $X, Y \in \text{Obj}(\mathbf{Sets}_*)$.

00D5

4. *As a Weighted Limit.* We have

$$A \pitchfork X \cong \lim^{[A]}(X),$$

where in the right hand side we write:

- A for the functor $A: \text{pt} \rightarrow \text{Sets}$ picking $A \in \text{Obj}(\text{Sets})$;
- X for the functor $X: \text{pt} \rightarrow \text{Sets}_*$ picking $(X, x_0) \in \text{Obj}(\text{Sets}_*)$.

00D6

5. *Iterated Cotensors.* We have an isomorphism of pointed sets

$$A \pitchfork (B \pitchfork X) \cong (A \times B) \pitchfork X,$$

natural in $A, B \in \text{Obj}(\text{Sets})$ and $(X, x_0) \in \text{Obj}(\text{Sets}_*)$.

00D7

6. *Commutativity With Homs.* We have natural isomorphisms

$$\begin{aligned} A \pitchfork \text{Sets}_*(X, -) &\cong \text{Sets}_*(A \odot X, -), \\ A \pitchfork \text{Sets}_*(-, Y) &\cong \text{Sets}_*(-, A \pitchfork Y). \end{aligned}$$

00D8

7. *The Cotensor Evaluation Map.* For each $X, Y \in \text{Obj}(\text{Sets}_*)$, we have a map

$$\text{ev}_{X,Y}^{\pitchfork}: X \rightarrow \text{Sets}_*(X, Y) \pitchfork Y,$$

natural in $X, Y \in \text{Obj}(\text{Sets}_*)$, and given by

$$\text{ev}_{X,Y}^{\pitchfork}(x) \stackrel{\text{def}}{=} \left[(f(x))_{f \in \text{Sets}_*(X,Y)} \right]$$

for each $x \in X$.

00D9

8. *The Cotensor Coevaluation Map.* For each $X \in \text{Obj}(\text{Sets}_*)$ and each $A \in \text{Obj}(\text{Sets})$, we have a map

$$\text{coev}_{A,X}^{\pitchfork}: A \rightarrow \text{Sets}_*(A \pitchfork X, X),$$

natural in $X \in \text{Obj}(\text{Sets}_*)$ and $A \in \text{Obj}(\text{Sets})$, and given by

$$\text{coev}_{A,X}^{\pitchfork}(a) \stackrel{\text{def}}{=} \llbracket [(x_b)_{b \in A}] \mapsto x_a \rrbracket$$

for each $a \in A$.

PROOF 4.2.2.7 ► PROOF OF PROPOSITION 4.2.2.6

Item 1: Functoriality

This is the special case of ?? of ?? for when $C = \text{Sets}_*$.

Item 2: Adjointness I

This is simply a rephrasing of **Definition 4.2.2.1**.

Item 3: : Adjointness II

This is the special case of ?? of ?? for when $C = \text{Sets}_*$.

Item 4: As a Weighted Limit

This is the special case of ?? of ?? for when $C = \text{Sets}_*$.

Item 5: Iterated Cotensors

This is the special case of ?? of ?? for when $C = \text{Sets}_*$.


Item 6: Commutativity With Homs

This is the special case of ?? of ?? for when $C = \text{Sets}_*$.

Item 7: The Cotensor Evaluation Map

This is the special case of ?? of ?? for when $C = \text{Sets}_*$.

Item 8: The Cotensor Coevaluation Map

This is the special case of ?? of ?? for when $C = \text{Sets}_*$. 

00DA **4.3 The Left Tensor Product of Pointed Sets**00DB **4.3.1 Foundations**

Let (X, x_0) and (Y, y_0) be pointed sets.

00DC **DEFINITION 4.3.1.1 ► THE LEFT TENSOR PRODUCT OF POINTED SETS**

The **left tensor product of pointed sets** is the functor¹

$$\triangleleft : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

defined as the composition

$$\text{Sets}_* \times \text{Sets}_* \xrightarrow{\text{id} \times \omega} \text{Sets}_* \times \text{Sets} \xrightarrow{\beta_{\text{Sets}_*, \text{Sets}}^{\text{Cats}_2}} \text{Sets} \times \text{Sets}_* \xrightarrow{\odot} \text{Sets}_*,$$

where:

· $\omega : \text{Sets}_* \rightarrow \text{Sets}$ is the forgetful functor from pointed sets to sets.

- $\beta_{\text{Sets}_*, \text{Sets}}^{\text{Cats}_2} : \text{Sets}_* \times \text{Sets} \xrightarrow{\cong} \text{Sets} \times \text{Sets}_*$ is the braiding of Cats_2 , i.e. the functor witnessing the isomorphism

$$\text{Sets}_* \times \text{Sets} \cong \text{Sets} \times \text{Sets}_*.$$

- $\odot : \text{Sets} \times \text{Sets}_* \rightarrow \text{Sets}_*$ is the tensor functor of **Item 1 of Proposition 4.2.1.9**.

¹Further Notation: Also written $\triangleleft_{\text{Sets}_*}$.

00DD

REMARK 4.3.1.2 ► UNWINDING DEFINITION 4.3.1.1: UNIVERSAL PROPERTY I

The left tensor product of pointed sets satisfies the following natural bijection:

$$\text{Sets}_*(X \triangleleft Y, Z) \cong \text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z).$$

That is to say, the following data are in natural bijection:

1. Pointed maps $f : X \triangleleft Y \rightarrow Z$.
2. Maps of sets $f : X \times Y \rightarrow Z$ satisfying $f(x_0, y) = z_0$ for each $y \in Y$.

00DE

REMARK 4.3.1.3 ► UNWINDING DEFINITION 4.3.1.1: UNIVERSAL PROPERTY II

The left tensor product of pointed sets may be described as follows:

- The left tensor product of (X, x_0) and (Y, y_0) is the pair $((X \triangleleft Y, x_0 \triangleleft y_0), \iota)$ consisting of
 - A pointed set $(X \triangleleft Y, x_0 \triangleleft y_0)$;
 - A left bilinear morphism of pointed sets $\iota : (X \times Y, (x_0, y_0)) \rightarrow X \triangleleft Y$;

satisfying the following universal property:

(UP) Given another such pair $((Z, z_0), f)$ consisting of

- * A pointed set (Z, z_0) ;
- * A left bilinear morphism of pointed sets $f : (X \times Y, (x_0, y_0)) \rightarrow Z$;

there exists a unique morphism of pointed sets $X \triangleleft Y \xrightarrow{\exists!} Z$

making the diagram

$$\begin{array}{ccc} & & X \triangleleft Y \\ & \nearrow \iota & \vdots \exists! \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commute.

00DF

CONSTRUCTION 4.3.1.4 ► THE LEFT TENSOR PRODUCT OF POINTED SETS

In detail, the **left tensor product of** (X, x_0) **and** (Y, y_0) is the pointed set $(X \triangleleft Y, [x_0])$ consisting of

- *The Underlying Set.* The set $X \triangleleft Y$ defined by

$$\begin{aligned} X \triangleleft Y &\stackrel{\text{def}}{=} |Y| \odot X \\ &\cong \bigvee_{y \in Y} (X, x_0), \end{aligned}$$

where $|Y|$ denotes the underlying set of (Y, y_0) ;

- *The Underlying Basepoint.* The point $[(y_0, x_0)]$ of $\bigvee_{y \in Y} (X, x_0)$, which is equal to $[(y, x_0)]$ for any $y \in Y$.

00DG

NOTATION 4.3.1.5 ► ELEMENTS OF LEFT TENSOR PRODUCTS OF POINTED SETS

We write¹ $x \triangleleft y$ for the element $[(y, x)]$ of

$$X \triangleleft Y \cong |Y| \odot X.$$

¹Further Notation: Also written $x \triangleleft_{\text{Sets.}} y$.

00DH

REMARK 4.3.1.6 ► BASEPOINTS OF LEFT TENSOR PRODUCTS OF POINTED SETS

Employing the notation introduced in [Notation 4.3.1.5](#), we have

$$x_0 \triangleleft y_0 = x_0 \triangleleft y$$

for each $y \in Y$, and

$$x_0 \triangleleft y = x_0 \triangleleft y'$$

for each $y, y' \in Y$.

00DJ **PROPOSITION 4.3.1.7 ► PROPERTIES OF LEFT TENSOR PRODUCTS OF POINTED SETS**

Let (X, x_0) and (Y, y_0) be pointed sets.

00DK 1. *Functoriality.* The assignments $X, Y, (X, Y) \mapsto X \triangleleft Y$ define functors

$$\begin{aligned} X \triangleleft - &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ - \triangleleft Y &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ -_1 \triangleleft -_2 &: \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f &: (X, x_0) \rightarrow (A, a_0), \\ g &: (Y, y_0) \rightarrow (B, b_0), \end{aligned}$$

the induced map

$$f \triangleleft g: X \triangleleft Y \rightarrow A \triangleleft B$$

is given by

$$[f \triangleleft g](x \triangleleft y) \stackrel{\text{def}}{=} f(x) \triangleleft g(y)$$

for each $x \triangleleft y \in X \triangleleft Y$.

00DL 2. *Adjointness I.* We have an adjunction

$$\left(- \triangleleft Y \dashv [Y, -]_{\mathbf{Sets}_*}^{\triangleleft} \right): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \triangleleft Y} \\ \perp \\ \xleftarrow{[Y, -]_{\mathbf{Sets}_*}^{\triangleleft}} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}\left(X, [Y, Z]_{\mathbf{Sets}_*}^{\triangleleft}\right)$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$, where $[X, Y]_{\mathbf{Sets}_*}^{\triangleleft}$ is the pointed set of [Definition 4.3.2.1](#).

00DM 3. *Adjointness II.* The functor

$$X \triangleleft -: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

does not admit a right adjoint.

00DN 4. *Adjointness III.* We have a bijection of sets

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(|Y|, \mathbf{Sets}_*(X, Z))$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$.

PROOF 4.3.1.8 ► PROOF OF PROPOSITION 4.3.1.7

Item 1: Functoriality

Clear.

Item 2: Adjointness I

This follows from **Item 3** of **Proposition 4.2.1.9**.

Item 3: Adjointness II

For $X \triangleleft -$ to admit a right adjoint would require it to preserve colimits by ?? of ???. However, we have

$$\begin{aligned} X \triangleleft \text{pt} &\stackrel{\text{def}}{=} | \text{pt} | \odot X \\ &\cong X \\ &\neq \text{pt}, \end{aligned}$$

and thus we see that $X \triangleleft -$ does not have a right adjoint.

Item 4: Adjointness III

This follows from **Item 2** of **Proposition 4.2.1.9**. 

00DP

REMARK 4.3.1.9 ► ON THE FAILURE OF $X \triangleleft -$ TO BE A LEFT ADJOINT

Here is some intuition on why $X \triangleleft -$ fails to be a left adjoint. **Item 4** of **Proposition 4.3.1.7** states that we have a natural bijection

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(|Y|, \mathbf{Sets}_*(X, Z)),$$

so it would be reasonable to wonder whether a natural bijection of the form

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(Y, \mathbf{Sets}_*(X, Z)),$$

also holds, which would give $X \triangleleft - \dashv \mathbf{Sets}_*(X, -)$. However, such a bijection would require every map

$$f: X \triangleleft Y \rightarrow Z$$

to satisfy

$$f(x \triangleleft y_0) = z_0$$

for each $x \in X$, whereas we are imposing such a basepoint preservation condition only for elements of the form $x_0 \triangleleft y$. Thus $\mathbf{Sets}_*(X, -)$ can't be a right adjoint for $X \triangleleft -$, and as shown by **Item 3** of **Proposition 4.3.1.7**, no functor can.¹

¹The functor $\mathbf{Sets}_*(X, -)$ is instead right adjoint to $X \wedge -$, the smash product of pointed sets of **Definition 4.5.1.1**. See **Item 2** of **Proposition 4.5.1.10**.

00DQ 4.3.2 The Left Internal Hom of Pointed Sets

Let (X, x_0) and (Y, y_0) be pointed sets.

00DR DEFINITION 4.3.2.1 ► THE LEFT INTERNAL HOM OF POINTED SETS

The **left internal Hom of pointed sets** is the functor

$$[-, -]_{\mathbf{Sets}_*}^{\triangleleft} : \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

defined as the composition

$$\mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \xrightarrow{\text{forget} \times \text{id}} \mathbf{Sets}^{\text{op}} \times \mathbf{Sets}_* \xrightarrow{\pitchfork} \mathbf{Sets}_*,$$

where:

- $\text{forget} : \mathbf{Sets}_* \rightarrow \mathbf{Sets}$ is the forgetful functor from pointed sets to sets.
- $\pitchfork : \mathbf{Sets}^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$ is the cotensor functor of [Item 1 of Proposition 4.2.2.6](#).

PROOF 4.3.2.2 ► PROOF OF DEFINITION 4.3.2.1

For a proof that $[-, -]_{\mathbf{Sets}_*}^{\triangleleft}$ is indeed the left internal Hom of \mathbf{Sets}_* with respect to the left tensor product of pointed sets, see [Item 2 of Proposition 4.3.1.7](#).



00DS REMARK 4.3.2.3 ► UNWINDING DEFINITION 4.3.2.1, I: UNIVERSAL PROPERTY

The left internal Hom of pointed sets satisfies the following universal property:

$$\mathbf{Sets}_*(X \triangleleft Y, Z) \cong \mathbf{Sets}_*\left(X, [Y, Z]_{\mathbf{Sets}_*}^{\triangleleft}\right)$$

That is to say, the following data are in bijection:

1. Pointed maps $f : X \triangleleft Y \rightarrow Z$.
2. Pointed maps $f : X \rightarrow [Y, Z]_{\mathbf{Sets}_*}^{\triangleleft}$.

00DT REMARK 4.3.2.4 ► UNWINDING DEFINITION 4.3.2.1, II: EXPLICIT DESCRIPTION

In detail, the **left internal Hom of (X, x_0) and (Y, y_0)** is the pointed set $\left([X, Y]_{\mathbf{Sets}_*}^{\triangleleft}, [(y_0)_{x \in X}]\right)$ consisting of

• *The Underlying Set.* The set $[X, Y]_{\mathbf{Sets}_*}^{\triangleleft}$ defined by

$$[X, Y]_{\mathbf{Sets}_*}^{\triangleleft} \stackrel{\text{def}}{=} |X| \pitchfork Y \\ \cong \bigwedge_{x \in X} (Y, y_0),$$

where $|X|$ denotes the underlying set of (X, x_0) ;

• *The Underlying Basepoint.* The point $[(y_0)_{x \in X}]$ of $\bigwedge_{x \in X} (Y, y_0)$.

00DU

PROPOSITION 4.3.2.5 ► PROPERTIES OF LEFT INTERNAL HOMS OF POINTED SETS

Let (X, x_0) and (Y, y_0) be pointed sets.

00DV

1. *Functoriality.* The assignments $X, Y, (X, Y) \mapsto [X, Y]_{\mathbf{Sets}_*}^{\triangleleft}$ define functors

$$[X, -]_{\mathbf{Sets}_*}^{\triangleleft} : \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ [-, Y]_{\mathbf{Sets}_*}^{\triangleleft} : \mathbf{Sets}_*^{\text{op}} \rightarrow \mathbf{Sets}_*, \\ [-1, -2]_{\mathbf{Sets}_*}^{\triangleleft} : \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*.$$

In particular, given pointed maps

$$f : (X, x_0) \rightarrow (A, a_0), \\ g : (Y, y_0) \rightarrow (B, b_0),$$

the induced map

$$[f, g]_{\mathbf{Sets}_*}^{\triangleleft} : [A, Y]_{\mathbf{Sets}_*}^{\triangleleft} \rightarrow [X, B]_{\mathbf{Sets}_*}^{\triangleleft}$$

is given by

$$[f, g]_{\mathbf{Sets}_*}^{\triangleleft}([(y_a)_{a \in A}]) \stackrel{\text{def}}{=} [(g(y_{f(x)}))_{x \in X}]$$

for each $[(y_a)_{a \in A}] \in [A, Y]_{\mathbf{Sets}_*}^{\triangleleft}$.

00DW

2. *Adjointness I.* We have an adjunction

$$\left(- \triangleleft Y \dashv [Y, -]_{\mathbf{Sets}_*}^{\triangleleft} \right) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \triangleleft Y} \\ \perp \\ \xleftarrow{[Y, -]_{\mathbf{Sets}_*}^{\triangleleft}} \end{array} \mathbf{Sets}_*$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Sets}_*}(X \triangleleft Y, Z) \cong \text{Hom}_{\text{Sets}_*}(X, [Y, Z]_{\text{Sets}_*}^{\triangleleft})$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$

00DX

3. *Adjointness II.* The functor

$$X \triangleleft -: \text{Sets}_* \rightarrow \text{Sets}_*$$

does not admit a right adjoint.

PROOF 4.3.2.6 ► PROOF OF PROPOSITION 4.3.2.5

Item 1: Functoriality

Clear.

Item 2: Adjointness I

This is a repetition of **Item 2** of **Proposition 4.3.1.7**, and is proved there.

Item 3: Adjointness II

This is a repetition of **Item 3** of **Proposition 4.3.1.7**, and is proved there. 

00DY 4.3.3 The Left Skew Unit

00DZ

DEFINITION 4.3.3.1 ► THE LEFT SKEW UNIT OF \triangleleft

The **left skew unit of the left tensor product of pointed sets** is the functor

$$\mathbb{1}^{\text{Sets}_*, \triangleleft} : \text{pt} \rightarrow \text{Sets}_*$$

defined by

$$\mathbb{1}_{\text{Sets}_*}^{\triangleleft} \stackrel{\text{def}}{=} S^0.$$

00E0 4.3.4 The Left Skew Associator

00E1

DEFINITION 4.3.4.1 ► THE LEFT SKEW ASSOCIATOR OF \triangleleft

The **skew associator of the left tensor product of pointed sets** is the natural transformation

$$\alpha^{\text{Sets}_*, \triangleleft} : \triangleleft \circ (\triangleleft \times \text{id}_{\text{Sets}_*}) \Longrightarrow \triangleleft \circ (\text{id}_{\text{Sets}_*} \times \triangleleft) \circ \alpha_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}^{\text{Cats}}$$

as in the diagram

$$\begin{array}{ccc}
 & \text{Sets}_* \times (\text{Sets}_* \times \text{Sets}_*) & \\
 \alpha_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}^{\text{Cats}} \dashrightarrow & & \text{id} \times \triangleleft \\
 (\text{Sets}_* \times \text{Sets}_*) \times \text{Sets}_* & \xrightarrow{\alpha_{\text{Sets}_*, \triangleleft}} & \text{Sets}_* \times \text{Sets}_* \\
 \triangleleft \times \text{id} \searrow & & \searrow \triangleleft \\
 \text{Sets}_* \times \text{Sets}_* & \xrightarrow{\triangleleft} & \text{Sets}_*
 \end{array}$$

whose component

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft} : (X \triangleleft Y) \triangleleft Z \rightarrow X \triangleleft (Y \triangleleft Z)$$


at $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ is given by

$$\begin{aligned}
 (X \triangleleft Y) \triangleleft Z &\stackrel{\text{def}}{=} |Z| \odot (X \triangleleft Y) \\
 &\stackrel{\text{def}}{=} |Z| \odot (|Y| \odot X) \\
 &\cong \bigvee_{z \in Z} |Y| \odot X \\
 &\cong \bigvee_{z \in Z} \left(\bigvee_{y \in Y} X \right) \\
 &\rightarrow \bigvee_{[(z,y)] \in \bigvee_{z \in Z} Y} X \\
 &\cong \bigvee_{[(z,y)] \in |Z| \odot Y} X \\
 &\cong ||Z| \odot Y| \odot X \\
 &\stackrel{\text{def}}{=} |Y \triangleleft Z| \odot X \\
 &\stackrel{\text{def}}{=} X \triangleleft (Y \triangleleft Z),
 \end{aligned}$$

where the map

$$\bigvee_{z \in Z} \left(\bigvee_{y \in Y} X \right) \rightarrow \bigvee_{(z,y) \in \bigvee_{z \in Z} Y} X$$

is given by $[(z, [(y, x)])] \mapsto [([(z, y)], x)]$.

PROOF 4.3.4.2 ► PROOF OF DEFINITION 4.3.4.1(Proven below in a bit.) **00E2 REMARK 4.3.4.3 ► UNWINDING DEFINITION 4.3.4.1**

Unwinding the notation for elements, we have

$$\begin{aligned} [(z, [(y, x)])] &\stackrel{\text{def}}{=} [(z, x \triangleleft y)] \\ &\stackrel{\text{def}}{=} (x \triangleleft y) \triangleleft z \end{aligned}$$

and

$$\begin{aligned} [([(z, y)], x)] &\stackrel{\text{def}}{=} [(y \triangleleft z, x)] \\ &\stackrel{\text{def}}{=} x \triangleleft (y \triangleleft z). \end{aligned}$$

So, in other words, $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}$ acts on elements via

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}((x \triangleleft y) \triangleleft z) \stackrel{\text{def}}{=} x \triangleleft (y \triangleleft z)$$

for each $(x \triangleleft y) \triangleleft z \in (X \triangleleft Y) \triangleleft Z$.**00E3 REMARK 4.3.4.4 ► NON-INVERTIBILITY OF THE SKEW ASSOCIATOR OF \triangleleft** Taking $y = y_0$, we see that the morphism $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}$ acts on elements as

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}((x \triangleleft y_0) \triangleleft z) \stackrel{\text{def}}{=} x \triangleleft (y_0 \triangleleft z).$$

However, by the definition of \triangleleft , we have $y_0 \triangleleft z = y_0 \triangleleft z'$ for all $z, z' \in Z$, preventing $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}$ from being non-invertible.**PROOF 4.3.4.5 ► PROOF OF DEFINITION 4.3.4.1**Firstly, note that, given $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$, the map

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft} : (X \triangleleft Y) \triangleleft Z \rightarrow X \triangleleft (Y \triangleleft Z)$$

is indeed a morphism of pointed sets, as we have

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}((x_0 \triangleleft y_0) \triangleleft z_0) = x_0 \triangleleft (y_0 \triangleleft z_0).$$

Next, we claim that $\alpha^{\text{Sets}_*, \triangleleft}$ is a natural transformation. We need to show that, given morphisms of pointed sets


$$\begin{aligned} f &: (X, x_0) \rightarrow (X', x'_0), \\ g &: (Y, y_0) \rightarrow (Y', y'_0), \\ h &: (Z, z_0) \rightarrow (Z', z'_0) \end{aligned}$$

the diagram

$$\begin{array}{ccc} (X \triangleleft Y) \triangleleft Z & \xrightarrow{(f \triangleleft g) \triangleleft h} & (X' \triangleleft Y') \triangleleft Z' \\ \alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft} \downarrow & & \downarrow \alpha_{X',Y',Z'}^{\text{Sets}_*, \triangleleft} \\ X \triangleleft (Y \triangleleft Z) & \xrightarrow{f \triangleleft (g \triangleleft h)} & X' \triangleleft (Y' \triangleleft Z') \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (x \triangleleft y) \triangleleft z & \longmapsto & (f(x) \triangleleft g(y)) \triangleleft h(z) \\ \downarrow & & \downarrow \\ x \triangleleft (y \triangleleft z) & \longmapsto & f(x) \triangleleft (g(y) \triangleleft h(z)) \end{array}$$

and hence indeed commutes, showing $\alpha^{\text{Sets}_*, \triangleleft}$ to be a natural transformation. This finishes the proof. 

00E4 4.3.5 The Left Skew Left Unitor

00E5 DEFINITION 4.3.5.1 ► THE LEFT SKEW LEFT UNITOR OF \triangleleft

The **skew left unitor of the left tensor product of pointed sets** is the natural transformation

$$\lambda^{\text{Sets}_*, \triangleleft} : \triangleleft \circ (\mathbb{1}_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \xrightarrow{\sim} \lambda_{\text{Sets}_*}^{\text{Cats}_2}$$

whose component

$$\lambda_X^{\text{Sets}_*, \triangleleft} : S^0 \triangleleft X \rightarrow X$$

at $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ is given by the composition

$$\begin{aligned} S^0 \triangleleft X &\cong |X| \odot S^0 \\ &\cong \bigvee_{x \in X} S^0 \\ &\rightarrow X, \end{aligned}$$

where $\bigvee_{x \in X} S^0 \rightarrow X$ is the map given by

$$\begin{aligned} [(x, 0)] &\mapsto x_0, \\ [(x, 1)] &\mapsto x. \end{aligned}$$

PROOF 4.3.5.2 ► PROOF OF DEFINITION 4.3.5.1

(Proven below in a bit.)

00E6 REMARK 4.3.5.3 ► UNWINDING DEFINITION 4.3.5.1

In other words, $\lambda_X^{\text{Sets}_*, \triangleleft}$ acts on elements as

$$\begin{aligned} \lambda_X^{\text{Sets}_*, \triangleleft}(0 \triangleleft x) &\stackrel{\text{def}}{=} x_0, \\ \lambda_X^{\text{Sets}_*, \triangleleft}(1 \triangleleft x) &\stackrel{\text{def}}{=} x \end{aligned}$$

for each $1 \triangleleft x \in S^0 \triangleleft X$.

00E7

REMARK 4.3.5.4 ► NON-INVERTIBILITY OF THE SKEW LEFT UNITOR OF \triangleleft

The morphism $\lambda_X^{\text{Sets}_*, \triangleleft}$ is almost invertible, with its would-be-inverse

$$\phi_X: X \rightarrow S^0 \triangleleft X$$

given by

$$\phi_X(x) \stackrel{\text{def}}{=} 1 \triangleleft x$$

for each $x \in X$. Indeed, we have

$$\begin{aligned} \left[\lambda_X^{\text{Sets}_*, \triangleleft} \circ \phi \right](x) &= \lambda_X^{\text{Sets}_*, \triangleleft}(\phi(x)) \\ &= \lambda_X^{\text{Sets}_*, \triangleleft}(1 \triangleleft x) \\ &= x \\ &= [\text{id}_X](x) \end{aligned}$$

so that

$$\lambda_X^{\text{Sets}_*, \triangleleft} \circ \phi = \text{id}_X$$

and

$$\begin{aligned} \left[\phi \circ \lambda_X^{\text{Sets}_*, \triangleleft} \right](1 \triangleleft x) &= \phi\left(\lambda_X^{\text{Sets}_*, \triangleleft}(1 \triangleleft x)\right) \\ &= \phi(x) \\ &= 1 \triangleleft x \\ &= [\text{id}_{S^0 \triangleleft X}](1 \triangleleft x), \end{aligned}$$

but

$$\begin{aligned} \left[\phi \circ \lambda_X^{\text{Sets}_*, \triangleleft} \right](0 \triangleleft x) &= \phi\left(\lambda_X^{\text{Sets}_*, \triangleleft}(0 \triangleleft x)\right) \\ &= \phi(x_0) \\ &= 1 \triangleleft x_0, \end{aligned}$$

where $0 \triangleleft x \neq 1 \triangleleft x_0$. Thus

$$\phi \circ \lambda_X^{\text{Sets}_*, \triangleleft} \stackrel{?}{=} \text{id}_{S^0 \triangleleft X}$$

holds for all elements in $S^0 \triangleleft X$ except one.

PROOF 4.3.5.5 ► PROOF OF DEFINITION 4.3.5.1

Firstly, note that, given $(X, x_0) \in \text{Obj}(\text{Sets}_*)$, the map

$$\lambda_X^{\text{Sets}_*, \triangleleft} : S^0 \triangleleft X \rightarrow X$$

is indeed a morphism of pointed sets, as we have

$$\lambda_X^{\text{Sets}_*, \triangleleft}(0 \triangleleft x_0) = x_0.$$

Next, we claim that $\lambda^{\text{Sets}_*, \triangleleft}$ is a natural transformation. We need to show that, given a morphism of pointed sets

$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram


$$\begin{array}{ccc} S^0 \triangleleft X & \xrightarrow{\text{id}_{S^0} \triangleleft f} & S^0 \triangleleft Y \\ \lambda_X^{\text{Sets}_*, \triangleleft} \downarrow & & \downarrow \lambda_Y^{\text{Sets}_*, \triangleleft} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} 0 \triangleleft x & & 0 \triangleleft x \mapsto 0 \triangleleft f(x) \\ \downarrow & & \downarrow \\ x_0 \mapsto & f(x_0) & y_0 \end{array}$$

and

$$\begin{array}{ccc} 1 \triangleleft x \mapsto & 1 \triangleleft f(x) & \\ \downarrow & \downarrow & \\ x \mapsto & f(x) & \end{array}$$

and hence indeed commutes, showing $\lambda^{\text{Sets}_*, \triangleleft}$ to be a natural transformation. This finishes the proof. 

00E8 4.3.6 The Left Skew Right Unitor

00E9

DEFINITION 4.3.6.1 ▶ THE LEFT SKEW RIGHT UNITOR OF \triangleleft

The **skew right unitor of the left tensor product of pointed sets** is the natural transformation

$$\rho^{\text{Sets}_*, \triangleleft} : \rho_{\text{Sets}_*}^{\text{Cats}_2} \xrightarrow{\sim} \triangleleft \circ (\text{id} \times \mathbb{1}_{\text{Sets}_*}),$$

The diagram shows a commutative square with a diagonal arrow. The top-left node is $\text{Sets}_* \times \text{pt}$, the top-right node is $\text{Sets}_* \times \text{Sets}_*$, and the bottom-right node is Sets_* . A horizontal arrow labeled $\text{id} \times \mathbb{1}_{\text{Sets}_*}$ points from $\text{Sets}_* \times \text{pt}$ to $\text{Sets}_* \times \text{Sets}_*$. A vertical arrow labeled \triangleleft points from $\text{Sets}_* \times \text{Sets}_*$ to Sets_* . A dashed curved arrow labeled $\rho_{\text{Sets}_*}^{\text{Cats}_2}$ points from $\text{Sets}_* \times \text{pt}$ to Sets_* . A solid double arrow labeled $\rho^{\text{Sets}_*, \triangleleft}$ points from $\text{Sets}_* \times \text{pt}$ to Sets_* .

whose component

$$\rho_X^{\text{Sets}_*, \triangleleft} : X \rightarrow X \triangleleft S^0$$

at $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ is given by the composition

$$\begin{aligned} X &\rightarrow X \vee X \\ &\cong |S^0| \odot X \\ &\cong X \triangleleft S^0, \end{aligned}$$

where $X \rightarrow X \vee X$ is the map sending X to the second factor of X in $X \vee X$.

PROOF 4.3.6.2 ▶ PROOF OF DEFINITION 4.3.6.1

(Proven below in a bit.)



00EA

REMARK 4.3.6.3 ▶ UNWINDING DEFINITION 4.3.6.1

In other words, $\rho_X^{\text{Sets}_*, \triangleleft}$ acts on elements as

$$\rho_X^{\text{Sets}_*, \triangleleft}(x) \stackrel{\text{def}}{=} [(1, x)]$$

i.e. by

$$\rho_X^{\text{Sets}_*, \triangleleft}(x) \stackrel{\text{def}}{=} x \triangleleft 1$$

for each $x \in X$.

00EB

REMARK 4.3.6.4 ► NON-INVERTIBILITY OF THE SKEW RIGHT UNITOR OF \triangleleft

The morphism $\rho_X^{\text{Sets}_*, \triangleleft}$ is non-invertible, as it is non-surjective when viewed as a map of sets, since the elements $x \triangleleft 0$ of $X \triangleleft S^0$ with $x \neq x_0$ are outside the image of $\rho_X^{\text{Sets}_*, \triangleleft}$, which sends x to $x \triangleleft 1$.

PROOF 4.3.6.5 ► PROOF OF DEFINITION 4.3.6.1

Firstly, note that, given $(X, x_0) \in \text{Obj}(\text{Sets}_*)$, the map

$$\rho_X^{\text{Sets}_*, \triangleleft} : X \rightarrow X \triangleleft S^0$$

is indeed a morphism of pointed sets as we have

$$\begin{aligned} \rho_X^{\text{Sets}_*, \triangleleft}(x_0) &= x_0 \triangleleft 1 \\ &= x_0 \triangleleft 0. \end{aligned}$$

Next, we claim that $\rho^{\text{Sets}_*, \triangleleft}$ is a natural transformation. We need to show that, given a morphism of pointed sets


$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \rho_X^{\text{Sets}_*, \triangleleft} \downarrow & & \downarrow \rho_Y^{\text{Sets}_*, \triangleleft} \\ X \triangleleft S^0 & \xrightarrow{f \triangleleft \text{id}_{S^0}} & Y \triangleleft S^0 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \longmapsto & f(x) \\ \downarrow & & \downarrow \\ x \triangleleft 0 & \longmapsto & f(x) \triangleleft 0 \end{array}$$

and hence indeed commutes, showing $\rho^{\text{Sets}_*, \triangleleft}$ to be a natural transformation. This finishes the proof. 

00EC 4.3.7 The Diagonal

00ED

DEFINITION 4.3.7.1 ▶ THE DIAGONAL OF \triangleleft

The **diagonal of the left tensor product of pointed sets** is the natural transformation

$$\Delta^{\triangleleft} : \text{id}_{\text{Sets}_*} \implies \triangleleft \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2},$$

whose component

$$\Delta_X^{\triangleleft} : (X, x_0) \rightarrow (X \triangleleft X, x_0 \triangleleft x_0)$$

at $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ is given by

$$\Delta_X^{\triangleleft}(x) \stackrel{\text{def}}{=} x \triangleleft x$$

for each $x \in X$.

PROOF 4.3.7.2 ▶ PROOF OF DEFINITION 4.3.7.1

Being a Morphism of Pointed Sets

We have

$$\Delta_X^{\triangleleft}(x_0) \stackrel{\text{def}}{=} x_0 \triangleleft x_0,$$

and thus Δ_X^{\triangleleft} is a morphism of pointed sets.

Naturality

We need to show that, given a morphism of pointed sets


$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X^{\triangleleft} \downarrow & & \downarrow \Delta_Y^{\triangleleft} \\ X \triangleleft X & \xrightarrow{f \triangleleft f} & Y \triangleleft Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \xrightarrow{\quad} & f(x) \\ \downarrow & & \downarrow \\ x \triangleleft x & \xrightarrow{\quad} & f(x) \triangleleft f(x) \end{array}$$

and hence indeed commutes, showing Δ^{\triangleleft} to be natural. 

00EE 4.3.8 The Left Skew Monoidal Structure on Pointed Sets Associated to \triangleleft

00EF

PROPOSITION 4.3.8.1 ► THE LEFT SKEW MONOIDAL STRUCTURE ON POINTED SETS ASSOCIATED TO \triangleleft

The category \mathbf{Sets}_* admits a left-closed left skew monoidal category structure consisting of

- *The Underlying Category.* The category \mathbf{Sets}_* of pointed sets;
- *The Left Skew Monoidal Product.* The left tensor product functor

$$\triangleleft : \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

of [Definition 4.3.1.1](#);

- *The Left Internal Skew Hom.* The left internal Hom functor

$$[-, -]_{\mathbf{Sets}_*}^{\triangleleft} : \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

of [Definition 4.3.2.1](#);

- *The Left Skew Monoidal Unit.* The functor

$$\mathbb{1}^{\mathbf{Sets}_*, \triangleleft} : \text{pt} \rightarrow \mathbf{Sets}_*$$

of [Definition 4.3.3.1](#);

- *The Left Skew Associators.* The natural transformation

$$\alpha^{\mathbf{Sets}_*, \triangleleft} : \triangleleft \circ (\triangleleft \times \text{id}_{\mathbf{Sets}_*}) \Longrightarrow \triangleleft \circ (\text{id}_{\mathbf{Sets}_*} \times \triangleleft) \circ \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\text{Cats}}$$

of [Definition 4.3.4.1](#);

- *The Left Skew Left Unitors.* The natural transformation

$$\lambda^{\text{Sets}_*, \triangleleft} : \triangleleft \circ (\mathbb{1}^{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \xrightarrow{\sim} \lambda_{\text{Sets}_*}^{\text{Cats}_2}$$

of Definition 4.3.5.1;

- *The Left Skew Right Unitors.* The natural transformation


$$\rho^{\text{Sets}_*, \triangleleft} : \rho_{\text{Sets}_*}^{\text{Cats}_2} \xrightarrow{\sim} \triangleleft \circ (\text{id} \times \mathbb{1}^{\text{Sets}_*})$$

of Definition 4.3.6.1.


PROOF 4.3.8.2 ► PROOF OF PROPOSITION 4.3.8.1

The Pentagon Identity

Let (W, w_0) , (X, x_0) , (Y, y_0) and (Z, z_0) be pointed sets. We have to show that the diagram




commutes. Indeed, this diagram acts on elements as




and thus we see that the pentagon identity is satisfied.

The Left Skew Left Triangle Identity


Let (X, x_0) and (Y, y_0) be pointed sets. We have to show that the diagram



commutes. Indeed, this diagram acts on elements as



and



and hence indeed commutes. Thus the left skew triangle identity is satisfied.

The Left Skew Right Triangle Identity

Let (X, x_0) and (Y, y_0) be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleleft Y & & \\
 \rho_{X \triangleleft Y}^{\text{Sets}_*, \triangleleft} \downarrow & \searrow \text{id}_X \triangleleft \rho_Y^{\text{Sets}_*, \triangleleft} & \\
 (X \triangleleft Y) \triangleleft S^0 & \xrightarrow{\alpha_{X, Y, S^0}^{\text{Sets}_*, \triangleleft}} & X \triangleleft (Y \triangleleft S^0)
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleleft y & & \\
 \downarrow & \searrow & \\
 (x \triangleleft y) \triangleleft 1 & \mapsto & x \triangleleft (y \triangleleft 1)
 \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

The Left Skew Middle Triangle Identity

Let (X, x_0) and (Y, y_0) be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleleft Y & \xlongequal{\quad} & X \triangleleft Y \\
 \rho_X^{\text{Sets}_*, \triangleleft} \triangleleft \text{id}_Y \downarrow & & \uparrow \text{id}_A \triangleleft \lambda_Y^{\text{Sets}_*, \triangleleft} \\
 (X \triangleleft S^0) \triangleleft Y & \xrightarrow{\alpha_{X, S^0, Y}^{\text{Sets}_*, \triangleleft}} & X \triangleleft (S^0 \triangleleft Y)
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleleft y & \xrightarrow{\quad} & x \triangleleft y \\
 \downarrow & & \uparrow \\
 (x \triangleleft 1) \triangleleft y & \xrightarrow{\quad} & x \triangleleft (1 \triangleleft y)
 \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

The Zig-Zag Identity

We have to show that the diagram

$$\begin{array}{ccc}
 S^0 & \xrightarrow{\rho_{S^0}^{\text{Sets}_*, \triangleleft}} & S^0 \triangleleft S^0 \\
 \parallel & & \downarrow \lambda_{S^0}^{\text{Sets}_*, \triangleleft} \\
 & & S^0
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 0 & \mapsto & 0 \triangleleft 1 \\
 \swarrow & & \downarrow \\
 & & 0
 \end{array}$$

and

$$\begin{array}{ccc}
 1 & \mapsto & 1 \triangleleft 1 \\
 \swarrow & & \downarrow \\
 & & 1
 \end{array}$$

and hence indeed commutes. Thus the zig-zag identity is satisfied.

Left Skew Monoidal Left-Closedness

This follows from **Item 2** of **Proposition 4.3.1.7**. ☐

00EG 4.3.9 Monoids With Respect to the Left Tensor Product of Pointed Sets

00EH PROPOSITION 4.3.9.1 ► MONOIDS WITH RESPECT TO \triangleleft

The category of monoids on $(\text{Sets}_*, \triangleleft, S^0)$ is isomorphic to the category of “monoids with left zero”¹ and morphisms between them.

¹A monoid with left zero is defined similarly as the monoids with zero of ???. Succinctly, they are monoids (A, μ_A, η_A) with a special element 0_A satisfying

$$0_A a = 0_A$$

for each $a \in A$.

PROOF 4.3.9.2 ► PROOF OF PROPOSITION 4.3.9.1

Monoids on $(\text{Sets}_*, \triangleleft, S^0)$

A monoid on $(\text{Sets}_*, \triangleleft, S^0)$ consists of:

- *The Underlying Object.* A pointed set $(A, 0_A)$.
- *The Multiplication Morphism.* A morphism of pointed sets

$$\mu_A: A \triangleleft A \rightarrow A,$$

determining a left bilinear morphism of pointed sets

$$\begin{aligned} A \times A &\longrightarrow A \\ (a, b) &\longmapsto ab. \end{aligned}$$


- *The Unit Morphism.* A morphism of pointed sets

$$\eta_A: S^0 \rightarrow A$$


picking an element 1_A of A .

satisfying the following conditions:

1. *Associativity.* The diagram



2. *Left Unitality.* The diagram



commutes.

3. *Right Unitality.* The diagram

$$\begin{array}{ccc}
 A & \xrightarrow{\rho_A^{\text{Sets}, \triangleleft}} & A \triangleleft S^0 \\
 \parallel & & \downarrow \text{id}_A \times \eta_A \\
 A & \xleftarrow{\mu_A} & A \triangleleft A
 \end{array}$$


commutes.

Being a left-bilinear morphism of pointed sets, the multiplication map satisfies

$$0_A a = 0_A$$

for each $a \in A$. Now, the associativity, left unitality, and right unitality conditions act on elements as follows:

1. *Associativity.* The associativity condition acts as




This gives

$$(ab)c = a(bc)$$


for each $a, b, c \in A$.

2. *Left Unitality.* The left unitality condition acts:

(a) On $0 \triangleleft a$ as



(b) On $1 \triangleleft a$ as




This gives

$$\begin{aligned}
 1_A a &= a, \\
 0_A a &= 0_A
 \end{aligned}$$

for each $a \in A$.

3. *Right Unitality*. The right unitality condition acts as



This gives

$$a 1_A = a$$

for each $a \in A$.

Thus we see that monoids with respect to \triangleleft are exactly monoids with left zero.


Morphisms of Monoids on $(\text{Sets}_*, \triangleleft, S^0)$

A morphism of monoids on $(\text{Sets}_*, \triangleleft, S^0)$ from $(A, \mu_A, \eta_A, 0_A)$ to $(B, \mu_B, \eta_B, 0_B)$ is a morphism of pointed sets

$$f: (A, 0_A) \rightarrow (B, 0_B)$$

satisfying the following conditions:

1. *Compatibility With the Multiplication Morphisms*. The diagram



commutes.

2. *Compatibility With the Unit Morphisms.* The diagram

$$\begin{array}{ccc} S^0 & \xrightarrow{\eta_A} & A \\ & \searrow \eta_B & \downarrow f \\ & & B \end{array}$$

commutes.

These act on elements as

$$\begin{array}{ccc} a \triangleleft b & & a \triangleleft b \mapsto f(a) \triangleleft f(b) \\ \downarrow & & \downarrow \\ ab \mapsto f(ab) & & f(a)f(b) \end{array}$$

and

$$\begin{array}{ccc} 0 & & 0 \mapsto 0_A \\ & \searrow & \downarrow \\ & & 0_B \quad f(0_A) \end{array}$$

and

$$\begin{array}{ccc} 1 & & 1 \mapsto 1_A \\ & \searrow & \downarrow \\ & & 1_B \quad f(1_A) \end{array}$$

giving

$$\begin{aligned} f(ab) &= f(a)f(b), \\ f(0_A) &= 0_B, \\ f(1_A) &= 1_B, \end{aligned}$$

for each $a, b \in A$, which is exactly a morphism of monoids with left zero.

Identities and Composition

Similarly, the identities and composition of $\text{Mon}(\text{Sets}_*, \triangleleft, S^0)$ can be easily seen to agree with those of monoids with left zero, which finishes the proof. \square

00EJ 4.4 The Right Tensor Product of Pointed Sets

00EK 4.4.1 Foundations

Let (X, x_0) and (Y, y_0) be pointed sets.

00EL DEFINITION 4.4.1.1 ► THE RIGHT TENSOR PRODUCT OF POINTED SETS

The **right tensor product of pointed sets** is the functor¹

$$\triangleright : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

defined as the composition

$$\text{Sets}_* \times \text{Sets}_* \xrightarrow{\text{forgetful}} \text{Sets} \times \text{Sets}_* \xrightarrow{\otimes} \text{Sets}_*,$$

where:

- $\text{forgetful} : \text{Sets}_* \rightarrow \text{Sets}$ is the forgetful functor from pointed sets to sets.
- $\otimes : \text{Sets} \times \text{Sets}_* \rightarrow \text{Sets}_*$ is the tensor functor of [Item 1 of Proposition 4.2.1.9](#).

¹Further Notation: Also written $\triangleright_{\text{Sets}_*}$.

00EM REMARK 4.4.1.2 ► UNWINDING DEFINITION 4.4.1.1: UNIVERSAL PROPERTY I

The right tensor product of pointed sets satisfies the following natural bijection:

$$\text{Sets}_*(X \triangleright Y, Z) \cong \text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z).$$

That is to say, the following data are in natural bijection:

1. Pointed maps $f : X \triangleright Y \rightarrow Z$.
2. Maps of sets $f : X \times Y \rightarrow Z$ satisfying $f(x, y_0) = z_0$ for each $x \in X$.

00EN REMARK 4.4.1.3 ► UNWINDING DEFINITION 4.4.1.1: UNIVERSAL PROPERTY II

The right tensor product of pointed sets may be described as follows:

- The right tensor product of (X, x_0) and (Y, y_0) is the pair $((X \triangleright Y, x_0 \triangleright y_0), \iota)$ consisting of
 - A pointed set $(X \triangleright Y, x_0 \triangleright y_0)$;
 - A right bilinear morphism of pointed sets $\iota : (X \times Y, (x_0, y_0)) \rightarrow$

$$X \triangleright Y;$$

satisfying the following universal property:

(UP) Given another such pair $((Z, z_0), f)$ consisting of

- * A pointed set (Z, z_0) ;
- * A right bilinear morphism of pointed sets $f: (X \times Y, (x_0, y_0)) \rightarrow X \triangleright Y$;

there exists a unique morphism of pointed sets $X \triangleright Y \xrightarrow{\exists!} Z$ making the diagram

$$\begin{array}{ccc} & X \triangleright Y & \\ & \nearrow \iota & \vdots \exists! \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commute.

00EP

CONSTRUCTION 4.4.1.4 ► THE RIGHT TENSOR PRODUCT OF POINTED SETS

In detail, the **right tensor product of** (X, x_0) **and** (Y, y_0) is the pointed set $(X \triangleright Y, [y_0])$ consisting of:

- *The Underlying Set.* The set $X \triangleright Y$ defined by

$$\begin{aligned} X \triangleright Y &\stackrel{\text{def}}{=} |X| \odot Y \\ &\cong \bigvee_{x \in X} (Y, y_0), \end{aligned}$$

where $|X|$ denotes the underlying set of (X, x_0) .

- *The Underlying Basepoint.* The point $[(x_0, y_0)]$ of $\bigvee_{x \in X} (Y, y_0)$, which is equal to $[(x, y_0)]$ for any $x \in X$.

00EQ

NOTATION 4.4.1.5 ► ELEMENTS OF RIGHT TENSOR PRODUCTS OF POINTED SETS

We write¹ $x \triangleright y$ for the element $[(x, y)]$ of

$$X \triangleright Y \cong |X| \odot Y.$$

¹Further Notation: Also written $x \triangleright_{\text{Sets}_*} y$.

00ER **REMARK 4.4.1.6 ► BASEPOINTS OF RIGHT TENSOR PRODUCTS OF POINTED SETS**

Employing the notation introduced in [Notation 4.4.1.5](#), we have

$$x_0 \triangleright y_0 = x \triangleright y_0$$

for each $x \in X$, and

$$x \triangleright y_0 = x' \triangleright y_0$$

for each $x, x' \in X$.

00ES **PROPOSITION 4.4.1.7 ► PROPERTIES OF RIGHT TENSOR PRODUCTS OF POINTED SETS**

Let (X, x_0) and (Y, y_0) be pointed sets.

00ET 1. *Functoriality.* The assignments $X, Y, (X, Y) \mapsto X \triangleright Y$ define functors

$$X \triangleright -: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

$$- \triangleright Y: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

$$-_1 \triangleright -_2: \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*.$$

In particular, given pointed maps

$$f: (X, x_0) \rightarrow (A, a_0),$$

$$g: (Y, y_0) \rightarrow (B, b_0),$$

the induced map

$$f \triangleright g: X \triangleright Y \rightarrow A \triangleright B$$

is given by

$$[f \triangleright g](x \triangleright y) \stackrel{\text{def}}{=} f(x) \triangleright g(y)$$

for each $x \triangleright y \in X \triangleright Y$.

00EU 2. *Adjointness I.* We have an adjunction

$$\left(X \triangleright - \dashv [X, -]_{\mathbf{Sets}_*}^{\triangleright} \right): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \triangleright -} \\ \perp \\ \xleftarrow{[X, -]_{\mathbf{Sets}_*}^{\triangleright}} \end{array} \mathbf{Sets}_*$$

witnessed by a bijection of sets

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleright Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}\left(Y, [X, Z]_{\mathbf{Sets}_*}^{\triangleright}\right)$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$, where $[X, Y]_{\mathbf{Sets}_*}^{\triangleright}$ is the pointed set of [Definition 4.4.2.1](#).

00EV

3. *Adjointness II.* The functor

$$- \triangleright Y : \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

does not admit a right adjoint.

00EW

4. *Adjointness III.* We have a bijection of sets

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleright Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}}(|X|, \mathbf{Sets}_*(Y, Z))$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$.**PROOF 4.4.1.8 ► PROOF OF PROPOSITION 4.4.1.7****Item 1: Functoriality**

Clear.

Item 2: Adjointness IThis follows from **Item 3** of **Proposition 4.2.1.9**.**Item 3: Adjointness II**For $- \triangleright Y$ to admit a right adjoint would require it to preserve colimits by ?? of ??. However, we have

$$\begin{aligned} \mathrm{pt} \triangleright X &\stackrel{\mathrm{def}}{=} |\mathrm{pt}| \odot X \\ &\cong X \\ &\neq \mathrm{pt}, \end{aligned}$$

and thus we see that $- \triangleright Y$ does not have a right adjoint.**Item 4: Adjointness III**This follows from **Item 2** of **Proposition 4.2.1.9**. 

00EX

REMARK 4.4.1.9 ► ON THE FAILURE OF $- \triangleright Y$ TO BE A LEFT ADJOINTHere is some intuition on why $- \triangleright Y$ fails to be a left adjoint. **Item 4** of **Proposition 4.3.1.7** states that we have a natural bijection

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleright Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}}(|X|, \mathbf{Sets}_*(Y, Z)),$$

so it would be reasonable to wonder whether a natural bijection of the form

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleright Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)),$$

also holds, which would give $- \triangleright Y \dashv \mathbf{Sets}_*(Y, -)$. However, such a bijection would require every map

$$f: X \triangleright Y \rightarrow Z$$

to satisfy

$$f(x_0 \triangleright y) = z_0$$

for each $x \in X$, whereas we are imposing such a basepoint preservation condition only for elements of the form $x \triangleright y_0$. Thus $\mathbf{Sets}_*(Y, -)$ can't be a right adjoint for $- \triangleright Y$, and as shown by [Item 3 of Proposition 4.4.1.7](#), no functor can.¹

¹The functor $\mathbf{Sets}_*(Y, -)$ is instead right adjoint to $- \wedge Y$, the smash product of pointed sets of [Definition 4.5.1.1](#). See [Item 2 of Proposition 4.5.1.10](#).

00EY 4.4.2 The Right Internal Hom of Pointed Sets

Let (X, x_0) and (Y, y_0) be pointed sets.

00EZ

DEFINITION 4.4.2.1 ► THE RIGHT INTERNAL HOM OF POINTED SETS

The **right internal Hom of pointed sets** is the functor

$$[-, -]_{\mathbf{Sets}_*}^{\triangleright} : \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$


defined as the composition

$$\mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \xrightarrow{\overline{\omega} \times \text{id}} \mathbf{Sets}^{\text{op}} \times \mathbf{Sets}_* \xrightarrow{\pitchfork} \mathbf{Sets}_*,$$

where:

- $\overline{\omega} : \mathbf{Sets}_* \rightarrow \mathbf{Sets}$ is the forgetful functor from pointed sets to sets.
- $\pitchfork : \mathbf{Sets}^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$ is the cotensor functor of [Item 1 of Proposition 4.2.2.6](#).

PROOF 4.4.2.2 ► PROOF OF DEFINITION 4.4.2.1

For a proof that $[-, -]_{\mathbf{Sets}_*}^{\triangleright}$ is indeed the right internal Hom of \mathbf{Sets}_* with respect to the right tensor product of pointed sets, see [Item 2 of Proposition 4.4.1.7](#). 

00F0	REMARK 4.4.2.3 ▶ UNWINDING DEFINITION 4.4.2.1, I: COMPARISON WITH $[-, -]_{\mathbf{Sets}_*}^{\triangleleft}$
	<p>We have</p> $[-, -]_{\mathbf{Sets}_*}^{\triangleleft} = [-, -]_{\mathbf{Sets}_*}^{\triangleright}.$
00F1	REMARK 4.4.2.4 ▶ UNWINDING DEFINITION 4.4.2.1, II: UNIVERSAL PROPERTY
	<p>The right internal Hom of pointed sets satisfies the following universal property:</p> $\mathbf{Sets}_*(X \triangleright Y, Z) \cong \mathbf{Sets}_*(Y, [X, Z]_{\mathbf{Sets}_*}^{\triangleright})$ <p>That is to say, the following data are in bijection:</p> <ol style="list-style-type: none"> 1. Pointed maps $f: X \triangleright Y \rightarrow Z$. 2. Pointed maps $f: Y \rightarrow [X, Z]_{\mathbf{Sets}_*}^{\triangleright}$.
00F2	REMARK 4.4.2.5 ▶ UNWINDING DEFINITION 4.4.2.1, III: EXPLICIT DESCRIPTION
	<p>In detail, the right internal Hom of (X, x_0) and (Y, y_0) is the pointed set $([X, Y]_{\mathbf{Sets}_*}^{\triangleright}, [(y_0)_{x \in X}])$ consisting of</p> <ul style="list-style-type: none"> • <i>The Underlying Set.</i> The set $[X, Y]_{\mathbf{Sets}_*}^{\triangleright}$ defined by $[X, Y]_{\mathbf{Sets}_*}^{\triangleright} \stackrel{\text{def}}{=} X \pitchfork Y \cong \bigwedge_{x \in X} (Y, y_0),$ where X denotes the underlying set of (X, x_0); • <i>The Underlying Basepoint.</i> The point $[(y_0)_{x \in X}]$ of $\bigwedge_{x \in X} (Y, y_0)$.
00F3	PROPOSITION 4.4.2.6 ▶ PROPERTIES OF RIGHT INTERNAL HOMS OF POINTED SETS
	<p>Let (X, x_0) and (Y, y_0) be pointed sets.</p>
00F4	<ol style="list-style-type: none"> 1. <i>Functoriality.</i> The assignments $X, Y, (X, Y) \mapsto [X, Y]_{\mathbf{Sets}_*}^{\triangleright}$ define func-

tors

$$\begin{aligned} [X, -]_{\mathbf{Sets}_*}^{\triangleright} &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ [-, Y]_{\mathbf{Sets}_*}^{\triangleright} &: \mathbf{Sets}_*^{\text{op}} \rightarrow \mathbf{Sets}_*, \\ [-_1, -_2]_{\mathbf{Sets}_*}^{\triangleright} &: \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f &: (X, x_0) \rightarrow (A, a_0), \\ g &: (Y, y_0) \rightarrow (B, b_0), \end{aligned}$$

the induced map

$$[f, g]_{\mathbf{Sets}_*}^{\triangleright} : [A, Y]_{\mathbf{Sets}_*}^{\triangleright} \rightarrow [X, B]_{\mathbf{Sets}_*}^{\triangleright}$$

is given by

$$[f, g]_{\mathbf{Sets}_*}^{\triangleright} \left([(y_a)_{a \in A}] \right) \stackrel{\text{def}}{=} [(g(y_{f(x)}))_{x \in X}]$$

for each $[(y_a)_{a \in A}] \in [A, Y]_{\mathbf{Sets}_*}^{\triangleright}$.

00F5

2. *Adjointness I.* We have an adjunction

$$\left(X \triangleright - \dashv [X, -]_{\mathbf{Sets}_*}^{\triangleright} \right) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \triangleright -} \\ \perp \\ \xleftarrow{[X, -]_{\mathbf{Sets}_*}^{\triangleright}} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleright Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(Y, [X, Z]_{\mathbf{Sets}_*}^{\triangleright})$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$, where $[X, Y]_{\mathbf{Sets}_*}^{\triangleright}$ is the pointed set of [Definition 4.4.2.1](#).

00F6

3. *Adjointness II.* The functor


$$- \triangleright Y : \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

does not admit a right adjoint.

PROOF 4.4.2.7 ▶ PROOF OF PROPOSITION 4.4.2.6

Item 1: Functoriality
 Clear.

Item 2: Adjointness I
 This is a repetition of **Item 2** of **Proposition 4.4.1.7**, and is proved there.

Item 3: Adjointness II
 This is a repetition of **Item 3** of **Proposition 4.4.1.7**, and is proved there. 

00F7 4.4.3 The Right Skew Unit

00F8 DEFINITION 4.4.3.1 ▶ THE RIGHT SKEW UNIT OF ▷

The **right skew unit of the right tensor product of pointed sets** is the functor

$$\mathbb{1}^{\text{Sets}_*, \triangleright} : \text{pt} \rightarrow \text{Sets}_*$$

defined by

$$\mathbb{1}_{\text{Sets}_*}^{\triangleright} \stackrel{\text{def}}{=} S^0.$$

00F9 4.4.4 The Right Skew Associator

00FA DEFINITION 4.4.4.1 ▶ THE RIGHT SKEW ASSOCIATOR OF ▷

The **skew associator of the right tensor product of pointed sets** is the natural transformation

$$\alpha^{\text{Sets}_*, \triangleright} : \triangleright \circ (\text{id}_{\text{Sets}_*} \times \triangleright) \implies \triangleright \circ (\triangleright \times \text{id}_{\text{Sets}_*}) \circ \alpha_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}^{\text{Cats}, -1}$$

as in the diagram

whose component

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright} : X \triangleright (Y \triangleright Z) \rightarrow (X \triangleright Y) \triangleright Z$$

at $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ is given by


$$\begin{aligned} X \triangleright (Y \triangleright Z) &\stackrel{\text{def}}{=} |X| \odot (Y \triangleright Z) \\ &\stackrel{\text{def}}{=} |X| \odot (|Y| \odot Z) \\ &\cong \bigvee_{x \in X} (|Y| \odot Z) \\ &\cong \bigvee_{x \in X} \left(\bigvee_{y \in Y} Z \right) \\ &\rightarrow \bigvee_{[(x,y)] \in \bigvee_{x \in X} Y} Z \\ &\cong \bigvee_{[(x,y)] \in |X| \odot Y} Z \\ &\cong ||X| \odot Y| \odot Z \\ &\stackrel{\text{def}}{=} |X \triangleright Y| \odot Z \\ &\stackrel{\text{def}}{=} (X \triangleright Y) \triangleright Z, \end{aligned}$$

where the map

$$\bigvee_{x \in X} \left(\bigvee_{y \in Y} Z \right) \rightarrow \bigvee_{[(x,y)] \in \bigvee_{x \in X} Y} Z$$

is given by $[(x, [(y, z)])] \mapsto [([(x, y)], z)]$.

PROOF 4.4.4.2 ► PROOF OF DEFINITION 4.4.4.1

(Proven below in a bit.) 

00FB

REMARK 4.4.4.3 ► UNWINDING DEFINITION 4.4.4.1

Unwinding the notation for elements, we have

$$\begin{aligned} [(x, [(y, z)])] &\stackrel{\text{def}}{=} [(x, y \triangleright z)] \\ &\stackrel{\text{def}}{=} x \triangleright (y \triangleright z) \end{aligned}$$

and

$$\begin{aligned} [([x, y], z)] &\stackrel{\text{def}}{=} [(x \triangleright y, z)] \\ &\stackrel{\text{def}}{=} (x \triangleright y) \triangleright z. \end{aligned}$$

So, in other words, $\alpha_{X,Y,Z}^{\text{Sets}_{*,\triangleright}}$ acts on elements via

$$\alpha_{X,Y,Z}^{\text{Sets}_{*,\triangleright}}(x \triangleright (y \triangleright z)) \stackrel{\text{def}}{=} (x \triangleright y) \triangleright z$$

for each $x \triangleright (y \triangleright z) \in X \triangleright (Y \triangleright Z)$.

00FC

REMARK 4.4.4.4 ▶ NON-INVERTIBILITY OF THE SKEW ASSOCIATOR OF \triangleright

Taking $y = y_0$, we see that the morphism $\alpha_{X,Y,Z}^{\text{Sets}_{*,\triangleright}}$ acts on elements as

$$\alpha_{X,Y,Z}^{\text{Sets}_{*,\triangleright}}(x \triangleright (y_0 \triangleright z)) \stackrel{\text{def}}{=} (x \triangleright y_0) \triangleright z.$$

However, by the definition of \triangleright , we have $x \triangleright y_0 = x' \triangleright y_0$ for all $x, x' \in X$, preventing $\alpha_{X,Y,Z}^{\text{Sets}_{*,\triangleright}}$ from being non-invertible.

PROOF 4.4.4.5 ▶ PROOF OF DEFINITION 4.4.4.1

Firstly, note that, given $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_{*})$, the map

$$\alpha_{X,Y,Z}^{\text{Sets}_{*,\triangleright}} : X \triangleright (Y \triangleright Z) \rightarrow (X \triangleright Y) \triangleright Z$$

is indeed a morphism of pointed sets, as we have

$$\alpha_{X,Y,Z}^{\text{Sets}_{*,\triangleright}}(x_0 \triangleright (y_0 \triangleright z_0)) = (x_0 \triangleright y_0) \triangleright z_0.$$

Next, we claim that $\alpha^{\text{Sets}_{*,\triangleright}}$ is a natural transformation. We need to show that, given morphisms of pointed sets

$$\begin{aligned} f &: (X, x_0) \rightarrow (X', x'_0), \\ g &: (Y, y_0) \rightarrow (Y', y'_0), \\ h &: (Z, z_0) \rightarrow (Z', z'_0) \end{aligned}$$

the diagram

$$\begin{array}{ccc}
 X \triangleright (Y \triangleright Z) & \xrightarrow{f \triangleright (g \triangleright h)} & X' \triangleright (Y' \triangleright Z') \\
 \alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright} \downarrow & & \downarrow \alpha_{X',Y',Z'}^{\text{Sets}_*, \triangleright} \\
 (X \triangleright Y) \triangleright Z & \xrightarrow{(f \triangleright g) \triangleright h} & (X' \triangleright Y') \triangleright Z'
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleright (y \triangleright z) & \longmapsto & f(x) \triangleright (g(y) \triangleright h(z)) \\
 \downarrow & & \downarrow \\
 (x \triangleright y) \triangleright z & \longmapsto & (f(x) \triangleright g(y)) \triangleright h(z)
 \end{array}$$

and hence indeed commutes, showing $\alpha^{\text{Sets}_*, \triangleright}$ to be a natural transformation. This finishes the proof. ▢

00FD 4.4.5 The Right Skew Left Unitor

00FE DEFINITION 4.4.5.1 ▶ THE RIGHT SKEW LEFT UNITOR OF \triangleright

The **skew left unitor of the right tensor product of pointed sets** is the natural transformation

$$\lambda^{\text{Sets}_*, \triangleright} : \lambda_{\text{Sets}_*}^{\text{Cats}_2} \xrightarrow{\sim} \triangleright \circ (\mathbb{1}_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*})$$

whose component


$$\lambda_X^{\text{Sets}_*, \triangleright} : X \rightarrow S^0 \triangleright X$$

at $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ is given by the composition

$$\begin{aligned} X &\rightarrow X \vee X \\ &\cong |S^0| \odot X \\ &\cong S^0 \triangleright X, \end{aligned}$$

where $X \rightarrow X \vee X$ is the map sending X to the second factor of X in $X \vee X$.

PROOF 4.4.5.2 ▶ PROOF OF DEFINITION 4.4.5.1

(Proven below in a bit.) 

00FF REMARK 4.4.5.3 ▶ UNWINDING DEFINITION 4.4.5.1

In other words, $\lambda_X^{\text{Sets}_*, \triangleright}$ acts on elements as

$$\lambda_X^{\text{Sets}_*, \triangleright}(x) \stackrel{\text{def}}{=} [(1, x)]$$

i.e. by

$$\lambda_X^{\text{Sets}_*, \triangleright}(x) \stackrel{\text{def}}{=} 1 \triangleright x$$

for each $x \in X$.

00FG REMARK 4.4.5.4 ▶ NON-INVERTIBILITY OF THE SKEW LEFT UNITOR OF \triangleright

The morphism $\lambda_X^{\text{Sets}_*, \triangleright}$ is non-invertible, as it is non-surjective when viewed as a map of sets, since the elements $0 \triangleright x$ of $S^0 \triangleright X$ with $x \neq x_0$ are outside the image of $\lambda_X^{\text{Sets}_*, \triangleright}$, which sends x to $1 \triangleright x$.

PROOF 4.4.5.5 ▶ PROOF OF DEFINITION 4.4.5.1

Firstly, note that, given $(X, x_0) \in \text{Obj}(\text{Sets}_*)$, the map

$$\lambda_X^{\text{Sets}_*, \triangleright} : X \rightarrow S^0 \triangleright X$$

is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \lambda_X^{\text{Sets}_*, \triangleright}(x_0) &= 1 \triangleright x_0 \\ &= 0 \triangleright x_0. \end{aligned}$$

Next, we claim that $\lambda^{\text{Sets}_*, \triangleright}$ is a natural transformation. We need to show that, given a morphism of pointed sets

$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc}
 X & \xrightarrow{f} & Y \\
 \lambda_X^{\text{Sets}_*, \triangleright} \downarrow & & \downarrow \lambda_Y^{\text{Sets}_*, \triangleright} \\
 S^0 \triangleright X & \xrightarrow{\text{id}_{S^0} \triangleright f} & S^0 \triangleright Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x & \longmapsto & f(x) \\
 \downarrow & & \downarrow \\
 1 \triangleright x & \longmapsto & 1 \triangleright f(x)
 \end{array}$$

and hence indeed commutes, showing $\lambda^{\text{Sets}_*, \triangleright}$ to be a natural transformation. This finishes the proof. ▢

00FH 4.4.6 The Right Skew Right Unitor

00FJ DEFINITION 4.4.6.1 ▶ THE RIGHT SKEW RIGHT UNITOR OF \triangleright

The **skew right unitor of the right tensor product of pointed sets** is the natural transformation

$$\rho^{\text{Sets}_*, \triangleright} : \triangleright \circ (\text{id} \times \mathbb{1}^{\text{Sets}_*}) \xrightarrow{\sim} \rho_{\text{Sets}_*}^{\text{Cats}_2}$$

whose component

$$\rho_X^{\text{Sets}_*, \triangleright} : X \triangleright S^0 \rightarrow X$$


at $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ is given by the composition

$$\begin{aligned} X \triangleright S^0 &\cong |X| \odot S^0 \\ &\cong \bigvee_{x \in X} S^0 \\ &\rightarrow X, \end{aligned}$$

where $\bigvee_{x \in X} S^0 \rightarrow X$ is the map given by

$$\begin{aligned} [(x, 0)] &\mapsto x_0, \\ [(x, 1)] &\mapsto x. \end{aligned}$$

PROOF 4.4.6.2 ▶ PROOF OF DEFINITION 4.4.6.1

(Proven below in a bit.) 

00FK

REMARK 4.4.6.3 ▶ UNWINDING DEFINITION 4.4.6.1

In other words, $\rho_X^{\text{Sets}_*, \triangleright}$ acts on elements as

$$\begin{aligned} \rho_X^{\text{Sets}_*, \triangleright}(x \triangleright 0) &\stackrel{\text{def}}{=} x_0, \\ \rho_X^{\text{Sets}_*, \triangleright}(x \triangleright 1) &\stackrel{\text{def}}{=} x \end{aligned}$$

for each $x \triangleright 1 \in X \triangleright S^0$.

00FL

REMARK 4.4.6.4 ▶ NON-INVERTIBILITY OF THE SKEW RIGHT UNITOR OF \triangleright

The morphism $\rho_X^{\text{Sets}_*, \triangleright}$ is almost invertible, with its would-be-inverse

$$\phi_X: X \rightarrow X \triangleright S^0$$

given by

$$\phi_X(x) \stackrel{\text{def}}{=} x \triangleright 1$$

for each $x \in X$. Indeed, we have

$$\begin{aligned} \left[\rho_X^{\text{Sets}_*, \triangleright} \circ \phi \right](x) &= \rho_X^{\text{Sets}_*, \triangleright}(\phi(x)) \\ &= \rho_X^{\text{Sets}_*, \triangleright}(x \triangleright 1) \\ &= x \\ &= [\text{id}_X](x) \end{aligned}$$

so that

$$\rho_X^{\text{Sets}_{*,\triangleright}} \circ \phi = \text{id}_X$$

and

$$\begin{aligned} [\phi \circ \rho_X^{\text{Sets}_{*,\triangleright}}](x \triangleright 1) &= \phi(\rho_X^{\text{Sets}_{*,\triangleright}}(x \triangleright 1)) \\ &= \phi(x) \\ &= x \triangleright 1 \\ &= [\text{id}_{X \triangleright S^0}](x \triangleright 1), \end{aligned}$$

but

$$\begin{aligned} [\phi \circ \rho_X^{\text{Sets}_{*,\triangleright}}](x \triangleright 0) &= \phi(\rho_X^{\text{Sets}_{*,\triangleright}}(x \triangleright 0)) \\ &= \phi(x_0) \\ &= 1 \triangleright x_0, \end{aligned}$$

where $x \triangleright 0 \neq 1 \triangleright x_0$. Thus

$$\phi \circ \rho_X^{\text{Sets}_{*,\triangleright}} \stackrel{?}{=} \text{id}_{X \triangleright S^0}$$

holds for all elements in $X \triangleright S^0$ except one.

PROOF 4.4.6.5 ► PROOF OF DEFINITION 4.4.6.1

Firstly, note that, given $(X, x_0) \in \text{Obj}(\text{Sets}_{*})$, the map

$$\rho_X^{\text{Sets}_{*,\triangleright}} : X \triangleright S^0 \rightarrow X$$

is indeed a morphism of pointed sets as we have

$$\rho_X^{\text{Sets}_{*,\triangleright}}(x_0 \triangleright 0) = x_0.$$

Next, we claim that $\rho^{\text{Sets}_{*,\triangleright}}$ is a natural transformation. We need to show that, given a morphism of pointed sets

$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram


$$\begin{array}{ccc} X \triangleright S^0 & \xrightarrow{f \triangleright \text{id}_{S^0}} & Y \triangleright S^0 \\ \rho_X^{\text{Sets}_{*,\triangleright}} \downarrow & & \downarrow \rho_Y^{\text{Sets}_{*,\triangleright}} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleright 0 & & x \triangleright 0 \mapsto f(x) \triangleright 0 \\
 \downarrow & & \downarrow \\
 x_0 \mapsto f(x_0) & & y_0
 \end{array}$$

and


$$\begin{array}{ccc}
 x \triangleright 1 \mapsto f(x) \triangleright 1 & & \\
 \downarrow & & \downarrow \\
 x \mapsto f(x) & &
 \end{array}$$

and hence indeed commutes, showing $\rho^{\text{Sets}_*, \triangleright}$ to be a natural transformation. This finishes the proof. 

00FM 4.4.7 The Diagonal

00FN DEFINITION 4.4.7.1 ▶ THE DIAGONAL OF \triangleright

The **diagonal of the right tensor product of pointed sets** is the natural transformation

$$\Delta^\triangleright : \text{id}_{\text{Sets}_*} \implies \triangleright \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2}$$


whose component

$$\Delta_X^\triangleright : (X, x_0) \rightarrow (X \triangleright X, x_0 \triangleright x_0)$$

at $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ is given by

$$\Delta_X^\triangleright(x) \stackrel{\text{def}}{=} x \triangleright x$$

for each $x \in X$.

PROOF 4.4.7.2 ► PROOF OF DEFINITION 4.4.7.1

Being a Morphism of Pointed Sets

We have

$$\Delta_X^\triangleright(x_0) \stackrel{\text{def}}{=} x_0 \triangleright x_0,$$

and thus Δ_X^\triangleright is a morphism of pointed sets.

Naturality

We need to show that, given a morphism of pointed sets


$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X^\triangleright \downarrow & & \downarrow \Delta_Y^\triangleright \\ X \triangleright X & \xrightarrow{f \triangleright f} & Y \triangleright Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \longmapsto & f(x) \\ \downarrow & & \downarrow \\ x \triangleright x & \longmapsto & f(x) \triangleright f(x) \end{array}$$

and hence indeed commutes, showing Δ^\triangleright to be natural. 

4.4.8 The Right Skew Monoidal Structure on Pointed Sets Associated to

00FP

►

PROPOSITION 4.4.8.1 ► THE RIGHT SKEW MONOIDAL STRUCTURE ON POINTED SETS ASSOCIATED TO ►

00FQ

The category Sets_* admits a right-closed right skew monoidal category structure consisting of

- *The Underlying Category.* The category Sets_* of pointed sets;
- *The Right Skew Monoidal Product.* The right tensor product functor

$$\triangleright : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of [Definition 4.4.1.1](#);

- *The Right Internal Skew Hom.* The right internal Hom functor

$$[-, -]_{\mathbf{Sets}_*}^{\triangleright} : \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

of [Definition 4.4.2.1](#);

- *The Right Skew Monoidal Unit.* The functor

$$\mathbb{1}^{\mathbf{Sets}_*, \triangleright} : \text{pt} \rightarrow \mathbf{Sets}_*$$

of [Definition 4.4.3.1](#);

- *The Right Skew Associators.* The natural transformation

$$\alpha^{\mathbf{Sets}_*, \triangleright} : \triangleright \circ (\text{id}_{\mathbf{Sets}_*} \times \triangleright) \Longrightarrow \triangleright \circ (\triangleright \times \text{id}_{\mathbf{Sets}_*}) \circ \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\text{Cats}_*, -1}$$

of [Definition 4.4.4.1](#);

- *The Right Skew Left Unitors.* The natural transformation

$$\lambda^{\mathbf{Sets}_*, \triangleright} : \lambda_{\mathbf{Sets}_*}^{\text{Cats}_2} \xrightarrow{\sim} \triangleright \circ (\mathbb{1}^{\mathbf{Sets}_*} \times \text{id}_{\mathbf{Sets}_*})$$

of [Definition 4.4.5.1](#);

- *The Right Skew Right Unitors.* The natural transformation

$$\rho^{\mathbf{Sets}_*, \triangleright} : \triangleright \circ (\text{id} \times \mathbb{1}^{\mathbf{Sets}_*}) \xrightarrow{\sim} \rho_{\mathbf{Sets}_*}^{\text{Cats}_2}$$


of [Definition 4.4.6.1](#).

PROOF 4.4.8.2 ► PROOF OF PROPOSITION 4.4.8.1


The Pentagon Identity

Let (W, w_0) , (X, x_0) , (Y, y_0) and (Z, z_0) be pointed sets. We have to show

that the diagram



commutes. Indeed, this diagram acts on elements as



and thus we see that the pentagon identity is satisfied.

The Right Skew Left Triangle Identity

Let (X, x_0) and (Y, y_0) be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleright Y & & \\
 \lambda_{X \triangleright Y}^{\text{Sets}_*, \triangleright} \downarrow & \searrow \lambda_X^{\text{Sets}_*, \triangleright} \triangleright \text{id}_Y & \\
 S^0 \triangleright (X \triangleright Y) & \xrightarrow{\alpha_{S^0, X, Y}^{\text{Sets}_*, \triangleright}} & (S^0 \triangleright X) \triangleright Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleright y & & \\
 \downarrow & \searrow & \\
 1 \triangleright (x \triangleright y) & \mapsto & (1 \triangleright x) \triangleright y
 \end{array}$$

and hence indeed commutes. Thus the left skew triangle identity is satisfied.

The Right Skew Right Triangle Identity

Let (X, x_0) and (Y, y_0) be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleright (Y \triangleright S^0) & \xrightarrow{\text{id}_X \triangleright \rho_Y^{\text{Sets}_*, \triangleright}} & (X \triangleright Y) \triangleright S^0 \\
 & \searrow \alpha_{S^0, X, Y}^{\text{Sets}_*, \triangleright} & \downarrow \rho_{X \triangleright Y}^{\text{Sets}_*, \triangleright} \\
 & & X \triangleright Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleright (y \triangleright 0) & \mapsto & (x \triangleright y) \triangleright 0 \\
 & \searrow & \downarrow \\
 & & x \triangleright y_0 = x_0 \triangleright y_0
 \end{array}$$

and

$$\begin{array}{ccc}
 x \triangleright (y \triangleright 1) & \mapsto & (x \triangleright y) \triangleright 1 \\
 & \searrow & \downarrow \\
 & & x \triangleright y
 \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

The Right Skew Middle Triangle Identity

Let (X, x_0) and (Y, y_0) be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleright Y & \xlongequal{\quad} & X \triangleright Y \\
 \text{id}_X \triangleright \lambda_Y^{\text{Sets}_*, \triangleright} \downarrow & & \uparrow \rho_X^{\text{Sets}_*, \triangleright} \triangleright \text{id}_Y \\
 X \triangleright (S^0 \triangleright Y) & \xrightarrow{\alpha_{X, S^0, Y}^{\text{Sets}_*, \triangleright}} & (X \triangleright S^0) \triangleright Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleright y & \xrightarrow{\quad} & x \triangleright y \\
 \downarrow & & \uparrow \\
 x \triangleright (1 \triangleright y) & \xrightarrow{\quad} & (x \triangleright 1) \triangleright y
 \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

The Zig-Zag Identity

We have to show that the diagram

$$\begin{array}{ccc}
 S^0 & \xrightarrow{\lambda_{S^0}^{\text{Sets}_*, \triangleright}} & S^0 \triangleright S^0 \\
 \searrow & & \downarrow \rho_{S^0}^{\text{Sets}_*, \triangleright} \\
 & & S^0
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 0 & \xrightarrow{\quad} & 1 \triangleright 0 \\
 \swarrow & & \downarrow \\
 & & 0
 \end{array}$$

and

$$\begin{array}{ccc}
 1 & \xrightarrow{\quad} & 1 \triangleright 1 \\
 \swarrow & & \downarrow \\
 & & 1
 \end{array}$$

and hence indeed commutes. Thus the zig-zag identity is satisfied.

Right Skew Monoidal Right-Closedness

This follows from **Item 2** of **Proposition 4.4.1.7**.



00FR 4.4.9 Monoids With Respect to the Right Tensor Product of Pointed Sets

00FS PROPOSITION 4.4.9.1 ► MONOIDS WITH RESPECT TO ▷

The category of monoids on $(\text{Sets}_*, \triangleright, S^0)$ is isomorphic to the category of “monoids with right zero”¹ and morphisms between them.

¹A monoid with right zero is defined similarly as the monoids with zero of ???. Succinctly, they are monoids (A, μ_A, η_A) with a special element 0_A satisfying

$$0_A a = 0_A$$

for each $a \in A$.

PROOF 4.4.9.2 ► PROOF OF PROPOSITION 4.4.9.1

Monoids on $(\text{Sets}_*, \triangleright, S^0)$

A monoid on $(\text{Sets}_*, \triangleright, S^0)$ consists of:

- *The Underlying Object.* A pointed set $(A, 0_A)$.
- *The Multiplication Morphism.* A morphism of pointed sets

$$\mu_A: A \triangleright A \rightarrow A,$$

determining a right bilinear morphism of pointed sets

$$\begin{aligned} A \times A &\longrightarrow A \\ (a, b) &\longmapsto ab. \end{aligned}$$


- *The Unit Morphism.* A morphism of pointed sets

$$\eta_A: S^0 \rightarrow A$$

picking an element 1_A of A .

satisfying the following conditions:

1. *Associativity.* The diagram



2. *Left Unitality*. The diagram

$$\begin{array}{ccc}
 A & \xrightarrow{\lambda_A^{\text{Sets}_{*,\triangleright}}} & S^0 \triangleright A \\
 \parallel & & \downarrow \eta_A \times \text{id}_A \\
 A & \xleftarrow{\mu_A} & A \triangleright A
 \end{array}$$

commutes.

3. *Right Unitality*. The diagram

$$\begin{array}{ccc}
 A \triangleright S^0 & \xrightarrow{\text{id}_A \times \eta_A} & A \triangleright A \\
 \searrow \rho_A^{\text{Sets}_{*,\triangleright}} & & \downarrow \mu_A \\
 & & A
 \end{array}$$

commutes.

Being a right-bilinear morphism of pointed sets, the multiplication map satisfies

$$0_A a = 0_A$$

for each $a \in A$. Now, the associativity, left unitality, and right unitality conditions act on elements as follows:

1. *Associativity*. The associativity condition acts as

$$\begin{array}{ccc}
 & & a \triangleright (b \triangleright c) \\
 & \swarrow & \searrow \\
 (a \triangleright b) \triangleright c & & (a \triangleright b) \triangleright c \quad a \triangleright bc \\
 \searrow & & \swarrow \\
 ab \triangleright c & \xrightarrow{\quad} & (ab)c \\
 & & \downarrow \\
 & & a(bc)
 \end{array}$$

This gives

$$(ab)c = a(bc)$$

for each $a, b, c \in A$.

2. *Left Unitality.* The left unitality condition acts as

$$\begin{array}{ccc}
 a & \xrightarrow{\quad} & 1 \triangleright a \\
 \downarrow & & \downarrow \\
 a & & 1_A a \longleftarrow 1_A \triangleright a
 \end{array}$$

This gives

$$1_A a = a$$

for each $a \in A$.

3. *Right Unitality.* The right unitality condition acts:

(a) On $1 \triangleright 0$ as

$$\begin{array}{ccc}
 1 \triangleright 0 & \xrightarrow{\quad} & a \triangleright 0 \\
 \searrow & & \searrow \quad \downarrow \\
 & & a 0_A
 \end{array}$$

(b) On $a \triangleright 1$ as

$$\begin{array}{ccc}
 a \triangleright 1 & \xrightarrow{\quad} & a \triangleright 1_A \\
 \searrow & & \searrow \quad \downarrow \\
 & & a 1_A
 \end{array}$$

This gives

$$\begin{aligned}
 a 1_A &= a, \\
 a 0_A &= 0_A
 \end{aligned}$$

for each $a \in A$.

Thus we see that monoids with respect to \triangleright are exactly monoids with right zero.

Morphisms of Monoids on $(\text{Sets}_*, \triangleright, S^0)$

A morphism of monoids on $(\text{Sets}_*, \triangleright, S^0)$ from $(A, \mu_A, \eta_A, 0_A)$ to $(B, \mu_B, \eta_B, 0_B)$ is a morphism of pointed sets

$$f: (A, 0_A) \rightarrow (B, 0_B)$$

satisfying the following conditions:

1. *Compatibility With the Multiplication Morphisms.* The diagram

$$\begin{array}{ccc}
 A \triangleright A & \xrightarrow{f \triangleright f} & B \triangleright B \\
 \mu_A \downarrow & & \downarrow \mu_B \\
 A & \xrightarrow{f} & B
 \end{array}$$

commutes.

2. *Compatibility With the Unit Morphisms.* The diagram

$$\begin{array}{ccc}
 S^0 & \xrightarrow{\eta_A} & A \\
 & \searrow \eta_B & \downarrow f \\
 & & B
 \end{array}$$

commutes.

These act on elements as

$$\begin{array}{ccc}
 a \triangleright b & & a \triangleright b \mapsto f(a) \triangleright f(b) \\
 \downarrow & & \downarrow \\
 ab \mapsto f(ab) & & f(a)f(b)
 \end{array}$$

and

$$\begin{array}{ccc}
 0 & \mapsto & 0_A \\
 \searrow & & \downarrow \\
 & & f(0_A)
 \end{array}$$

and

$$\begin{array}{ccc}
 1 & \mapsto & 1_A \\
 \searrow & & \downarrow \\
 & & f(1_A)
 \end{array}$$

giving

$$\begin{aligned} f(ab) &= f(a)f(b), \\ f(0_A) &= 0_B, \\ f(1_A) &= 1_B, \end{aligned}$$

for each $a, b \in A$, which is exactly a morphism of monoids with right zero.

Identities and Composition

Similarly, the identities and composition of $\text{Mon}(\text{Sets}_*, \triangleright, S^0)$ can be easily seen to agree with those of monoids with right zero, which finishes the proof.



00FT 4.5 The Smash Product of Pointed Sets

00FU 4.5.1 Foundations

Let (X, x_0) and (Y, y_0) be pointed sets.

00FV DEFINITION 4.5.1.1 ► SMASH PRODUCTS OF POINTED SETS

The **smash product of (X, x_0) and (Y, y_0)** ¹ is the pointed set $X \wedge Y$ ² satisfying the bijection

$$\text{Sets}_*(X \wedge Y, Z) \cong \text{Hom}_{\text{Sets}_*}^{\otimes}(X \times Y, Z),$$

naturally in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$.

¹*Further Terminology:* In the context of monoids with zero as models for \mathbb{F}_1 -algebras, the smash product $X \wedge Y$ is also called the **tensor product of \mathbb{F}_1 -modules of (X, x_0) and (Y, y_0)** or the **tensor product of (X, x_0) and (Y, y_0) over \mathbb{F}_1** .

²*Further Notation:* In the context of monoids with zero as models for \mathbb{F}_1 -algebras, the smash product $X \wedge Y$ is also denoted $X \otimes_{\mathbb{F}_1} Y$.

00FW REMARK 4.5.1.2 ► UNWINDING DEFINITION 4.5.1.1: THE UNIVERSAL PROPERTY I

That is to say, the smash product of pointed sets is defined so as to induce a bijection between the following data:

- Pointed maps $f: X \wedge Y \rightarrow Z$.

· Maps of sets $f: X \times Y \rightarrow Z$ satisfying

$$f(x_0, y) = z_0,$$

$$f(x, y_0) = z_0$$

for each $x \in X$ and each $y \in Y$.

00FX **REMARK 4.5.1.3 ► UNWINDING DEFINITION 4.5.1.1: THE UNIVERSAL PROPERTY II**

The smash product of pointed sets may be described as follows:

- The smash product of (X, x_0) and (Y, y_0) is the pair $((X \wedge Y, x_0 \wedge y_0), \iota)$ consisting of
 - A pointed set $(X \wedge Y, x_0 \wedge y_0)$;
 - A bilinear morphism of pointed sets $\iota: (X \times Y, (x_0, y_0)) \rightarrow X \wedge Y$;

satisfying the following universal property:

(UP) Given another such pair $((Z, z_0), f)$ consisting of

- * A pointed set (Z, z_0) ;
- * A bilinear morphism of pointed sets $f: (X \times Y, (x_0, y_0)) \rightarrow Z$;

there exists a unique morphism of pointed sets $X \wedge Y \xrightarrow{\exists!} Z$ making the diagram

$$\begin{array}{ccc}
 & X \wedge Y & \\
 \iota \nearrow & & \downarrow \exists! \\
 X \times Y & \xrightarrow{f} & Z
 \end{array}$$

commute.

00FY **CONSTRUCTION 4.5.1.4 ► SMASH PRODUCTS OF POINTED SETS**

Concretely, the **smash product of (X, x_0) and (Y, y_0)** is the pointed set $(X \wedge Y, x_0 \wedge y_0)$ consisting of

- *The Underlying Set.* The set $X \wedge Y$ defined by

$$X \wedge Y \cong (X \times Y) / \sim_R,$$

where \sim_R is the equivalence relation on $X \times Y$ obtained by declaring

$$\begin{aligned} (x_0, y) &\sim_R (x_0, y'), \\ (x, y_0) &\sim_R (x', y_0) \end{aligned}$$

for each $x, x' \in X$ and each $y, y' \in Y$;

- *The Basepoint.* The element $[(x_0, y_0)]$ of $X \wedge Y$ given by the equivalence class of (x_0, y_0) under the equivalence relation \sim on $X \times Y$.

PROOF 4.5.1.5 ► PROOF OF CONSTRUCTION 4.5.1.4

By [Item 6](#) of [Proposition 7.5.2.3](#), we have a natural bijection

$$\text{Sets}_*(X \wedge Y, Z) \cong \text{Hom}_{\text{Sets}}^R(X \times Y, Z).$$

Now, by definition, $\text{Hom}_{\text{Sets}}^R(X \times Y, Z)$ is the set

$$\text{Hom}_{\text{Sets}}^R(X \times Y, Z) \stackrel{\text{def}}{=} \left\{ f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \left| \begin{array}{l} \text{for each } x, y \in X, \text{ if} \\ (x, y) \sim_R (x', y'), \text{ then} \\ f(x, y) = f(x', y') \end{array} \right. \right\}.$$

However, the condition $(x, y) \sim_R (x', y')$ only holds when:

1. We have $x = x'$ and $y = y'$.
2. The following conditions are satisfied:
 - (a) We have $x = x_0$ or $y = y_0$.
 - (b) We have $x' = x_0$ or $y' = y_0$.

So, given $f \in \text{Hom}_{\text{Sets}}(X \times Y, Z)$ with a corresponding $\bar{f}: X \wedge Y \rightarrow Z$, the latter case above implies

$$\begin{aligned} f(x_0, y) &= f(x, y_0) \\ &= f(x_0, y_0), \end{aligned}$$

and since $\bar{f}: X \wedge Y \rightarrow Z$ is a pointed map, we have

$$\begin{aligned} f(x_0, y_0) &= \bar{f}(x_0, y_0) \\ &= z_0. \end{aligned}$$

Thus the elements f in $\text{Hom}_{\text{Sets}}(X \times Y, Z)$ are precisely those functions $f: X \times Y \rightarrow Z$ satisfying the equalities


$$\begin{aligned} f(x_0, y) &= z_0, \\ f(x, y_0) &= z_0 \end{aligned}$$

for each $x \in X$ and each $y \in Y$, giving an equality

$$\text{Hom}_{\text{Sets}}^R(X \times Y, Z) = \text{Hom}_{\text{Sets}_*}^\otimes(X \times Y, Z)$$

of sets, which when composed with our earlier isomorphism

$$\text{Sets}_*(X \wedge Y, Z) \cong \text{Hom}_{\text{Sets}}^R(X \times Y, Z)$$

gives our desired natural bijection, finishing the proof. 

00FZ REMARK 4.5.1.6 ► ON THE CONSTRUCTION OF THE SMASH PRODUCT OF POINTED SETS

It is also somewhat common to write

$$X \wedge Y \stackrel{\text{def}}{=} \frac{X \times Y}{X \vee Y},$$

identifying $X \vee Y$ with the subspace $(\{x_0\} \times Y) \cup (X \times \{y_0\})$ of $X \times Y$, and having the quotient be defined by declaring $(x, y) \sim (x', y')$ iff we have $(x, y), (x', y') \in X \vee Y$.

00G0 NOTATION 4.5.1.7 ► ELEMENTS OF SMASH PRODUCTS OF POINTED SETS

We write $x \wedge y$ for the element $[(x, y)]$ of

$$X \wedge Y \cong X \times Y / \sim.$$

00G1 REMARK 4.5.1.8 ► BASEPOINTS OF SMASH PRODUCTS OF POINTED SETS

Employing the notation introduced in [Notation 4.5.1.7](#), we have

$$\begin{aligned} x_0 \wedge y_0 &= x \wedge y_0, \\ &= x_0 \wedge y \end{aligned}$$

for each $x \in X$ and each $y \in Y$, and

$$x \wedge y_0 = x' \wedge y_0,$$

$$x_0 \wedge y = x_0 \wedge y'$$

for each $x, x' \in X$ and each $y, y' \in Y$.

00G2 **EXAMPLE 4.5.1.9 ► EXAMPLES OF SMASH PRODUCTS OF POINTED SETS**

Here are some examples of smash products of pointed sets.

- 00G3 1. *Smashing With pt.* For any pointed set X , we have isomorphisms of pointed sets

$$\text{pt} \wedge X \cong \text{pt},$$

$$X \wedge \text{pt} \cong \text{pt}.$$

- 00G4 2. *Smashing With S^0 .* For any pointed set X , we have isomorphisms of pointed sets

$$S^0 \wedge X \cong X,$$

$$X \wedge S^0 \cong X.$$

00G5 **PROPOSITION 4.5.1.10 ► PROPERTIES OF SMASH PRODUCTS OF POINTED SETS**

Let (X, x_0) and (Y, y_0) be pointed sets.

- 00G6 1. *Functoriality.* The assignments $X, Y, (X, Y) \mapsto X \wedge Y$ define functors

$$X \wedge -: \text{Sets}_* \rightarrow \text{Sets}_*,$$

$$- \wedge Y: \text{Sets}_* \rightarrow \text{Sets}_*,$$

$$-_1 \wedge -_2: \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*.$$

In particular, given pointed maps

$$f: (X, x_0) \rightarrow (A, a_0),$$

$$g: (Y, y_0) \rightarrow (B, b_0),$$

the induced map

$$f \wedge g: X \wedge Y \rightarrow A \wedge B$$

is given by

$$[f \wedge g](x \wedge y) \stackrel{\text{def}}{=} f(x) \wedge g(y)$$

for each $x \wedge y \in X \wedge Y$.

00G7

2. *Adjointness.* We have adjunctions

$$(X \wedge - \dashv \mathbf{Sets}_*(X, -)) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*$$

$$(- \wedge Y \dashv \mathbf{Sets}_*(Y, -)) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(Y, -)} \end{array} \mathbf{Sets}_*$$

witnessed by bijections

$$\text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)),$$

$$\text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(A, Z)),$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$.

00G8

3. *Enriched Adjointness.* We have \mathbf{Sets}_* -enriched adjunctions

$$(X \wedge - \dashv \mathbf{Sets}_*(X, -)) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*$$

$$(- \wedge Y \dashv \mathbf{Sets}_*(Y, -)) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(Y, -)} \end{array} \mathbf{Sets}_*$$

witnessed by isomorphisms of pointed sets

$$\mathbf{Sets}_*(X \wedge Y, Z) \cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(Y, Z)),$$

$$\mathbf{Sets}_*(X \wedge Y, Z) \cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(A, Z)),$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$.

00G9

4. *As a Pushout.* We have an isomorphism

$$X \wedge Y \cong \text{pt} \coprod_{X \vee Y} (X \times Y),$$

$$\begin{array}{ccc} X \wedge Y & \longleftarrow & X \times Y \\ \uparrow \ulcorner & & \uparrow \lrcorner \\ \text{pt} & \longleftarrow & X \vee Y \end{array}$$

00GA

natural in $X, Y \in \text{Obj}(\text{Sets}_*)$, where the pushout is taken in Sets , and the embedding $\iota: X \vee Y \hookrightarrow X \times Y$ is defined following [Remark 4.5.1.6](#).

5. *Distributivity Over Wedge Sums*. We have isomorphisms of pointed sets

$$X \wedge (Y \vee Z) \cong (X \wedge Y) \vee (X \wedge Z),$$

$$(X \vee Y) \wedge Z \cong (X \wedge Z) \vee (Y \wedge Z),$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$.

PROOF 4.5.1.11 ► PROOF OF PROPOSITION 4.5.1.10

Item 1: Functoriality

The map $f \wedge g$ comes from [Item 4 of Proposition 7.5.2.3](#) via the map

$$f \wedge g: X \times Y \rightarrow A \wedge B$$

sending (x, y) to $f(x) \wedge g(y)$, which we need to show satisfies

$$[f \wedge g](x, y) = [f \wedge g](x', y')$$

for each $(x, y), (x', y') \in X \times Y$ with $(x, y) \sim_R (x', y')$, where \sim_R is the relation constructing $X \wedge Y$ as

$$X \wedge Y \cong (X \times Y) / \sim_R$$

in [Construction 4.5.1.4](#). The condition defining \sim is that at least one of the following conditions is satisfied:

1. We have $x = x'$ and $y = y'$;
2. Both of the following conditions are satisfied:
 - (a) We have $x = x_0$ or $y = y_0$.
 - (b) We have $x' = x_0$ or $y' = y_0$.

We have five cases:

1. In the first case, we clearly have

$$[f \wedge g](x, y) = [f \wedge g](x', y')$$

since $x = x'$ and $y = y'$.

2. If $x = x_0$ and $x' = x_0$, we have

$$\begin{aligned} [f \wedge g](x_0, y) &\stackrel{\text{def}}{=} f(x_0) \wedge g(y) \\ &= a_0 \wedge g(y) \\ &= a_0 \wedge g(y') \\ &= f(x_0) \wedge g(y') \\ &\stackrel{\text{def}}{=} [f \wedge g](x_0, y'). \end{aligned}$$

3. If $x = x_0$ and $y' = y_0$, we have

$$\begin{aligned} [f \wedge g](x_0, y) &\stackrel{\text{def}}{=} f(x_0) \wedge g(y) \\ &= a_0 \wedge g(y) \\ &= a_0 \wedge b_0 \\ &= f(x') \wedge b_0 \\ &= f(x') \wedge g(y_0) \\ &\stackrel{\text{def}}{=} [f \wedge g](x', y_0). \end{aligned}$$

4. If $y = y_0$ and $x' = x_0$, we have

$$\begin{aligned} [f \wedge g](x, y_0) &\stackrel{\text{def}}{=} f(x) \wedge g(y_0) \\ &= f(x) \wedge b_0 \\ &= a_0 \wedge b_0 \\ &= a_0 \wedge g(y') \\ &= f(x_0) \wedge g(y') \\ &\stackrel{\text{def}}{=} [f \wedge g](x_0, y'). \end{aligned}$$

5. If $y = y_0$ and $y' = y_0$, we have

$$\begin{aligned} [f \wedge g](x, y_0) &\stackrel{\text{def}}{=} f(x) \wedge g(y_0) \\ &= f(x) \wedge b_0 \\ &= f(x') \wedge b_0 \\ &= f(x) \wedge g(y_0) \\ &\stackrel{\text{def}}{=} [f \wedge g](x', y_0). \end{aligned}$$

Thus $f \wedge g$ is well-defined. Next, we claim that \wedge preserves identities and composition:

• *Preservation of Identities.* We have

$$\begin{aligned} [\text{id}_X \wedge \text{id}_Y](x \wedge y) &\stackrel{\text{def}}{=} \text{id}_X(x) \wedge \text{id}_Y(y) \\ &= x \wedge y \\ &= [\text{id}_{X \wedge Y}](x \wedge y) \end{aligned}$$

for each $x \wedge y \in X \wedge Y$, and thus

$$\text{id}_X \wedge \text{id}_Y = \text{id}_{X \wedge Y}.$$

• *Preservation of Composition.* Given pointed maps

$$\begin{aligned} f: (X, x_0) &\rightarrow (X', x'_0), \\ h: (X', x'_0) &\rightarrow (X'', x''_0), \\ g: (Y, y_0) &\rightarrow (Y', y'_0), \\ k: (Y', y'_0) &\rightarrow (Y'', y''_0), \end{aligned}$$

we have

$$\begin{aligned} [(h \circ f) \wedge (k \circ g)](x \wedge y) &\stackrel{\text{def}}{=} h(f(x)) \wedge k(g(y)) \\ &\stackrel{\text{def}}{=} [h \wedge k](f(x) \wedge g(y)) \\ &\stackrel{\text{def}}{=} [h \wedge k]([f \wedge g](x \wedge y)) \\ &\stackrel{\text{def}}{=} [(h \wedge k) \circ (f \wedge g)](x \wedge y) \end{aligned}$$

for each $x \wedge y \in X \wedge Y$, and thus

$$(h \circ f) \wedge (k \circ g) = (h \wedge k) \circ (f \wedge g).$$

This finishes the proof.

Item 2: Adjointness

We prove only the adjunction $- \wedge Y \dashv \mathbf{Sets}_*(Y, -)$, witnessed by a natural bijection

$$\text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)),$$

as the proof of the adjunction $X \wedge - \dashv \mathbf{Sets}_*(X, -)$ is similar. We claim we have a bijection

$$\text{Hom}_{\mathbf{Sets}_*}^{\otimes}(X \times Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z))$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$, implying the desired adjunction. Indeed, this bijection is a restriction of the bijection

$$\mathbf{Sets}(X \times Y, Z) \cong \mathbf{Sets}(X, \mathbf{Sets}(Y, Z))$$

of **Item 2** of **Proposition 2.1.3.3**:

- A map

$$\xi: X \times Y \rightarrow Z$$

in $\text{Hom}_{\mathbf{Sets}_*}^{\otimes}(X \times Y, Z)$ gets sent to the pointed map

$$\begin{aligned} \xi^\dagger: (X, x_0) &\rightarrow (\mathbf{Sets}_*(Y, Z), \Delta_{z_0}), \\ x &\longmapsto \left(\xi_x^\dagger: Y \rightarrow Z \right), \end{aligned}$$

where $\xi_x^\dagger: Y \rightarrow Z$ is the map defined by

$$\xi_x^\dagger(y) \stackrel{\text{def}}{=} \xi(x, y)$$

for each $y \in Y$, where:

- The map ξ^\dagger is indeed pointed, as we have

$$\begin{aligned} \xi_{x_0}^\dagger(y) &\stackrel{\text{def}}{=} \xi(x_0, y) \\ &\stackrel{\text{def}}{=} z_0 \end{aligned}$$

for each $y \in Y$. Thus $\xi_{x_0}^\dagger = \Delta_{z_0}$ and ξ^\dagger is pointed.

- The map ξ_x^\dagger indeed lies in $\mathbf{Sets}_*(Y, Z)$, as we have

$$\begin{aligned} \xi_x^\dagger(y_0) &\stackrel{\text{def}}{=} \xi(x, y_0) \\ &\stackrel{\text{def}}{=} z_0. \end{aligned}$$

- Conversely, a map

$$\begin{aligned} \xi: (X, x_0) &\rightarrow (\mathbf{Sets}_*(Y, Z), \Delta_{z_0}), \\ x &\longmapsto (\xi_x: Y \rightarrow Z), \end{aligned}$$

in $\text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z))$ gets sent to the map

$$\xi^\dagger: X \times Y \rightarrow Z$$

defined by

$$\xi^\dagger(x, y) \stackrel{\text{def}}{=} \xi_x(y)$$

for each $(x, y) \in X \times Y$, which indeed lies in $\text{Hom}_{\mathbf{Sets}_*}^{\otimes}(X \times Y, Z)$, as:

– *Left Bilinearity.* We have

$$\begin{aligned}\xi^\dagger(x_0, y) &\stackrel{\text{def}}{=} \xi_{x_0}(y) \\ &\stackrel{\text{def}}{=} \Delta_{z_0}(y) \\ &\stackrel{\text{def}}{=} z_0\end{aligned}$$

for each $y \in Y$, since $\xi_{x_0} = \Delta_{z_0}$ as ξ is assumed to be a pointed map.

– *Right Bilinearity.* We have

$$\begin{aligned}\xi^\dagger(x, y_0) &\stackrel{\text{def}}{=} \xi_x(y_0) \\ &\stackrel{\text{def}}{=} z_0\end{aligned}$$

for each $x \in X$, since $\xi_x \in \mathbf{Sets}_*(Y, Z)$ is a morphism of pointed sets.

This finishes the proof.

Item 3: Enriched Adjointness

This follows from [Item 2](#) and ?? of ??.

Item 4: As a Pushout

Following the description of [Remark 2.2.4.3](#), we have

$$\text{pt} \coprod_{X \vee Y} (X \times Y) \cong (\text{pt} \times (X \times Y)) / \sim,$$

where \sim identifies the element \star in pt with all elements of the form (x_0, y) and (x, y_0) in $X \times Y$. Thus [Item 4](#) of [Proposition 7.5.2.3](#) coupled with [Remark 4.5.1.8](#) then gives us a well-defined map


$$\text{pt} \coprod_{X \vee Y} (X \times Y) \rightarrow X \wedge Y$$

via $[(\star, (x, y))] \mapsto x \wedge y$, with inverse

$$X \wedge Y \rightarrow \text{pt} \coprod_{X \vee Y} (X \times Y)$$

given by $x \wedge y \mapsto [(\star, (x, y))]$.

Item 5: Distributivity Over Wedge Sums

This follows from [Proposition 4.5.9.1](#), ?? of ??, and the fact that \vee is the coproduct in \mathbf{Sets}_* ([Definition 3.3.3.1](#)). 

4.5.2 The Internal Hom of Pointed Sets

Let (X, x_0) and (Y, y_0) be pointed sets.



00GC

The **internal Hom**¹ of pointed sets from (X, x_0) to (Y, y_0) is the pointed set $\mathbf{Sets}_*((X, x_0), (Y, y_0))$ ² consisting of:

- *The Underlying Set.* The set $\mathbf{Sets}_*((X, x_0), (Y, y_0))$ of morphisms of pointed sets from (X, x_0) to (Y, y_0) .
- *The Basepoint.* The element

$$\Delta_{y_0} : (X, x_0) \rightarrow (Y, y_0)$$

of $\mathbf{Sets}_*((X, x_0), (Y, y_0))$ given by


$$\Delta_{y_0}(x) \stackrel{\text{def}}{=} y_0$$

for each $x \in X$.

¹The pointed set $\mathbf{Sets}_*(X, Y)$ is the internal **Hom** of \mathbf{Sets}_* with respect to the smash product of [Definition 4.5.1.1](#); see [Item 2](#) of [Proposition 4.5.1.10](#).

²*Further Notation:* Also written $\mathbf{Hom}_{\mathbf{Sets}_*}(X, Y)$.

PROOF 4.5.2.2 ► PROOF OF DEFINITION 4.5.2.1

For a proof that \mathbf{Sets}_* is indeed the internal Hom of \mathbf{Sets}_* with respect to the smash product of pointed sets, see [Item 2](#) of [Proposition 4.5.1.10](#). 

00GD PROPOSITION 4.5.2.3 ► PROPERTIES OF THE INTERNAL HOM OF POINTED SETS

Let (X, x_0) and (Y, y_0) be pointed sets.

- 00GE 1. *Functoriality.* The assignments $X, Y, (X, Y) \mapsto \mathbf{Sets}_*(X, Y)$ define functors

$$\mathbf{Sets}_*(X, -) : \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

$$\mathbf{Sets}_*(-, Y) : \mathbf{Sets}_*^{\text{op}} \rightarrow \mathbf{Sets}_*,$$

$$\mathbf{Sets}_*(-, -) : \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*.$$

In particular, given pointed maps

$$f : (X, x_0) \rightarrow (A, a_0),$$

$$g : (Y, y_0) \rightarrow (B, b_0),$$

the induced map

$$\mathbf{Sets}_*(f, g) : \mathbf{Sets}_*(A, Y) \rightarrow \mathbf{Sets}_*(X, B)$$

is given by

$$[\mathbf{Sets}_*(f, g)](\phi) \stackrel{\text{def}}{=} g \circ \phi \circ f$$

for each $\phi \in \mathbf{Sets}_*(A, Y)$.

00GF

2. *Adjointness.* We have adjunctions

$$(X \wedge - \dashv \mathbf{Sets}_*(X, -)) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*$$

$$(- \wedge Y \dashv \mathbf{Sets}_*(Y, -)) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(Y, -)} \end{array} \mathbf{Sets}_*$$

witnessed by bijections

$$\text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)),$$

$$\text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(A, Z)),$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$.

00GG

3. *Enriched Adjointness.* We have \mathbf{Sets}_* -enriched adjunctions

$$(X \wedge - \dashv \mathbf{Sets}_*(X, -)) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*$$

$$(- \wedge Y \dashv \mathbf{Sets}_*(Y, -)) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(Y, -)} \end{array} \mathbf{Sets}_*$$

witnessed by isomorphisms of pointed sets

$$\mathbf{Sets}_*(X \wedge Y, Z) \cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(Y, Z)),$$

$$\mathbf{Sets}_*(X \wedge Y, Z) \cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(A, Z)),$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$.

PROOF 4.5.2.4 ► PROOF OF PROPOSITION 4.5.2.3

Item 1: Functoriality

This follows from **Item 1** of **Proposition 2.3.5.2** and from the equalities

$$\begin{aligned} g \circ \Delta_{y_0} &= \Delta_{z_0}, \\ \Delta_{y_0} \circ f &= \Delta_{y_0} \end{aligned}$$

for morphisms $f: (K, k_0) \rightarrow (X, x_0)$ and $g: (Y, y_0) \rightarrow (Z, z_0)$, which guarantee pre- and postcomposition by morphisms of pointed sets to also be morphisms of pointed sets.

Item 2: Adjointness

This is a repetition of **Item 2** of **Proposition 4.5.1.10**, and is proved there.

Item 3: Enriched Adjointness

This is a repetition of **Item 3** of **Proposition 4.5.1.10**, and is proved there. 

00GH 4.5.3 The Monoidal Unit

00GJ DEFINITION 4.5.3.1 ► THE MONOIDAL UNIT OF \wedge

The **monoidal unit of the smash product of pointed sets** is the functor

$$\mathbb{1}^{\text{Sets}_*} : \text{pt} \rightarrow \text{Sets}_*$$

defined by

$$\mathbb{1}_{\text{Sets}_*} \stackrel{\text{def}}{=} S^0.$$


00GK 4.5.4 The Associator

00GL DEFINITION 4.5.4.1 ► THE ASSOCIATOR OF \wedge

The **associator of the smash product of pointed sets** is the natural isomorphism

$$\alpha^{\text{Sets}_*} : \wedge \circ (\wedge \times \text{id}_{\text{Sets}_*}) \xrightarrow{\sim} \wedge \circ (\text{id}_{\text{Sets}_*} \times \wedge) \circ \alpha_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}^{\text{Cats}}$$

as in the diagram



whose component

$$\alpha_{X,Y,Z}^{\text{Sets}_*}: (X \wedge Y) \wedge Z \xrightarrow{\cong} X \wedge (Y \wedge Z)$$

at $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$ is given by

$$\alpha_{X,Y,Z}^{\text{Sets}_*}((x \wedge y) \wedge z) \stackrel{\text{def}}{=} x \wedge (y \wedge z)$$

for each $(x \wedge y) \wedge z \in (X \wedge Y) \wedge Z$.

PROOF 4.5.4.2 ► PROOF OF DEFINITION 4.5.4.1

Well-Definedness

Let $[(x, y), z] = [(x', y'), z']$ be an element in $(X \wedge Y) \wedge Z$. Then either:

1. We have $x = x', y = y',$ and $z = z'$.
2. Both of the following conditions are satisfied:
 - (a) We have $x = x_0$ or $y = y_0$ or $z = z_0$.
 - (b) We have $x' = x_0$ or $y' = y_0$ or $z' = z_0$.

In the first case, $\alpha_{X,Y,Z}^{\text{Sets}_*}$ clearly sends both elements to the same element in $X \wedge (Y \wedge Z)$. Meanwhile, in the latter case both elements are equal to the basepoint $(x_0 \wedge y_0) \wedge z_0$ of $(X \wedge Y) \wedge Z$, which gets sent to the basepoint $x_0 \wedge (y_0 \wedge z_0)$ of $X \wedge (Y \wedge Z)$.

Being a Morphism of Pointed Sets

As just mentioned, we have

$$\alpha_{X,Y,Z}^{\text{Sets}_*}((x_0 \wedge y_0) \wedge z_0) \stackrel{\text{def}}{=} x_0 \wedge (y_0 \wedge z_0),$$

and thus $\alpha_{X,Y,Z}^{\text{Sets}_*}$ is a morphism of pointed sets.

Invertibility

Clearly, the inverse of $\alpha_{X,Y,Z}^{\text{Sets}_*}$ is given by the morphism

$$\alpha_{X,Y,Z}^{\text{Sets}_*, -1} : X \wedge (Y \wedge Z) \xrightarrow{\cong} (X \wedge Y) \wedge Z$$

defined by

$$\alpha_{X,Y,Z}^{\text{Sets}_*, -1}(x \wedge (y \wedge z)) \stackrel{\text{def}}{=} (x \wedge y) \wedge z$$

for each $x \wedge (y \wedge z) \in X \wedge (Y \wedge Z)$.

Naturality

We need to show that, given morphisms of pointed sets

$$f : (X, x_0) \rightarrow (X', x'_0),$$

$$g : (Y, y_0) \rightarrow (Y', y'_0),$$

$$h : (Z, z_0) \rightarrow (Z', z'_0)$$

the diagram


$$\begin{array}{ccc} (X \wedge Y) \wedge Z & \xrightarrow{(f \wedge g) \wedge h} & (X' \wedge Y') \wedge Z' \\ \alpha_{X,Y,Z}^{\text{Sets}_*} \downarrow & & \downarrow \alpha_{X',Y',Z'}^{\text{Sets}_*} \\ X \wedge (Y \wedge Z) & \xrightarrow{f \wedge (g \wedge h)} & X' \wedge (Y' \wedge Z') \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (x \wedge y) \wedge z & \longmapsto & (f(x) \wedge g(y)) \wedge h(z) \\ \downarrow & & \downarrow \\ x \wedge (y \wedge z) & \longmapsto & f(x) \wedge (g(y) \wedge h(z)) \end{array}$$

and hence indeed commutes, showing α^{Sets_*} to be a natural transformation.

Being a Natural Isomorphism


Since α^{Sets_*} is natural and $\alpha^{\text{Sets}_*, -1}$ is a componentwise inverse to α^{Sets_*} , it follows from [Item 2 of Proposition 8.8.6.2](#) that $\alpha^{\text{Sets}_*, -1}$ is also natural. Thus α^{Sets_*} is a natural isomorphism. 

00GM 4.5.5 The Left Unitor

00GN

DEFINITION 4.5.5.1 ► THE LEFT UNITOR OF \wedge

The **left unitor of the smash product of pointed sets** is the natural isomorphism



whose component

$$\lambda_X^{\text{Sets}_*} : S^0 \wedge X \xrightarrow{\cong} X$$

at $X \in \text{Obj}(\text{Sets}_*)$ is given by

$$\begin{aligned} 0 \wedge x &\mapsto x_0, \\ 1 \wedge x &\mapsto x. \end{aligned}$$

PROOF 4.5.5.2 ► PROOF OF DEFINITION 4.5.5.1

Well-Definedness

Let $[(x, y)] = [(x', y')]$ be an element in $S^0 \wedge X$. Then either:

1. We have $x = x'$ and $y = y'$.
2. Both of the following conditions are satisfied:
 - (a) We have $x = 0$ or $y = x_0$.
 - (b) We have $x' = 0$ or $y' = x_0$.

In the first case, $\lambda_X^{\text{Sets}_*}$ clearly sends both elements to the same element in X . Meanwhile, in the latter case both elements are equal to the basepoint $0 \wedge x_0$ of $S^0 \wedge X$, which gets sent to the basepoint x_0 of X .

Being a Morphism of Pointed Sets

As just mentioned, we have

$$\lambda_X^{\text{Sets}_*}(0 \wedge x_0) \stackrel{\text{def}}{=} x_0,$$

and thus $\lambda_X^{\text{Sets}_*}$ is a morphism of pointed sets.

Invertibility

The inverse of $\lambda_X^{\text{Sets}_*}$ is the morphism

$$\lambda_X^{\text{Sets}_*, -1}: X \xrightarrow{\cong} S^0 \wedge X$$

defined by

$$\lambda_X^{\text{Sets}_*, -1}(x) \stackrel{\text{def}}{=} 1 \wedge x$$

for each $x \in X$. Indeed:

· *Invertibility I.* We have

$$\begin{aligned} \left[\lambda_X^{\text{Sets}_*, -1} \circ \lambda_X^{\text{Sets}_*} \right] (0 \wedge x) &= \lambda_X^{\text{Sets}_*, -1} \left(\lambda_X^{\text{Sets}_*} (0 \wedge x) \right) \\ &= \lambda_X^{\text{Sets}_*, -1} (x_0) \\ &= 1 \wedge x_0 \\ &= 0 \wedge x, \end{aligned}$$

and

$$\begin{aligned} \left[\lambda_X^{\text{Sets}_*, -1} \circ \lambda_X^{\text{Sets}_*} \right] (1 \wedge x) &= \lambda_X^{\text{Sets}_*, -1} \left(\lambda_X^{\text{Sets}_*} (1 \wedge x) \right) \\ &= \lambda_X^{\text{Sets}_*, -1} (x) \\ &= 1 \wedge x \end{aligned}$$

for each $x \in X$, and thus we have

$$\lambda_X^{\text{Sets}_*, -1} \circ \lambda_X^{\text{Sets}_*} = \text{id}_{S^0 \wedge X}.$$

· *Invertibility II.* We have

$$\begin{aligned} \left[\lambda_X^{\text{Sets}_*} \circ \lambda_X^{\text{Sets}_*, -1} \right] (x) &= \lambda_X^{\text{Sets}_*} \left(\lambda_X^{\text{Sets}_*, -1} (x) \right) \\ &= \lambda_X^{\text{Sets}_*, -1} (1 \wedge x) \\ &= x \end{aligned}$$

for each $x \in X$, and thus we have

$$\lambda_X^{\text{Sets}_*} \circ \lambda_X^{\text{Sets}_*, -1} = \text{id}_X.$$

This shows $\lambda_X^{\text{Sets}_*}$ to be invertible.

Naturality

We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} S^0 \wedge X & \xrightarrow{\text{id}_{S^0} \wedge f} & S^0 \wedge Y \\ \lambda_X^{\text{Sets}_*} \downarrow & & \downarrow \lambda_Y^{\text{Sets}_*} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} 0 \wedge x & & 0 \wedge x \mapsto 0 \wedge f(x) \\ \downarrow & & \downarrow \\ x_0 & \mapsto & f(x_0) \\ & & y_0 \end{array}$$

and

$$\begin{array}{ccc} 1 \wedge x & \mapsto & 1 \wedge f(x) \\ \downarrow & & \downarrow \\ x & \mapsto & f(x) \end{array}$$

and hence indeed commutes, showing λ^{Sets_*} to be a natural transformation.

Being a Natural Isomorphism

Since λ^{Sets_*} is natural and $\lambda^{\text{Sets}_*, -1}$ is a componentwise inverse to λ^{Sets_*} , it follows from **Item 2 of Proposition 8.8.6.2** that $\lambda^{\text{Sets}_*, -1}$ is also natural. Thus λ^{Sets_*} is a natural isomorphism. ▢

00GP 4.5.6 The Right Unitor

00GQ DEFINITION 4.5.6.1 ► THE RIGHT UNITOR OF \wedge

The **right unitor of the smash product of pointed sets** is the natural isomorphism

$$\rho^{\text{Sets}_*} : \wedge \circ (\text{id} \times \mathbb{1}^{\text{Sets}_*}) \xrightarrow{\sim} \rho_{\text{Sets}_*}^{\text{Cats}_2}$$

whose component

$$\rho_X^{\text{Sets}_*} : X \wedge S^0 \xrightarrow{\cong} X$$

at $X \in \text{Obj}(\text{Sets}_*)$ is given by

$$\begin{aligned} x \wedge 0 &\mapsto x_0, \\ x \wedge 1 &\mapsto x. \end{aligned}$$

PROOF 4.5.6.2 ► PROOF OF DEFINITION 4.5.6.1

Well-Definedness

Let $[(x, y)] = [(x', y')]$ be an element in $X \wedge S^0$. Then either:

1. We have $x = x'$ and $y = y'$.
2. Both of the following conditions are satisfied:
 - (a) We have $x = x_0$ or $y = 0$.
 - (b) We have $x' = x_0$ or $y' = 0$.

In the first case, $\rho_X^{\text{Sets}_*}$ clearly sends both elements to the same element in X . Meanwhile, in the latter case both elements are equal to the basepoint $x_0 \wedge 0$ of $X \wedge S^0$, which gets sent to the basepoint x_0 of X .

Being a Morphism of Pointed Sets

As just mentioned, we have

$$\rho_X^{\text{Sets}_*}(x_0 \wedge 0) \stackrel{\text{def}}{=} x_0,$$

and thus $\rho_X^{\text{Sets}_*}$ is a morphism of pointed sets.

Invertibility

The inverse of $\rho_X^{\text{Sets}_*}$ is the morphism

$$\rho_X^{\text{Sets}_*, -1} : X \xrightarrow{\cong} X \wedge S^0$$

defined by

$$\rho_X^{\text{Sets}_*, -1}(x) \stackrel{\text{def}}{=} x \wedge 1$$

for each $x \in X$. Indeed:

· *Invertibility I.* We have

$$\begin{aligned} \left[\rho_X^{\text{Sets}_*, -1} \circ \rho_X^{\text{Sets}_*} \right] (x \wedge 0) &= \rho_X^{\text{Sets}_*, -1} \left(\rho_X^{\text{Sets}_*} (x \wedge 0) \right) \\ &= \rho_X^{\text{Sets}_*, -1} (x_0) \\ &= x_0 \wedge 1 \\ &= x \wedge 0, \end{aligned}$$

and

$$\begin{aligned} \left[\rho_X^{\text{Sets}_*, -1} \circ \rho_X^{\text{Sets}_*} \right] (x \wedge 1) &= \rho_X^{\text{Sets}_*, -1} \left(\rho_X^{\text{Sets}_*} (x \wedge 1) \right) \\ &= \rho_X^{\text{Sets}_*, -1} (x) \\ &= x \wedge 1 \end{aligned}$$

for each $x \in X$, and thus we have

$$\rho_X^{\text{Sets}_*, -1} \circ \rho_X^{\text{Sets}_*} = \text{id}_{X \wedge S^0}.$$

· *Invertibility II.* We have

$$\begin{aligned} \left[\rho_X^{\text{Sets}_*} \circ \rho_X^{\text{Sets}_*, -1} \right] (x) &= \rho_X^{\text{Sets}_*} \left(\rho_X^{\text{Sets}_*, -1} (x) \right) \\ &= \rho_X^{\text{Sets}_*} (x \wedge 1) \\ &= x \end{aligned}$$

for each $x \in X$, and thus we have

$$\rho_X^{\text{Sets}_*} \circ \rho_X^{\text{Sets}_*, -1} = \text{id}_X.$$

This shows $\rho_X^{\text{Sets}_*}$ to be invertible.

The **symmetry of the smash product of pointed sets** is the natural isomorphism

$$\sigma^{\mathbf{Sets}_*} : \wedge \xrightarrow{\sim} \wedge \circ \sigma_{\mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}_2},$$

whose component

$$\sigma_{X,Y}^{\mathbf{Sets}_*} : X \wedge Y \xrightarrow{\cong} Y \wedge X$$

at $X, Y \in \text{Obj}(\mathbf{Sets}_*)$ is defined by

$$\sigma_{X,Y}^{\mathbf{Sets}_*}(x \wedge y) \stackrel{\text{def}}{=} y \wedge x$$

for each $x \wedge y \in X \wedge Y$.

PROOF 4.5.7.2 ► PROOF OF DEFINITION 4.5.6.1

Well-Definedness

Let $[(x, y)] = [(x', y')]$ be an element in $X \wedge Y$. Then either:

1. We have $x = x'$ and $y = y'$.
2. Both of the following conditions are satisfied:
 - (a) We have $x = x_0$ or $y = y_0$.
 - (b) We have $x' = x_0$ or $y' = y_0$.

In the first case, $\sigma_X^{\mathbf{Sets}_*}$ clearly sends both elements to the same element in X . Meanwhile, in the latter case both elements are equal to the basepoint $x_0 \wedge y_0$ of $X \wedge Y$, which gets sent to the basepoint $y_0 \wedge x_0$ of $Y \wedge X$.

Being a Morphism of Pointed Sets

As just mentioned, we have

$$\sigma_X^{\mathbf{Sets}_*}(x_0 \wedge y_0) \stackrel{\text{def}}{=} y_0 \wedge x_0,$$

and thus $\sigma_X^{\mathbf{Sets}_*}$ is a morphism of pointed sets.

Invertibility

Clearly, the inverse of $\sigma_{X,Y}^{\text{Sets}_*}$ is given by the morphism

$$\sigma_{X,Y}^{\text{Sets}_*, -1}: Y \wedge X \xrightarrow{\cong} X \wedge Y$$

defined by

$$\sigma_{X,Y}^{\text{Sets}_*, -1}(y \wedge x) \stackrel{\text{def}}{=} x \wedge y$$

for each $y \wedge x \in Y \wedge X$.

Naturality

We need to show that, given morphisms of pointed sets

$$f: (X, x_0) \rightarrow (A, a_0),$$

$$g: (Y, y_0) \rightarrow (B, b_0)$$

the diagram


$$\begin{array}{ccc} X \wedge Y & \xrightarrow{f \wedge g} & A \wedge B \\ \sigma_{X,Y}^{\text{Sets}_*} \downarrow & & \downarrow \sigma_{A,B}^{\text{Sets}_*} \\ Y \wedge X & \xrightarrow{g \wedge f} & B \wedge A \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \wedge y & \longmapsto & f(x) \wedge g(y) \\ \downarrow & & \downarrow \\ y \wedge x & \longmapsto & g(y) \wedge f(x) \end{array}$$

and hence indeed commutes, showing σ^{Sets_*} to be a natural transformation.

Being a Natural Isomorphism

Since σ^{Sets_*} is natural and $\sigma^{\text{Sets}_*, -1}$ is a componentwise inverse to σ^{Sets_*} , it follows from [Item 2 of Proposition 8.8.6.2](#) that $\sigma^{\text{Sets}_*, -1}$ is also natural. Thus σ^{Sets_*} is a natural isomorphism. 

00GT 4.5.8 The Diagonal



DEFINITION 4.5.8.1 ► THE DIAGONAL OF \wedge

The **diagonal of the smash product of pointed sets** is the natural transformation

$$\Delta^\wedge : \text{id}_{\text{Sets}_*} \implies \wedge \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2}$$

whose component

$$\Delta_X^\wedge : (X, x_0) \rightarrow (X \wedge X, x_0 \wedge x_0)$$

at $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ is given by the composition

$$\begin{aligned} (X, x_0) &\xrightarrow{\Delta_X^\wedge} (X \times X, (x_0, x_0)) \\ &\longrightarrow ((X \times X)/\sim, [(x_0, x_0)]) \\ &\stackrel{\text{def}}{=} (X \wedge X, x_0 \wedge x_0) \end{aligned}$$

in Sets_* , and thus by

$$\Delta_X^\wedge(x) \stackrel{\text{def}}{=} x \wedge x$$

for each $x \in X$.

PROOF 4.5.8.2 ► PROOF OF DEFINITION 4.5.8.1

Being a Morphism of Pointed Sets

We have

$$\Delta_X^\wedge(x_0) \stackrel{\text{def}}{=} x_0 \wedge x_0,$$

and thus Δ_X^\wedge is a morphism of pointed sets.

Naturality

We need to show that, given a morphism of pointed sets

$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X^\wedge \downarrow & & \downarrow \Delta_Y^\wedge \\ X \wedge X & \xrightarrow{f \wedge f} & Y \wedge Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \longmapsto & f(x) \\ \downarrow & & \downarrow \\ x \wedge x & \longmapsto & f(x) \wedge f(x) \end{array}$$

and hence indeed commutes, showing Δ^\wedge to be natural. ▢

00GV PROPOSITION 4.5.8.3 ► PROPERTIES OF THE DIAGONAL OF \wedge

Let $(X, x_0) \in \text{Obj}(\text{Sets}_*)$.

00GW 1. *Monoidality.* The diagonal

$$\Delta^\wedge : \text{id}_{\text{Sets}_*} \implies \wedge \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2},$$

of the smash product of pointed sets is a monoidal natural transformation:

00GX (a) *Compatibility With Strong Monoidality Constraints.* For each $(X, x_0), (Y, y_0) \in \text{Obj}(\text{Sets}_*)$, the diagram

$$\begin{array}{ccc} X \wedge Y & \xrightarrow{\Delta_X^\wedge \wedge \Delta_Y^\wedge} & (X \wedge X) \wedge (Y \wedge Y) \\ & \searrow \Delta_{X \wedge Y}^\wedge & \downarrow \wr \\ & & (X \wedge Y) \wedge (X \wedge Y) \end{array}$$

commutes.

00GY

(b) *Compatibility With Strong Unitality Constraints.* The diagrams

$$\begin{array}{ccc}
 S^0 & \xrightarrow{\Delta_{S^0}^\wedge} & S^0 \wedge S^0 \\
 \parallel & & \downarrow \lambda_{S^0}^{\text{Sets}_*} \\
 & & S^0
 \end{array}
 \qquad
 \begin{array}{ccc}
 S^0 & \xrightarrow{\Delta_{S^0}^\wedge} & S^0 \wedge S^0 \\
 \parallel & & \downarrow \rho_{S^0}^{\text{Sets}_*} \\
 & & S^0
 \end{array}$$

commute, i.e. we have

$$\begin{aligned}
 \Delta_{S^0}^\wedge &= \lambda_{S^0}^{\text{Sets}_*, -1} \\
 &= \rho_{S^0}^{\text{Sets}_*, -1},
 \end{aligned}$$

where we recall that the equalities

$$\begin{aligned}
 \lambda_{S^0}^{\text{Sets}_*} &= \rho_{S^0}^{\text{Sets}_*}, \\
 \lambda_{S^0}^{\text{Sets}_*, -1} &= \rho_{S^0}^{\text{Sets}_*, -1}
 \end{aligned}$$

are always true in any monoidal category by ?? of ??.

00GZ

2. *The Diagonal of the Unit.* The component

$$\Delta_{S^0}^\wedge : S^0 \xrightarrow{\cong} S^0 \wedge S^0$$

of Δ^\wedge at S^0 is an isomorphism.

PROOF 4.5.8.4 ► PROOF OF PROPOSITION 4.5.8.3

Item 1: Monoidality

We claim that Δ^\wedge is indeed monoidal:

1. *Item 1a: Compatibility With Strong Monoidality Constraints:* We need to show that the diagram

$$\begin{array}{ccc}
 X \wedge Y & \xrightarrow{\Delta_X^\wedge \wedge \Delta_Y^\wedge} & (X \wedge X) \wedge (Y \wedge Y) \\
 \searrow \Delta_{X \wedge Y}^\wedge & & \downarrow \lambda \\
 & & (X \wedge Y) \wedge (X \wedge Y)
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \wedge y & \xrightarrow{\quad} & (x \wedge x) \wedge (y \wedge y) \\ & \searrow & \downarrow \\ & & (x \wedge y) \wedge (x \wedge y) \end{array}$$

and hence indeed commutes.

2. *Item 1b: Compatibility With Strong Unitality Constraints:* As shown in the proof of [Definition 4.5.5.1](#), the inverse of the left unitor of \mathbf{Sets}_* with respect to the smash product of pointed sets at $(X, x_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$ is given by

$$\lambda_X^{\mathbf{Sets}_*, -1}(x) \stackrel{\text{def}}{=} 1 \wedge x$$

for each $x \in X$, so when $X = S^0$, we have

$$\begin{aligned} \lambda_{S^0}^{\mathbf{Sets}_*, -1}(0) &\stackrel{\text{def}}{=} 1 \wedge 0, \\ \lambda_{S^0}^{\mathbf{Sets}_*, -1}(1) &\stackrel{\text{def}}{=} 1 \wedge 1. \end{aligned}$$


But since $1 \wedge 0 = 0 \wedge 0$ and

$$\begin{aligned} \Delta_{S^0}^\wedge(0) &\stackrel{\text{def}}{=} 0 \wedge 0, \\ \Delta_{S^0}^\wedge(1) &\stackrel{\text{def}}{=} 1 \wedge 1, \end{aligned}$$

it follows that we indeed have $\Delta_{S^0}^\wedge = \lambda_{S^0}^{\mathbf{Sets}_*, -1}$.

This finishes the proof.

Item 2: The Diagonal of the Unit

This follows from [Item 1](#) and the invertibility of the left/right unitor of \mathbf{Sets}_* with respect to \wedge , proved in the proof of [Definition 4.5.5.1](#) for the left unitor or the proof of [Definition 4.5.6.1](#) for the right unitor. 

00H0 4.5.9 The Monoidal Structure on Pointed Sets Associated to \wedge

00H1

PROPOSITION 4.5.9.1 ► THE MONOIDAL STRUCTURE ON POINTED SETS ASSOCIATED TO \wedge

The category \mathbf{Sets}_* admits a closed monoidal category with diagonals structure consisting of

- *The Underlying Category.* The category \mathbf{Sets}_* of pointed sets;

- *The Monoidal Product.* The smash product functor

$$\wedge : \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

of [Item 1](#) of [Proposition 4.5.1.10](#);

- *The Internal Hom.* The internal Hom functor

$$\mathbf{Sets}_* : \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

of [Item 1](#) of [Proposition 4.5.2.3](#);

- *The Monoidal Unit.* The functor

$$\mathbb{1}^{\mathbf{Sets}_*} : \text{pt} \rightarrow \mathbf{Sets}_*$$

of [Definition 4.5.3.1](#);

- *The Associators.* The natural isomorphism

$$\alpha^{\mathbf{Sets}_*} : \wedge \circ (\wedge \times \text{id}_{\mathbf{Sets}_*}) \xrightarrow{\sim} \wedge \circ (\text{id}_{\mathbf{Sets}_*} \times \wedge) \circ \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}}$$

of [Definition 4.5.4.1](#);

- *The Left Unitors.* The natural isomorphism

$$\lambda^{\mathbf{Sets}_*} : \wedge \circ (\mathbb{1}^{\mathbf{Sets}_*} \times \text{id}_{\mathbf{Sets}_*}) \xrightarrow{\sim} \lambda_{\mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

of [Definition 4.5.5.1](#);

- *The Right Unitors.* The natural isomorphism

$$\rho^{\mathbf{Sets}_*} : \wedge \circ (\text{id} \times \mathbb{1}^{\mathbf{Sets}_*}) \xrightarrow{\sim} \rho_{\mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

of [Definition 4.5.6.1](#);

- *The Symmetry.* The natural isomorphism

$$\sigma^{\mathbf{Sets}_*} : \wedge \xrightarrow{\sim} \wedge \circ \sigma_{\mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

of [Definition 4.5.7.1](#);

- *The Diagonals.* The monoidal natural transformation


$$\Delta^\wedge : \text{id}_{\mathbf{Sets}_*} \xrightarrow{\sim} \wedge \circ \Delta_{\mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

of [Definition 4.5.8.1](#).


PROOF 4.5.9.2 ► PROOF OF PROPOSITION 4.5.9.1

The Pentagon Identity

Let (W, w_0) , (X, x_0) , (Y, y_0) and (Z, z_0) be pointed sets. We have to show that the diagram



commutes. Indeed, this diagram acts on elements as



and thus we see that the pentagon identity is satisfied.

The Triangle Identity

Let (X, x_0) and (Y, y_0) be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 (X \wedge S^0) \wedge Y & \xrightarrow{\alpha_{X,S^0,Y}^{\text{Sets}^*}} & X \wedge (S^0 \wedge Y) \\
 \searrow \rho_X^{\text{Sets}^*} \wedge \text{id}_Y & & \swarrow \text{id}_X \wedge \lambda_Y^{\text{Sets}^*} \\
 & X \wedge Y &
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 (x \wedge 0) \wedge y & \xrightarrow{\quad} & x \wedge (0 \wedge y) \\
 \searrow & & \swarrow \\
 x_0 \wedge y & & x \wedge y_0
 \end{array}$$

and

$$\begin{array}{ccc}
 (x \wedge 1) \wedge y & \xrightarrow{\quad} & x \wedge (1 \wedge y) \\
 \searrow & & \swarrow \\
 & x \wedge y &
 \end{array}$$


and thus we see that the triangle identity is satisfied.

The Left Hexagon Identity

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 & (X \wedge Y) \wedge Z & \\
 \alpha_{X,Y,Z}^{\text{Sets}^*} \swarrow & & \searrow \beta_{X,Y}^{\text{Sets}^*} \wedge \text{id}_Z \\
 X \wedge (Y \wedge Z) & & (Y \wedge X) \wedge Z \\
 \beta_{X,Y \wedge Z}^{\text{Sets}^*} \downarrow & & \downarrow \alpha_{Y,X,Z}^{\text{Sets}^*} \\
 (Y \wedge Z) \wedge X & & Y \wedge (X \wedge Z) \\
 \alpha_{Y,Z,X}^{\text{Sets}^*} \swarrow & & \swarrow \text{id}_Y \wedge \beta_{X,Z}^{\text{Sets}^*} \\
 & Y \wedge (Z \wedge X) &
 \end{array}$$


commutes. Indeed, this diagram acts on elements as




and thus we see that the left hexagon identity is satisfied.

The Right Hexagon Identity

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets. We have to show that the diagram



commutes. Indeed, this diagram acts on elements as




and thus we see that the right hexagon identity is satisfied.

Monoidal Closedness

This follows from **Item 2** of **Proposition 4.5.1.10**.

Existence of Monoidal Diagonals

This follows from **Items 1** and **2** of **Proposition 4.5.8.3**. 

00H2 4.5.10 Universal Properties of the Smash Product of Pointed Sets I

00H3

THEOREM 4.5.10.1 ► UNIVERSAL PROPERTIES OF THE SMASH PRODUCT OF POINTED SETS I

The symmetric monoidal structure on the category \mathbf{Sets}_* is uniquely determined by the following requirements:

1. *Two-Sided Preservation of Colimits*. The smash product

$$\wedge : \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

of \mathbf{Sets}_* preserves colimits separately in each variable.

2. *The Unit Object Is S^0* . We have $\mathbb{1}_{\mathbf{Sets}_*} = S^0$.

PROOF 4.5.10.2 ► PROOF OF THEOREM 4.5.10.1

Omitted. 

00H4 4.5.11 Universal Properties of the Smash Product of Pointed Sets II

00H5

THEOREM 4.5.11.1 ► UNIVERSAL PROPERTIES OF THE SMASH PRODUCT OF POINTED SETS II

The symmetric monoidal structure on the category \mathbf{Sets}_* is the unique symmetric monoidal structure on \mathbf{Sets}_* such that the free pointed set functor

$$(-)^+ : \mathbf{Sets} \rightarrow \mathbf{Sets}_*$$

admits a symmetric monoidal structure.

PROOF 4.5.11.2 ► PROOF OF THEOREM 4.5.11.1

See [CGN15, Theorem 5.1].

00H6 4.5.12 Monoids With Respect to the Smash Product of Pointed Sets

00H7

PROPOSITION 4.5.12.1 ► MONOIDS WITH RESPECT TO \wedge

The category of monoids on $(\mathbf{Sets}_*, \wedge, S^0)$ is isomorphic to the category of monoids with zero and morphisms between them.

PROOF 4.5.12.2 ► PROOF OF PROPOSITION 4.5.12.1

See ??, in particular ??, ??, and ??.

00H8 4.5.13 Comonoids With Respect to the Smash Product of Pointed Sets

00H9

PROPOSITION 4.5.13.1 ► COMONOIDS WITH RESPECT TO \wedge

The symmetric monoidal functor

$$((-)^+, (-)^{+, \times}, (-)_{\perp}^{+, \times}) : (\mathbf{Sets}, \times, \text{pt}) \rightarrow (\mathbf{Sets}_*, \wedge, S^0),$$

of [Item 4](#) of [Proposition 3.4.1.2](#) lifts to an equivalence of categories

$$\begin{aligned} \text{CoMon}(\mathbf{Sets}_*, \wedge, S^0) &\stackrel{\text{eq.}}{\cong} \text{CoMon}(\mathbf{Sets}, \times, \text{pt}) \\ &\cong \mathbf{Sets}. \end{aligned}$$

PROOF 4.5.13.2 ► PROOF OF PROPOSITION 4.5.13.1

See [PS19, Lemma 2.4].

00HA 4.6 Miscellany

00HB 4.6.1 The Smash Product of a Family of Pointed Sets

Let $\{(X_i, x_0^i)\}_{i \in I}$ be a family of pointed sets.

00HC

DEFINITION 4.6.1.1 ► THE SMASH PRODUCT OF A FAMILY OF POINTED SETS

The **smash product of the family** $\{(X_i, x_0^i)\}_{i \in I}$ is the pointed set $\bigwedge_{i \in I} X_i$ consisting of:

- *The Underlying Set.* The set $\bigwedge_{i \in I} X_i$ defined by

$$\bigwedge_{i \in I} X_i \stackrel{\text{def}}{=} \left(\prod_{i \in I} X_i \right) / \sim,$$

where \sim is the equivalence relation on $\prod_{i \in I} X_i$ obtained by declaring

$$(x_i)_{i \in I} \sim (y_i)_{i \in I}$$

if there exist $i_0 \in I$ such that $x_{i_0} = x_0$ and $y_{i_0} = y_0$, for each $(x_i)_{i \in I}, (y_i)_{i \in I} \in \prod_{i \in I} X_i$.

- *The Basepoint.* The element $[(x_0)_{i \in I}]$ of $\bigwedge_{i \in I} X_i$.

Appendices

4.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets

Relations

5. Relations

6. Constructions With Relations

7. Equivalence Relations and Apartness Relations

Category Theory

8. Categories

Bicategories

9. Types of Morphisms in Bicategories

Part II

Relations

Chapter 5

Relations

00HD This chapter contains some material about relations. Notably, we discuss and explore:

1. The definition of relations ([Section 5.1.1](#)).
2. How relations may be viewed as decategorification of profunctors ([Section 5.1.2](#)).
3. The various kind of categories that relations form, namely:
 - (a) A category ([Section 5.2.1](#)).
 - (b) A monoidal category ([Section 5.2.2](#)).
 - (c) A 2-category ([Section 5.2.3](#)).
 - (d) A double category ([Section 5.2.4](#)).
4. The various categorical properties of the 2-category of relations, including:
 - (a) The self-duality of \mathbf{Rel} and \mathbf{Rel} ([Proposition 5.3.1.1](#)).
 - (b) Identifications of equivalences and isomorphisms in \mathbf{Rel} with bijections ([Proposition 5.3.2.1](#)).
 - (c) Identifications of adjunctions in \mathbf{Rel} with functions ([Proposition 5.3.3.1](#)).
 - (d) Identifications of monads in \mathbf{Rel} with preorders ([Proposition 5.3.4.1](#)).
 - (e) Identifications of comonads in \mathbf{Rel} with subsets ([Proposition 5.3.5.1](#)).
 - (f) A description of the monoids and comonoids in \mathbf{Rel} with respect to the Cartesian product ([Remark 5.3.6.1](#)).
 - (g) Characterisations of monomorphisms in \mathbf{Rel} ([Proposition 5.3.7.1](#)).
 - (h) Characterisations of 2-categorical notions of monomorphisms in \mathbf{Rel} ([Proposition 5.3.8.1](#)).
 - (i) Characterisations of epimorphisms in \mathbf{Rel} ([Proposition 5.3.9.1](#)).

- (j) Characterisations of 2-categorical notions of epimorphisms in **Rel** (Proposition 5.3.10.1).
 - (k) The partial co/completeness of **Rel** (Proposition 5.3.11.1).
 - (l) The existence or non-existence of Kan extensions and Kan lifts in **Rel** (Remark 5.3.12.1).
 - (m) The closedness of **Rel** (Proposition 5.3.13.1).
 - (n) The identification of **Rel** with the category of free algebras of the powerset monad on **Sets** (Proposition 5.3.14.1).
5. A description of two notions of “skew composition” on **Rel**(A, B), giving rise to left and right skew monoidal structures analogous to the left skew monoidal structure on $\text{Fun}(C, \mathcal{D})$ appearing in the definition of a relative monad (Sections 5.4 and 5.5).

Contents

5.1	Relations	274
5.1.1	Foundations	274
5.1.2	Relations as Decategorifications of Profunctors	278
5.1.3	Examples of Relations	279
5.1.4	Functional Relations	282
5.1.5	Total Relations	283
5.2	Categories of Relations	284
5.2.1	The Category of Relations	284
5.2.2	The Closed Symmetric Monoidal Category of Relations	284
5.2.3	The 2-Category of Relations	290
5.2.4	The Double Category of Relations	291
5.3	Properties of the 2-Category of Relations	299
5.3.1	Self-Duality	299
5.3.2	Isomorphisms and Equivalences in Rel	300
5.3.3	Adjunctions in Rel	301
5.3.4	Monads in Rel	304
5.3.5	Comonads in Rel	305
5.3.6	Co/Monoids in Rel	306
5.3.7	Monomorphisms in Rel	306
5.3.8	2-Categorical Monomorphisms in Rel	309
5.3.9	Epimorphisms in Rel	313
5.3.10	2-Categorical Epimorphisms in Rel	317
5.3.11	Co/Limits in Rel	321
5.3.12	Kan Extensions and Kan Lifts in Rel	321

5.3.13	Closedness of Rel	321
5.3.14	Rel as a Category of Free Algebras.....	322
5.4	The Left Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	323
5.4.1	The Left Skew Monoidal Product	323
5.4.2	The Left Skew Monoidal Unit.....	324
5.4.3	The Left Skew Associators	324
5.4.4	The Left Skew Left Unitors	325
5.4.5	The Left Skew Right Unitors	326
5.4.6	The Left Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	327
5.5	The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	328
5.5.1	The Right Skew Monoidal Product.....	328
5.5.2	The Right Skew Monoidal Unit.....	329
5.5.3	The Right Skew Associators	330
5.5.4	The Right Skew Left Unitors	331
5.5.5	The Right Skew Right Unitors	331
5.5.6	The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	332
5.A	Other Chapters	333

00HE 5.1 Relations

00HF 5.1.1 Foundations

Let A and B be sets.

00HG DEFINITION 5.1.1.1 ► RELATIONS

A **relation** $R: A \rightarrow B$ **from A to B** ^{1,2} is a subset R of $A \times B$.

¹*Further Terminology:* Also called a **multivalued function from A to B** , a **relation over A and B** , **relation on A and B** , a **binary relation over A and B** , or a **binary relation on A and B** .

²*Further Terminology:* When $A = B$, we also call $R \subset A \times A$ a **relation on A** .

00HH NOTATION 5.1.1.2 ► FURTHER NOTATION FOR RELATIONS

Let $R: A \rightarrow B$ be a relation.

1. Given elements $a \in A$ and $b \in B$ and a relation $R: A \rightarrow B$, we write $a \sim_R b$ to mean $(a, b) \in R$.

00HJ

2. Viewing R as a function

00HK

$$R: A \times B \rightarrow \{t, f\}$$

via [Remark 5.1.1.4](#), we write R_a^b for the value of R at (a, b) .¹

¹The choice R_a^b in place of R_b^a is to keep the notation consistent with the notation we will later employ for profunctors.

00HL DEFINITION 5.1.1.3 ► THE PO/SET OF RELATIONS OVER TWO SETS

Let A and B be sets.

- 00HM 1. The **set of relations from A to B** is the set $\text{Rel}(A, B)$ defined by

$$\text{Rel}(A, B) \stackrel{\text{def}}{=} \{\text{Relations from } A \text{ to } B\}.$$

- 00HN 2. The **poset of relations from A to B** is the poset

$$\mathbf{Rel}(A, B) \stackrel{\text{def}}{=} (\text{Rel}(A, B), \subset)$$

consisting of:

- *The Underlying Set.* The set $\text{Rel}(A, B)$ of [Item 1](#).
- *The Partial Order.* The partial order

$$\subset : \text{Rel}(A, B) \times \text{Rel}(A, B) \rightarrow \{\text{true}, \text{false}\}$$

on $\text{Rel}(A, B)$ given by inclusion of relations.

- 00HP 3. The **category of relations from A to B** is the posetal category $\mathbf{Rel}(A, B)$ ¹ associated to the poset $\mathbf{Rel}(A, B)$ of [Item 2](#) via [Definition 8.1.3.1](#).

¹Here we choose to slightly abuse notation by writing $\mathbf{Rel}(A, B)$ (instead of e.g. $\mathbf{Rel}(A, B)_{\text{pos}}$) for the posetal category of relations from A to B , even though the same notation is used for the poset of relations from A to B .

00HQ REMARK 5.1.1.4 ► EQUIVALENT DEFINITIONS OF RELATIONS

A relation from A to B is equivalently:¹

- 00HR 1. A subset of $A \times B$.
- 00HS 2. A function from $A \times B$ to $\{\text{true}, \text{false}\}$.
- 00HT 3. A function from A to $\mathcal{P}(B)$.
- 00HU 4. A function from B to $\mathcal{P}(A)$.
- 00HV 5. A cocontinuous morphism of posets from $(\mathcal{P}(A), \subset)$ to $(\mathcal{P}(B), \subset)$.

That is: we have bijections of sets

$$\begin{aligned} \text{Rel}(A, B) &\stackrel{\text{def}}{=} \mathcal{P}(A \times B), \\ &\cong \text{Hom}_{\text{Sets}}(A \times B, \{\text{true}, \text{false}\}), \\ &\cong \text{Hom}_{\text{Sets}}(A, \mathcal{P}(B)), \\ &\cong \text{Hom}_{\text{Sets}}(B, \mathcal{P}(A)), \\ &\cong \text{Hom}_{\text{Pos}}^{\text{cocont}}(\mathcal{P}(A), \mathcal{P}(B)), \end{aligned}$$

natural in $A, B \in \text{Obj}(\text{Sets})$.

¹*Intuition:* In particular, we may think of a relation $R: A \rightarrow \mathcal{P}(B)$ from A to B as a multivalued function from A to B (including the possibility of a given $a \in A$ having no value at all).

PROOF 5.1.1.5 ► PROOF OF REMARK 5.1.1.4

We claim that **Items 1 to 5** are indeed equivalent:

- **Item 1** \iff **Item 2**: This is a special case of **Items 1 and 2** of **Proposition 2.4.3.9**.
- **Item 2** \iff **Item 3**: This follows from the bijections

$$\begin{aligned} \text{Hom}_{\text{Sets}}(A \times B, \{\text{true}, \text{false}\}) &\cong \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(B, \{\text{true}, \text{false}\})) \\ &\cong \text{Hom}_{\text{Sets}}(A, \mathcal{P}(B)), \end{aligned}$$

where the last bijection is from **Items 1 and 2** of **Proposition 2.4.3.9**.

- **Item 2** \iff **Item 4**: This follows from the bijections

$$\begin{aligned} \text{Hom}_{\text{Sets}}(A \times B, \{\text{true}, \text{false}\}) &\cong \text{Hom}_{\text{Sets}}(B, \text{Hom}_{\text{Sets}}(A, \{\text{true}, \text{false}\})) \\ &\cong \text{Hom}_{\text{Sets}}(B, \mathcal{P}(A)), \end{aligned}$$

where again the last bijection is from **Items 1 and 2** of **Proposition 2.4.3.9**.

- **Item 2** \iff **Item 5**: This follows from the universal property of the powerset $\mathcal{P}(X)$ of a set X as the free cocompletion of X via the characteristic embedding

$$\chi_X: X \hookrightarrow \mathcal{P}(X)$$


of X into $\mathcal{P}(X)$, **Item 2** of **Proposition 2.4.3.12**.

In particular, the bijection

$$\text{Rel}(A, B) \cong \text{Hom}_{\text{Pos}}^{\text{cocont}}(\mathcal{P}(A), \mathcal{P}(B))$$

is given by taking a relation $R: A \rightarrow B$, passing to its associated function $f: A \rightarrow \mathcal{P}(B)$ from A to B and then extending f from A to all of $\mathcal{P}(A)$ by taking its left Kan extension along χ_X .

This coincides with the direct image function $f_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$ of [Definition 2.4.4.1](#).

This finishes the proof. 

00HW

PROPOSITION 5.1.1.6 ► PROPERTIES OF RELATIONS

Let A and B be sets and let $R, S: A \rightarrow B$ be relations.

00HX

1. *End Formula for the Set of Inclusions of Relations.* We have

$$\mathrm{Hom}_{\mathbf{Rel}(A,B)}(R, S) \cong \int_{a \in A} \int_{b \in B} \mathrm{Hom}_{\{t,f\}}(R_a^b, S_a^b).$$

PROOF 5.1.1.7 ► PROOF OF PROPOSITION 5.1.1.6

Item 1: End Formula for the Set of Inclusions of Relations

Unwinding the expression inside the end on the right hand side, we have


$$\int_{a \in A} \int_{b \in B} \mathrm{Hom}_{\{t,f\}}(R_a^b, S_a^b) \cong \begin{cases} \mathrm{pt} & \text{if, for each } a \in A \text{ and each } b \in B, \\ & \text{we have } \mathrm{Hom}_{\{t,f\}}(R_a^b, S_a^b) \cong \mathrm{pt} \\ \emptyset & \text{otherwise.} \end{cases}$$

Since we have $\mathrm{Hom}_{\{t,f\}}(R_a^b, S_a^b) = \{\mathrm{true}\} \cong \mathrm{pt}$ exactly when $R_a^b = \mathrm{false}$ or $R_a^b = S_a^b = \mathrm{true}$, we get

$$\int_{a \in A} \int_{b \in B} \mathrm{Hom}_{\{t,f\}}(R_a^b, S_a^b) \cong \begin{cases} \mathrm{pt} & \text{if, for each } a \in A \text{ and each } b \in B, \\ & \text{if } a \sim_R b, \text{ then } a \sim_S b, \\ \emptyset & \text{otherwise.} \end{cases}$$

On the left hand-side, we have

$$\mathrm{Hom}_{\mathbf{Rel}(A,B)}(R, S) \cong \begin{cases} \mathrm{pt} & \text{if } R \subset S, \\ \emptyset & \text{otherwise.} \end{cases}$$

It is then clear that the conditions for each set to evaluate to pt (up to isomorphism) are equivalent, implying that those two sets are isomorphic. 

00HY **5.1.2 Relations as Decategorifications of Profunctors**

00HZ **REMARK 5.1.2.1 ► RELATIONS AS DECATEGORIFICATIONS OF PROFUNCTORS I**

The notion of a relation is a decategorification of that of a profunctor:

1. A profunctor from a category C to a category \mathcal{D} is a functor

$$p: \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}.$$

2. A relation on sets A and B is a function

$$R: A \times B \rightarrow \{\text{true}, \text{false}\}.$$

Here we notice that:

- The opposite X^{op} of a set X is itself, as $(-)^{\text{op}}: \text{Cats} \rightarrow \text{Cats}$ restricts to the identity endofunctor on Sets.
- The values that profunctors and relations take are analogous:
 - A category is enriched over the category

$$\text{Sets} \stackrel{\text{def}}{=} \text{Cats}_0$$

of sets, with profunctors taking values on it.

- A set is enriched over the set

$$\{\text{true}, \text{false}\} \stackrel{\text{def}}{=} \text{Cats}_{-1}$$

of classical truth values, with relations taking values on it.

00J0 **REMARK 5.1.2.2 ► RELATIONS AS DECATEGORIFICATIONS OF PROFUNCTORS II**

Extending Remark 5.1.2.1, the equivalent definitions of relations in Remark 5.1.1.4 are also related to the corresponding ones for profunctors (??), which state that a profunctor $p: C \rightarrow \mathcal{D}$ is equivalently:

- 00J1 1. A functor $p: \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}$.
- 00J2 2. A functor $p: C \rightarrow \text{PSh}(\mathcal{D})$.
- 00J3 3. A functor $p: \mathcal{D}^{\text{op}} \rightarrow \text{Fun}(C, \text{Sets})$.
- 00J4 4. A colimit-preserving functor $p: \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$.

Indeed:

- The equivalence between **Items 1** and **2** (and also that between **Items 1** and **3**, which is proved analogously) is an instance of currying, both for profunctors as well as for relations, using the isomorphisms

$$\begin{aligned} \text{Sets}(A \times B, \{\text{true}, \text{false}\}) &\cong \text{Sets}(A, \text{Sets}(B, \{\text{true}, \text{false}\})) \\ &\cong \text{Sets}(A, \mathcal{P}(B)), \\ \text{Fun}(\mathcal{D}^{\text{op}} \times \mathcal{D}, \text{Sets}) &\cong \text{Fun}(C, \text{Fun}(\mathcal{D}^{\text{op}}, \text{Sets})) \\ &\cong \text{Fun}(C, \text{PSh}(\mathcal{D})). \end{aligned}$$

- The equivalence between **Items 1** and **3** follows from the universal properties of:

- The powerset $\mathcal{P}(X)$ of a set X as the free cocompletion of X via the characteristic embedding

$$\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$$

of X into $\mathcal{P}(X)$, as stated and proved in **Item 2** of **Proposition 2.4.3.12**.

- The category $\text{PSh}(C)$ of presheaves on a category C as the free cocompletion of C via the Yoneda embedding

$$\mathcal{Y} : C \hookrightarrow \text{PSh}(C)$$

of C into $\text{PSh}(C)$, as stated and proved in ?? of ??.

00J5 5.1.3 Examples of Relations

00J6 EXAMPLE 5.1.3.1 ► THE TRIVIAL RELATION

The **trivial relation on A and B** is the relation \sim_{triv} defined equivalently as follows:

1. As a subset of $A \times B$, we have

$$\sim_{\text{triv}} \stackrel{\text{def}}{=} A \times B.$$

2. As a function from $A \times B$ to $\{\text{true}, \text{false}\}$, the relation \sim_{triv} is the constant function

$$\Delta_{\text{true}} : A \times B \rightarrow \{\text{true}, \text{false}\}$$

from $A \times B$ to $\{\text{true}, \text{false}\}$ taking the value true.

3. As a function from A to $\mathcal{P}(B)$, the relation \sim_{triv} is the function

$$\Delta_{\text{true}}: A \rightarrow \mathcal{P}(B)$$

defined by

$$\Delta_{\text{true}}(a) \stackrel{\text{def}}{=} B$$

for each $a \in A$.

4. Lastly, it is the unique relation R on A and B such that we have $a \sim_R b$ for each $a \in A$ and each $b \in B$.

00J7

EXAMPLE 5.1.3.2 ► THE COTRIVIAL RELATION

The **cotrivial relation on A and B** is the relation \sim_{cotriv} defined equivalently as follows:

1. As a subset of $A \times B$, we have

$$\sim_{\text{cotriv}} \stackrel{\text{def}}{=} \emptyset.$$

2. As a function from $A \times B$ to $\{\text{true}, \text{false}\}$, the relation \sim_{cotriv} is the constant function

$$\Delta_{\text{false}}: A \times B \rightarrow \{\text{true}, \text{false}\}$$

from $A \times B$ to $\{\text{true}, \text{false}\}$ taking the value false.

3. As a function from A to $\mathcal{P}(B)$, the relation \sim_{cotriv} is the function

$$\Delta_{\text{false}}: A \rightarrow \mathcal{P}(B)$$

defined by

$$\Delta_{\text{false}}(a) \stackrel{\text{def}}{=} \emptyset$$

for each $a \in A$.

4. Lastly, it is the unique relation R on A and B such that we have $a \not\sim_R b$ for each $a \in A$ and each $b \in B$.

00J8 EXAMPLE 5.1.3.3 ► THE CHARACTERISTIC RELATION OF A SET

The characteristic relation

$$\chi_X(-1, -2): X \times X \rightarrow \{t, f\}$$

on X of Item 3 of Definition 2.4.1.1, defined by

$$\chi_X(x, y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each $x, y \in X$, is another example of a relation.

00J9 EXAMPLE 5.1.3.4 ► SQUARE ROOTS

Square roots are examples of relations:

1. *Square Roots in \mathbb{R}* . The assignment $x \mapsto \sqrt{x}$ defines a relation

$$\sqrt{-}: \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$$

from \mathbb{R} to itself, being explicitly given by

$$\sqrt{x} \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } x = 0, \\ \{-\sqrt{|x|}, \sqrt{|x|}\} & \text{if } x \neq 0. \end{cases}$$

2. *Square Roots in \mathbb{Q}* . Square roots in \mathbb{Q} are similar to square roots in \mathbb{R} , though now additionally it may also occur that $\sqrt{-}: \mathbb{Q} \rightarrow \mathcal{P}(\mathbb{Q})$ sends a rational number x (e.g. 2) to the empty set (since $\sqrt{2} \notin \mathbb{Q}$).

00JA EXAMPLE 5.1.3.5 ► COMPLEX LOGARITHMS

The complex logarithm defines a relation

$$\log: \mathbb{C} \rightarrow \mathcal{P}(\mathbb{C})$$

from \mathbb{C} to itself, where we have

$$\log(a + bi) \stackrel{\text{def}}{=} \left\{ \log\left(\sqrt{a^2 + b^2}\right) + i \arg(a + bi) + (2\pi i)k \mid k \in \mathbb{Z} \right\}$$

for each $a + bi \in \mathbb{C}$.

00JB **EXAMPLE 5.1.3.6 ► MORE EXAMPLES OF RELATIONS**

See [Wik24] for more examples of relations, such as antiderivation, inverse trigonometric functions, and inverse hyperbolic functions.

00JC **5.1.4 Functional Relations**

Let A and B be sets.

00JD **DEFINITION 5.1.4.1 ► FUNCTIONAL RELATIONS**

A relation $R: A \rightarrow B$ is **functional** if, for each $a \in A$, the set $R(a)$ is either empty or a singleton.

00JE **PROPOSITION 5.1.4.2 ► PROPERTIES OF FUNCTIONAL RELATIONS**

Let $R: A \rightarrow B$ be a relation.

00JF 1. *Characterisations.* The following conditions are equivalent:

00JG (a) The relation R is functional.

00JH (b) We have $R \diamond R^\dagger \subset \chi_B$.

PROOF 5.1.4.3 ► PROOF OF PROPOSITION 5.1.4.2

Item 1: Characterisations

We claim that **Items 1a** and **1b** are indeed equivalent:

• **Item 1a** \implies **Item 1b**: Let $(b, b') \in B \times B$. We need to show that

$$[R \diamond R^\dagger](b, b') \preceq_{\{t,f\}} \chi_B(b, b'),$$

i.e. that if there exists some $a \in A$ such that $b \sim_{R^\dagger} a$ and $a \sim_R b'$, then $b = b'$. But since $b \sim_{R^\dagger} a$ is the same as $a \sim_R b$, we have both $a \sim_R b$ and $a \sim_R b'$ at the same time, which implies $b = b'$ since R is functional.


• **Item 1b** \implies **Item 1a**: Suppose that we have $a \sim_R b$ and $a \sim_R b'$ for $b, b' \in B$. We claim that $b = b'$:

1. Since $a \sim_R b$, we have $b \sim_{R^\dagger} a$.

2. Since $R \diamond R^\dagger \subset \chi_B$, we have

$$[R \diamond R^\dagger](b, b') \preceq_{\{t,f\}} \chi_B(b, b'),$$

and since $b \sim_{R^\dagger} a$ and $a \sim_R b'$, it follows that $[R \diamond R^\dagger](b, b') = \text{true}$, and thus $\chi_B(b, b') = \text{true}$ as well, i.e. $b = b'$.

This finishes the proof. 

00JJ 5.1.5 Total Relations

Let A and B be sets.

00JK DEFINITION 5.1.5.1 ► TOTAL RELATIONS

A relation $R: A \rightarrow B$ is **total** if, for each $a \in A$, we have $R(a) \neq \emptyset$.

00JL PROPOSITION 5.1.5.2 ► PROPERTIES OF TOTAL RELATIONS

Let $R: A \rightarrow B$ be a relation.

- 00JM 1. *Characterisations.* The following conditions are equivalent:
- 00JN (a) The relation R is total.
- 00JP (b) We have $\chi_A \subset R^\dagger \diamond R$.

PROOF 5.1.5.3 ► PROOF OF PROPOSITION 5.1.5.2

Item 1: Characterisations

We claim that **Item 1a** and **1b** are indeed equivalent:

- **Item 1a** \implies **Item 1b**: We have to show that, for each $(a, a') \in A$, we have


$$\chi_A(a, a') \preceq_{\{t, f\}} [R^\dagger \diamond R](a, a'),$$

i.e. that if $a = a'$, then there exists some $b \in B$ such that $a \sim_R b$ and $b \sim_{R^\dagger} a'$ (i.e. $a \sim_R b$ again), which follows from the totality of R .

- **Item 1b** \implies **Item 1a**: Given $a \in A$, since $\chi_A \subset R^\dagger \diamond R$, we must have

$$\{a\} \subset [R^\dagger \diamond R](a),$$

implying that there must exist some $b \in B$ such that $a \sim_R b$ and $b \sim_{R^\dagger} a$ (i.e. $a \sim_R b$) and thus $R(a) \neq \emptyset$, as $b \in R(a)$.

This finishes the proof. 

00JQ 5.2 Categories of Relations

00JR 5.2.1 The Category of Relations

00JS DEFINITION 5.2.1.1 ► THE CATEGORY OF RELATIONS

The **category of relations** is the category Rel where

- *Objects.* The objects of Rel are sets.
- *Morphisms.* For each $A, B \in \text{Obj}(\text{Sets})$, we have

$$\text{Rel}(A, B) \stackrel{\text{def}}{=} \text{Rel}(A, B).$$

- *Identities.* For each $A \in \text{Obj}(\text{Rel})$, the unit map

$$\mathbb{1}_A^{\text{Rel}} : \text{pt} \rightarrow \text{Rel}(A, A)$$

of Rel at A is defined by

$$\text{id}_A^{\text{Rel}} \stackrel{\text{def}}{=} \chi_A(-1, -2),$$

where $\chi_A(-1, -2)$ is the characteristic relation of A of [Item 3 of Definition 2.4.1.1](#).

- *Composition.* For each $A, B, C \in \text{Obj}(\text{Rel})$, the composition map

$$\circ_{A,B,C}^{\text{Rel}} : \text{Rel}(B, C) \times \text{Rel}(A, B) \rightarrow \text{Rel}(A, C)$$

of Rel at (A, B, C) is defined by

$$S \circ_{A,B,C}^{\text{Rel}} R \stackrel{\text{def}}{=} S \diamond R$$

for each $(S, R) \in \text{Rel}(B, C) \times \text{Rel}(A, B)$, where $S \diamond R$ is the composition of S and R of [Definition 6.3.12.1](#).

00JT 5.2.2 The Closed Symmetric Monoidal Category of Relations

00JU 5.2.2.1 The Monoidal Product

00JV DEFINITION 5.2.2.1 ► THE MONOIDAL PRODUCT OF REL

The **monoidal product** of Rel is the functor

$$\times : \text{Rel} \times \text{Rel} \rightarrow \text{Rel}$$

where

- *Action on Objects.* For each $A, B \in \text{Obj}(\text{Rel})$, we have

$$\times(A, B) \stackrel{\text{def}}{=} A \times B,$$

where $A \times B$ is the Cartesian product of sets of [Definition 2.1.3.1](#).

- *Action on Morphisms.* For each $(A, C), (B, D) \in \text{Obj}(\text{Rel} \times \text{Rel})$, the action on morphisms

$$\times_{(A,C),(B,D)} : \text{Rel}(A, B) \times \text{Rel}(C, D) \rightarrow \text{Rel}(A \times C, B \times D)$$

of \times is given by sending a pair of morphisms (R, S) of the form

$$\begin{aligned} R &: A \rightarrow B, \\ S &: C \rightarrow D \end{aligned}$$

to the relation

$$R \times S : A \times C \rightarrow B \times D$$

of [Definition 6.3.9.1](#).

00JW 5.2.2.2 The Monoidal Unit

00JX DEFINITION 5.2.2.2 ► THE MONOIDAL UNIT OF Rel

The **monoidal unit** of Rel is the functor

$$\mathbb{1}^{\text{Rel}} : \text{pt} \rightarrow \text{Rel}$$

picking the set

$$\mathbb{1}_{\text{Rel}} \stackrel{\text{def}}{=} \text{pt}$$

of Rel.


00JY 5.2.2.3 The Associator

00JZ DEFINITION 5.2.2.3 ► THE ASSOCIATOR OF Rel

The **associator** of Rel is the natural isomorphism

$$\alpha^{\text{Rel}} : \times \circ ((\times) \times \text{id}) \xrightarrow{\sim} \times \circ (\text{id} \times (\times)) \circ \alpha_{\text{Rel}, \text{Rel}, \text{Rel}}^{\text{Cats}}$$

as in the diagram



whose component

$$\alpha_{A,B,C}^{\text{Rel}} : (A \times B) \times C \rightarrow A \times (B \times C)$$

at $A, B, C \in \text{Obj}(\text{Rel})$ is the relation defined by declaring


$$((a, b), c) \sim_{\alpha_{A,B,C}^{\text{Rel}}} (a', (b', c'))$$

iff $a = a', b = b',$ and $c = c'.$

00K0 5.2.2.4 The Left Unitor

00K1 DEFINITION 5.2.2.4 ► THE LEFT UNITOR OF Rel

The **left unitor** of Rel is the natural isomorphism



whose component

$$\lambda_A^{\text{Rel}} : \mathbb{1}_{\text{Rel}} \times A \rightarrow A$$

at A is defined by declaring

$$(\star, a) \sim_{\lambda_A^{\text{Rel}}} b$$

iff $a = b.$

00K2 **5.2.2.5 The Right Unitor**

00K3 **DEFINITION 5.2.2.5 ► THE RIGHT UNITOR OF Rel**

The **right unitor** of Rel is the natural isomorphism

$$\rho^{\text{Rel}} : \times \circ (\text{id} \times \mathbb{1}^{\text{Rel}}) \xrightarrow{\sim} \rho_{\text{Rel}}^{\text{Cats}_2},$$

whose component

$$\rho_A^{\text{Rel}} : A \times \mathbb{1}_{\text{Rel}} \rightarrow A$$

at A is defined by declaring

$$(a, \star) \sim_{\rho_A^{\text{Rel}}} b$$

iff $a = b$.

00K4 **5.2.2.6 The Symmetry**

00K5 **DEFINITION 5.2.2.6 ► THE SYMMETRY OF Rel**

The **symmetry** of Rel is the natural isomorphism

$$\sigma^{\text{Rel}} : \times \xrightarrow{\sim} \times \circ \sigma_{\text{Rel,Rel}}^{\text{Cats}_2},$$

whose component

$$\sigma_{A,B}^{\text{Rel}} : A \times B \rightarrow B \times A$$

at (A, B) is defined by declaring

$$(a, b) \sim_{\sigma_{A,B}^{\text{Rel}}} (b', a')$$

iff $a = a'$ and $b = b'$.

00K6 5.2.2.7 The Internal Hom

00K7 DEFINITION 5.2.2.7 ► THE INTERNAL HOM OF REL

The **internal Hom of Rel** is the functor

$$\text{Rel} : \text{Rel}^{\text{op}} \times \text{Rel} \rightarrow \text{Rel}$$

defined

- On objects by sending $A, B \in \text{Obj}(\text{Rel})$ to the set $\text{Rel}(A, B)$ of **Item 1** of **Definition 5.1.1.3**.
- On morphisms by pre/post-composition defined as in **Definition 6.3.12.1**.

00K8 PROPOSITION 5.2.2.8 ► PROPERTIES OF THE INTERNAL HOM OF REL

Let $A, B, C \in \text{Obj}(\text{Rel})$.

00K9 1. *Adjointness*. We have adjunctions

$$(A \times - \dashv \text{Rel}(A, -)) : \text{Rel} \begin{array}{c} \xrightarrow{A \times -} \\ \perp \\ \xleftarrow{\text{Rel}(A, -)} \end{array} \text{Rel},$$

$$(- \times B \dashv \text{Rel}(B, -)) : \text{Rel} \begin{array}{c} \xrightarrow{- \times B} \\ \perp \\ \xleftarrow{\text{Rel}(B, -)} \end{array} \text{Rel},$$

witnessed by bijections

$$\text{Rel}(A \times B, C) \cong \text{Rel}(A, \text{Rel}(B, C)),$$

$$\text{Rel}(A \times B, C) \cong \text{Rel}(B, \text{Rel}(A, C)),$$


natural in $A, B, C \in \text{Obj}(\text{Rel})$.

PROOF 5.2.2.9 ► PROOF OF PROPOSITION 5.2.2.8

Item 1: Adjointness

Indeed, we have

$$\begin{aligned}\text{Rel}(A \times B, C) &\stackrel{\text{def}}{=} \text{Sets}(A \times B \times C, \{\text{true}, \text{false}\}) \\ &\stackrel{\text{def}}{=} \text{Rel}(A, B \times C) \\ &\stackrel{\text{def}}{=} \text{Rel}(A, \text{Rel}(B, C)),\end{aligned}$$

and similarly for the bijection $\text{Rel}(A \times B, C) \cong \text{Rel}(B, \text{Rel}(A, C))$. 

00KA 5.2.2.8 The Closed Symmetric Monoidal Category of Relations

00KB

PROPOSITION 5.2.2.10 ► THE CLOSED SYMMETRIC MONOIDAL CATEGORY OF RELATIONS

The category Rel admits a closed symmetric monoidal category structure consisting of¹

- *The Underlying Category.* The category Rel of sets and relations of [Definition 5.2.1.1](#).
- *The Monoidal Product.* The functor

$$\times: \text{Rel} \times \text{Rel} \rightarrow \text{Rel}$$

of [Definition 5.2.2.1](#).

- *The Internal Hom.* The internal Hom functor

$$\mathbf{Rel}: \text{Rel}^{\text{op}} \times \text{Rel} \rightarrow \text{Rel}$$

of [Definition 5.2.2.7](#).

- *The Monoidal Unit.* The functor

$$\mathbb{1}^{\text{Rel}}: \text{pt} \rightarrow \text{Rel}$$

of [Definition 5.2.2.2](#).

- *The Associators.* The natural isomorphism

$$\alpha^{\text{Rel}}: \times \circ (\times \times \text{id}_{\text{Rel}}) \xrightarrow{\sim} \times \circ (\text{id}_{\text{Rel}} \times \times) \circ \alpha_{\text{Rel}, \text{Rel}, \text{Rel}}^{\text{Cats}}$$

of [Definition 5.2.2.3](#).

- *The Left Unitors.* The natural isomorphism

$$\lambda^{\mathbf{Rel}} : \times \circ (\mathbb{1}^{\mathbf{Rel}} \times \text{id}_{\mathbf{Rel}}) \xrightarrow{\sim} \lambda_{\mathbf{Rel}}^{\mathbf{Cats}_2}$$

of [Definition 5.2.2.4](#).

- *The Right Unitors.* The natural isomorphism


$$\rho^{\mathbf{Rel}} : \times \circ (\text{id} \times \mathbb{1}^{\mathbf{Rel}}) \xrightarrow{\sim} \rho_{\mathbf{Rel}}^{\mathbf{Cats}_2}$$

of [Definition 5.2.2.5](#).

- *The Symmetry.* The natural isomorphism

$$\sigma^{\mathbf{Rel}} : \times \xrightarrow{\sim} \times \circ \sigma_{\mathbf{Rel}, \mathbf{Rel}}^{\mathbf{Cats}_2}$$

of [Definition 5.2.2.6](#).

 *Warning:* This is not a Cartesian monoidal structure, as the product on \mathbf{Rel} is in fact given by the disjoint union of sets; see ??.

END TEXTDBEND

PROOF 5.2.2.11 ► PROOF OF PROPOSITION 5.2.2.10

Omitted. 

00KC 5.2.3 The 2-Category of Relations

00KD DEFINITION 5.2.3.1 ► THE 2-CATEGORY OF RELATIONS

The **2-category of relations** is the locally posetal 2-category \mathbf{Rel} where

- *Objects.* The objects of \mathbf{Rel} are sets.
- *Hom-Objects.* For each $A, B \in \text{Obj}(\mathbf{Sets})$, we have

$$\begin{aligned} \text{Hom}_{\mathbf{Rel}}(A, B) &\stackrel{\text{def}}{=} \mathbf{Rel}(A, B) \\ &\stackrel{\text{def}}{=} (\mathbf{Rel}(A, B), \subset). \end{aligned}$$

- *Identities.* For each $A \in \text{Obj}(\mathbf{Rel})$, the unit map

$$\mathbb{1}_A^{\mathbf{Rel}} : \text{pt} \rightarrow \mathbf{Rel}(A, A)$$

of \mathbf{Rel} at A is defined by

$$\mathrm{id}_A^{\mathbf{Rel}} \stackrel{\mathrm{def}}{=} \chi_A(-1, -2),$$

where $\chi_A(-1, -2)$ is the characteristic relation of A of [Item 3 of Definition 2.4.1.1](#).

- *Composition.* For each $A, B, C \in \mathrm{Obj}(\mathbf{Rel})$, the composition map¹

$$\circ_{A,B,C}^{\mathbf{Rel}} : \mathbf{Rel}(B, C) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, C)$$

of \mathbf{Rel} at (A, B, C) is defined by

$$S \circ_{A,B,C}^{\mathbf{Rel}} R \stackrel{\mathrm{def}}{=} S \diamond R$$

for each $(S, R) \in \mathbf{Rel}(B, C) \times \mathbf{Rel}(A, B)$, where $S \diamond R$ is the composition of S and R of [Definition 6.3.12.1](#).

¹Note that this is indeed a morphism of posets: given relations $R_1, R_2 \in \mathbf{Rel}(A, B)$ and $S_1, S_2 \in \mathbf{Rel}(B, C)$ such that

$$\begin{aligned} R_1 &\subset R_2, \\ S_1 &\subset S_2, \end{aligned}$$

we have also $S_1 \diamond R_1 \subset S_2 \diamond R_2$.

00KE 5.2.4 The Double Category of Relations

00KF 5.2.4.1 The Double Category of Relations

00KG DEFINITION 5.2.4.1 ► THE DOUBLE CATEGORY OF RELATIONS

The **double category of relations** is the locally posetal double category $\mathbf{Rel}^{\mathrm{dbl}}$ where

- *Objects.* The objects of $\mathbf{Rel}^{\mathrm{dbl}}$ are sets.
- *Vertical Morphisms.* The vertical morphisms of $\mathbf{Rel}^{\mathrm{dbl}}$ are maps of sets $f: A \rightarrow B$.
- *Horizontal Morphisms.* The horizontal morphisms of $\mathbf{Rel}^{\mathrm{dbl}}$ are relations $R: A \rightarrow X$.

- 2-Morphisms. A 2-cell

$$\begin{array}{ccc}
 A & \xrightarrow{R} & B \\
 f \downarrow & \Downarrow \alpha & \downarrow g \\
 X & \xrightarrow{S} & Y
 \end{array}$$

of Rel^{dbl} is either non-existent or an inclusion of relations of the form

$$R \subset S \circ (f \times g), \quad \begin{array}{ccc}
 A \times B & \xrightarrow{R} & \{\text{true}, \text{false}\} \\
 f \times g \downarrow & \subset & \downarrow \text{id}_{\{\text{true}, \text{false}\}} \\
 X \times Y & \xrightarrow{S} & \{\text{true}, \text{false}\}.
 \end{array}$$

- *Horizontal Identities.* The horizontal unit functor of Rel^{dbl} is the functor of [Definition 5.2.4.2](#).
- *Vertical Identities.* For each $A \in \text{Obj}(\text{Rel}^{\text{dbl}})$, we have

$$\text{id}_A^{\text{Rel}^{\text{dbl}}} \stackrel{\text{def}}{=} \text{id}_A.$$

- *Identity 2-Morphisms.* For each horizontal morphism $R: A \rightarrow B$ of Rel^{dbl} , the identity 2-morphism

$$\begin{array}{ccc}
 A & \xrightarrow{R} & B \\
 \text{id}_A \downarrow & \Downarrow \text{id}_R & \downarrow \text{id}_B \\
 A & \xrightarrow{R} & B
 \end{array}$$

of R is the identity inclusion

$$R \subset R, \quad \begin{array}{ccc}
 B \times A & \xrightarrow{R} & \{\text{true}, \text{false}\} \\
 \text{id}_B \times \text{id}_A \downarrow & \subset & \downarrow \text{id}_{\{\text{true}, \text{false}\}} \\
 B \times A & \xrightarrow{R} & \{\text{true}, \text{false}\}.
 \end{array}$$

- *Horizontal Composition.* The horizontal composition functor of Rel^{dbl} is the functor of [Definition 5.2.4.3](#).
- *Vertical Composition of 1-Morphisms.* For each composable pair $A \xrightarrow{F} B \xrightarrow{G} C$ of vertical morphisms of Rel^{dbl} , i.e. maps of sets, we have

$$g \circ^{\text{Rel}^{\text{dbl}}} f \stackrel{\text{def}}{=} g \circ f.$$

- *Vertical Composition of 2-Morphisms.* The vertical composition of 2-morphisms in Rel^{dbl} is defined as in [Definition 5.2.4.4](#).
- *Associators.* The associators of Rel^{dbl} is defined as in [Definition 5.2.4.5](#).
- *Left Unitors.* The left unitors of Rel^{dbl} is defined as in [Definition 5.2.4.6](#).
- *Right Unitors.* The right unitors of Rel^{dbl} is defined as in [Definition 5.2.4.7](#).

00KH 5.2.4.2 Horizontal Identities

00KJ DEFINITION 5.2.4.2 ► THE HORIZONTAL IDENTITIES OF Rel^{dbl}

The **horizontal unit functor** of Rel^{dbl} is the functor

$$\mathbb{1}^{\text{Rel}^{\text{dbl}}} : \text{Rel}_0^{\text{dbl}} \rightarrow \text{Rel}_1^{\text{dbl}}$$

of Rel^{dbl} is the functor where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Rel}_0^{\text{dbl}})$, we have

$$\mathbb{1}_A \stackrel{\text{def}}{=} \chi_A(-1, -2).$$

- *Action on Morphisms.* For each vertical morphism $f : A \rightarrow B$ of Rel^{dbl} , i.e. each map of sets f from A to B , the identity 2-morphism

$$\begin{array}{ccc} A & \xrightarrow{\mathbb{1}_A} & A \\ f \downarrow & \parallel & \downarrow f \\ B & \xrightarrow{\mathbb{1}_B} & B \end{array}$$

of f is the inclusion

$$\chi_B \circ (f \times f) \subset \chi_A, \quad \begin{array}{ccc} A \times A & \xrightarrow{\chi_A(-1,-2)} & \{\text{true}, \text{false}\} \\ f \times f \downarrow & \subset & \downarrow \text{id}_{\{\text{true}, \text{false}\}} \\ B \times B & \xrightarrow{\chi_B(-1,-2)} & \{\text{true}, \text{false}\} \end{array}$$

of **Item 1** of **Proposition 2.4.1.3**.

00KK 5.2.4.3 Horizontal Composition

00KL DEFINITION 5.2.4.3 ► THE HORIZONTAL COMPOSITION OF Rel^{dbl}

The **horizontal composition functor** of Rel^{dbl} is the functor

$$\odot^{\text{Rel}^{\text{dbl}}} : \text{Rel}_1^{\text{dbl}} \times_{\text{Rel}_0^{\text{dbl}}} \text{Rel}_1^{\text{dbl}} \rightarrow \text{Rel}_1^{\text{dbl}}$$

of Rel^{dbl} is the functor where

- *Action on Objects.* For each composable pair $A \xrightarrow{R} B \xrightarrow{S} C$ of horizontal morphisms of Rel^{dbl} , we have

$$S \odot R \stackrel{\text{def}}{=} S \diamond R,$$

where $S \diamond R$ is the composition of R and S of **Definition 6.3.12.1**.

- *Action on Morphisms.* For each horizontally composable pair

$$\begin{array}{ccc} A & \xrightarrow{R} & B \\ f \downarrow & \Downarrow \alpha & \downarrow g \\ X & \xrightarrow{T} & Y \end{array} \quad \begin{array}{ccc} B & \xrightarrow{S} & C \\ g \downarrow & \Downarrow \beta & \downarrow h \\ Y & \xrightarrow{U} & Z \end{array}$$

of 2-morphisms of Rel^{dbl} , i.e. for each pair

$$\begin{array}{ccc} A \times B & \xrightarrow{R} & \{\text{true}, \text{false}\} \\ f \times g \downarrow & \subset & \downarrow \text{id}_{\{\text{true}, \text{false}\}} \\ X \times Y & \xrightarrow{T} & \{\text{true}, \text{false}\} \end{array} \quad \begin{array}{ccc} B \times C & \xrightarrow{S} & \{\text{true}, \text{false}\} \\ g \times h \downarrow & \subset & \downarrow \text{id}_{\{\text{true}, \text{false}\}} \\ Y \times Z & \xrightarrow{U} & \{\text{true}, \text{false}\} \end{array}$$

of inclusions of relations, the horizontal composition

$$\begin{array}{ccc}
 A & \xrightarrow{S \circ R} & C \\
 f \downarrow & \parallel & \downarrow h \\
 & \beta \circ \alpha & \\
 & \Downarrow & \\
 X & \xrightarrow{U \circ T} & Z
 \end{array}$$

of α and β is the inclusion of relations¹

$$\begin{array}{ccc}
 A \times C & \xrightarrow{S \circ R} & \{\text{true}, \text{false}\} \\
 f \times h \downarrow & \subset & \downarrow \text{id}_{\{\text{true}, \text{false}\}} \\
 X \times Z & \xrightarrow{U \circ T} & \{\text{true}, \text{false}\}.
 \end{array}$$

$(U \circ T) \circ (f \times h) \subset (S \circ R)$

¹This is justified by noting that, given $(a, c) \in A \times C$, the statement

– We have $a \sim_{(U \circ T) \circ (f \times h)} c$, i.e. $f(a) \sim_{U \circ T} h(c)$, i.e. there exists some $y \in Y$ such that:

1. We have $f(a) \sim_T y$;
2. We have $y \sim_U h(c)$;

is implied by the statement

– We have $a \sim_{S \circ R} c$, i.e. there exists some $b \in B$ such that:

1. We have $a \sim_R b$;
2. We have $b \sim_S c$;


since:

- If $a \sim_R b$, then $f(a) \sim_T g(b)$, as $T \circ (f \times g) \subset R$;
- If $b \sim_S c$, then $g(b) \sim_U h(c)$, as $U \circ (g \times h) \subset S$.


00KM 5.2.4.4 Vertical Composition of 2-Morphisms

00KN DEFINITION 5.2.4.4 ► THE VERTICAL COMPOSITION OF 2-MORPHISMS IN Rel^{dbl}


The **vertical composition** in Rel^{dbl} is defined as follows: for each vertically composable pair




of 2-morphisms of Rel^{dbl} , i.e. for each each pair




of inclusions of relations, we define the vertical composition



of α and β as the inclusion of relations



given by the pasting of inclusions¹



¹This is justified by noting that, given $(a, x) \in A \times X$, the statement

- We have $h(f(a)) \sim_T k(g(x))$;

is implied by the statement

- We have $a \sim_R x$;

since

- If $a \sim_R x$, then $f(a) \sim_S g(x)$, as $S \circ (f \times g) \subset R$;
- If $b \sim_S y$, then $h(b) \sim_T k(y)$, as $T \circ (h \times k) \subset S$, and thus, in particular:
 - If $f(a) \sim_S g(x)$, then $h(f(a)) \sim_T k(g(x))$.

00KP 5.2.4.5 The Associators

00KQ DEFINITION 5.2.4.5 ► THE ASSOCIATORS OF Rel^{dbl}

For each composable triple

$$A \xrightarrow{R} B \xrightarrow{S} C \xrightarrow{T} D$$

of horizontal morphisms of Rel^{dbl} , the component

$$\alpha_{T,S,R}^{\text{Rel}^{\text{dbl}}} : (T \circ S) \circ R \xrightarrow{\sim} T \circ (S \circ R),$$

of the associator of Rel^{dbl} at (R, S, T) is the identity inclusion¹

$$(T \circ S) \circ R = T \circ (S \circ R)$$

¹This is justified by **Item 2** of **Proposition 6.3.12.3**.

00KR 5.2.4.6 The Left Unitors

00KS

DEFINITION 5.2.4.6 ▶ THE LEFT UNITORS OF Rel^{dbl}

For each horizontal morphism $R: A \rightarrow B$ of Rel^{dbl} , the component

$$\lambda_R^{\text{Rel}^{\text{dbl}}} : \mathbb{1}_B \odot R \xrightarrow{\sim} R,$$

$$\begin{array}{ccccc} A & \xrightarrow{R} & B & \xrightarrow{\mathbb{1}_B} & B \\ \text{id}_A \downarrow & & \lambda_R^{\text{Rel}^{\text{dbl}}} \downarrow & & \downarrow \text{id}_B \\ A & \xrightarrow{R} & B & & B \end{array}$$

of the left unitor of Rel^{dbl} at R is the identity inclusion¹

$$R = \chi_B \diamond R,$$

$$\begin{array}{ccc} A \times B & \xrightarrow{\chi_B \diamond R} & \{\text{true}, \text{false}\} \\ \parallel & \cong & \downarrow \text{id}_{\{\text{true}, \text{false}\}} \\ A \times B & \xrightarrow{R} & \{\text{true}, \text{false}\}. \end{array}$$

¹This is justified by [Item 3](#) of [Proposition 6.3.12.3](#).

00KT 5.2.4.7 The Right Unitors

00KU

DEFINITION 5.2.4.7 ▶ THE RIGHT UNITORS OF Rel^{dbl}

For each horizontal morphism $R: A \rightarrow B$ of Rel^{dbl} , the component

$$\rho_R^{\text{Rel}^{\text{dbl}}} : R \odot \mathbb{1}_A \xrightarrow{\sim} R,$$

$$\begin{array}{ccccc} A & \xrightarrow{\mathbb{1}_A} & A & \xrightarrow{R} & B \\ \text{id}_A \downarrow & & \rho_R^{\text{Rel}^{\text{dbl}}} \downarrow & & \downarrow \text{id}_B \\ A & \xrightarrow{R} & B & & B \end{array}$$

of the right unitor of Rel^{dbl} at R is the identity inclusion¹

$$R = R \diamond \chi_A,$$

$$\begin{array}{ccc} A \times B & \xrightarrow{R \diamond \chi_A} & \{\text{true}, \text{false}\} \\ \parallel & \cong & \downarrow \text{id}_{\{\text{true}, \text{false}\}} \\ A \times B & \xrightarrow{R} & \{\text{true}, \text{false}\}. \end{array}$$

¹This is justified by [Item 3](#) of [Proposition 6.3.12.3](#).

00KV 5.3 Properties of the 2-Category of Relations

00KW 5.3.1 Self-Duality

00KX PROPOSITION 5.3.1.1 ► SELF-DUALITY FOR THE (2-)CATEGORY OF RELATIONS

The (2-)category of relations is self-dual:

- 00KY 1. *Self-Duality I*. We have an isomorphism

$$\mathbf{Rel}^{\text{op}} \stackrel{\text{eq.}}{\cong} \mathbf{Rel}$$

of categories.

- 00KZ 2. *Self-Duality II*. We have a 2-isomorphism

$$\mathbf{Rel}^{\text{op}} \stackrel{\text{eq.}}{\cong} \mathbf{Rel}$$

of 2-categories.

PROOF 5.3.1.2 ► PROOF OF PROPOSITION 5.3.1.1

Item 1: Self-Duality I

We claim that the functor

$$F: \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{Rel}$$

given by the identity on objects and by $R \mapsto R^\dagger$ on morphisms is an isomorphism of categories.

By [Item 1 of Proposition 8.5.8.3](#), it suffices to show that F is bijective on objects (which is clear) and fully faithful. Indeed, the map

$$(-)^\dagger: \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(B, A)$$

defined by the assignment $R \mapsto R^\dagger$ is a bijection by [Item 5 of Proposition 6.3.11.3](#), showing F to be fully faithful.

Item 2: Self-Duality II


We claim that the 2-functor

$$F: \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{Rel}$$

given by the identity on objects, by $R \mapsto R^\dagger$ on morphisms, and by preserving inclusions on 2-morphisms via [Item 1 of Proposition 6.3.11.3](#), is an isomorphism of categories.

By ?? of ??, it suffices to show that F is:

- Bijective on objects, which is clear.
- Bijective on 1-morphisms, which was shown in **Item 1**.
- Bijective on 2-morphisms, which follows from **Item 1** of **Proposition 6.3.11.3**.

Thus F is indeed a 2-isomorphism of categories. 

00L0 5.3.2 Isomorphisms and Equivalences in Rel

Let $R: A \rightarrow B$ be a relation from A to B .

00L1 PROPOSITION 5.3.2.1 ► ISOMORPHISMS AND EQUIVALENCES IN Rel

The following conditions are equivalent:

- 00L2 1. The relation $R: A \rightarrow B$ is an equivalence in **Rel**, i.e.:
- (★) There exists a relation $R^{-1}: B \rightarrow A$ from B to A together with isomorphisms

$$\begin{aligned} R^{-1} \diamond R &\cong \chi_A, \\ R \diamond R^{-1} &\cong \chi_B. \end{aligned}$$

- 00L3 2. The relation $R: A \rightarrow B$ is an isomorphism in **Rel**, i.e.:
- (★) There exists a relation $R^{-1}: B \rightarrow A$ from B to A such that we have

$$\begin{aligned} R^{-1} \diamond R &= \chi_A, \\ R \diamond R^{-1} &= \chi_B. \end{aligned}$$

- 00L4 3. There exists a bijection $f: A \xrightarrow{\cong} B$ with $R = \text{Gr}(f)$.

PROOF 5.3.2.2 ► PROOF OF PROPOSITION 5.3.2.1

We claim that **Items 1 to 3** are indeed equivalent:

- **Item 1** \iff **Item 2**: This follows from the fact that **Rel** is locally posetal, so that natural isomorphisms and equalities of 1-morphisms in **Rel** coincide.

- **Item 2** \implies **Item 3**: The equalities in **Item 2** imply $R \dashv R^{-1}$, and thus by **Proposition 5.3.3.1**, there exists a function $f_R: A \rightarrow B$ associated to R , where, for each $a \in A$, the image $f_R(a)$ of a by f_R is the unique element of $R(a)$, which implies $R = \text{Gr}(f_R)$ in particular. Furthermore, we have $R^{-1} = f_R^{-1}$ (as in **Definition 6.3.2.1**). The conditions from **Item 2** then become the following:

$$\begin{aligned} f_R^{-1} \diamond f_R &= \chi_A, \\ f_R \diamond f_R^{-1} &= \chi_B. \end{aligned}$$


All that is left is to show then is that f_R is a bijection:

- *The Function f_R Is Injective.* Let $a, b \in A$ and suppose that $f_R(a) = f_R(b)$. Since $a \sim_R f_R(a)$ and $f_R(a) = f_R(b) \sim_{R^{-1}} b$, the condition $f_R^{-1} \diamond f_R = \chi_A$ implies that $a = b$, showing f_R to be injective.
- *The Function f_R Is Surjective.* Let $b \in B$. Applying the condition $f_R \diamond f_R^{-1} = \chi_B$ to (b, b) , it follows that there exists some $a \in A$ such that $f_R^{-1}(b) = a$ and $f_R(a) = b$. This shows f_R to be surjective.
- **Item 3** \implies **Item 2**: By **Item 2** of **Proposition 6.3.1.2**, we have an adjunction $\text{Gr}(f) \dashv f^{-1}$, giving inclusions

$$\begin{aligned} \chi_A &\subset f^{-1} \diamond \text{Gr}(f), \\ \text{Gr}(f) \diamond f^{-1} &\subset \chi_B. \end{aligned}$$

We claim the reverse inclusions are also true:

- $f^{-1} \diamond \text{Gr}(f) \subset \chi_A$: This is equivalent to the statement that if $f(a) = b$ and $f^{-1}(b) = a'$, then $a = a'$, which follows from the injectivity of f .
- $\chi_B \subset \text{Gr}(f) \diamond f^{-1}$: This is equivalent to the statement that given $b \in B$ there exists some $a \in A$ such that $f^{-1}(b) = a$ and $f(a) = b$, which follows from the surjectivity of f .

This finishes the proof. 

00L5 5.3.3 Adjunctions in Rel

Let A and B be sets.

00L6

PROPOSITION 5.3.3.1 ► ADJUNCTIONS IN Rel

We have a natural bijection

$$\left\{ \begin{array}{l} \text{Adjunctions in } \mathbf{Rel} \\ \text{from } A \text{ to } B \end{array} \right\} \cong \left\{ \begin{array}{l} \text{Functions} \\ \text{from } A \text{ to } B \end{array} \right\},$$

with every adjunction in **Rel** being of the form $\text{Gr}(f) \dashv f^{-1}$ for some function f .

PROOF 5.3.3.2 ► PROOF OF PROPOSITION 5.3.3.1

We proceed step by step:

1. *From Adjunctions in Rel to Functions.* An adjunction in **Rel** from A to B consists of a pair of relations

$$R: A \rightarrow B,$$

$$S: B \rightarrow A,$$

together with inclusions

$$\chi_A \subset S \diamond R,$$

$$R \diamond S \subset \chi_B.$$

We claim that these conditions imply that R is total and functional, i.e. that $R(a)$ is a singleton for each $a \in A$:

- (a) *$R(a)$ Has an Element.* Given $a \in A$, since $\chi_A \subset S \diamond R$, we must have $\{a\} \subset S(R(a))$, implying that there exists some $b \in B$ such that $a \sim_R b$ and $b \sim_S a$, and thus $R(a) \neq \emptyset$, as $b \in R(a)$.
- (b) *$R(a)$ Has No More Than One Element.* Suppose that we have $a \sim_R b$ and $a \sim_R b'$ for $b, b' \in B$. We claim that $b = b'$:
 - i. Since $\chi_A \subset S \diamond R$, there exists some $k \in B$ such that $a \sim_R k$ and $k \sim_S a$.
 - ii. Since $R \diamond S \subset \chi_B$, if $b'' \sim_S a'$ and $a' \sim_R b'''$, then $b'' = b'''$.
 - iii. Applying the above to $b'' = k$, $b''' = b$, and $a' = a$, since $k \sim_S a$ and $a \sim_R b'$, we have $k = b$.
 - iv. Similarly $k = b'$.
 - v. Thus $b = b'$.

Together, the above two items show $R(a)$ to be a singleton, being thus given by $\text{Gr}(f)$ for some function $f: A \rightarrow B$, which gives a map

$$\left\{ \begin{array}{l} \text{Adjunctions in } \mathbf{Rel} \\ \text{from } A \text{ to } B \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{Functions} \\ \text{from } A \text{ to } B \end{array} \right\}.$$

Moreover, by uniqueness of adjoints (?? of ??), this implies also that $S = f^{-1}$.

2. *From Functions to Adjunctions in Rel.* By **Item 2** of **Proposition 6.3.1.2**, every function $f: A \rightarrow B$ gives rise to an adjunction $\text{Gr}(f) \dashv f^{-1}$ in \mathbf{Rel} , giving a map

$$\left\{ \begin{array}{l} \text{Functions} \\ \text{from } A \text{ to } B \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{Adjunctions in } \mathbf{Rel} \\ \text{from } A \text{ to } B \end{array} \right\}.$$

3. *Invertibility: From Functions to Adjunctions Back to Functions.* We need to show that starting with a function $f: A \rightarrow B$, passing to $\text{Gr}(f) \dashv f^{-1}$, and then passing again to a function gives f again. This is clear however, since we have $a \sim_{\text{Gr}(f)} b$ iff $f(a) = b$.
4. *Invertibility: From Adjunctions to Functions Back to Adjunctions.* We need to show that, given an adjunction $R \dashv S$ in \mathbf{Rel} giving rise to a function $f_{R,S}: A \rightarrow B$, we have

$$\begin{aligned} \text{Gr}(f_{R,S}) &= R, \\ f_{R,S}^{-1} &= S. \end{aligned}$$

We check these explicitly:

- $\text{Gr}(f_{R,S}) = R$. We have

$$\begin{aligned} \text{Gr}(f_{R,S}) &\stackrel{\text{def}}{=} \{(a, f_{R,S}(a)) \in A \times B \mid a \in A\} \\ &\stackrel{\text{def}}{=} \{(a, R(a)) \in A \times B \mid a \in A\} \\ &= R. \end{aligned}$$

- $f_{R,S}^{-1} = S$. We first claim that, given $a \in A$ and $b \in B$, the following conditions are equivalent:

- We have $a \sim_R b$.
- We have $b \sim_S a$.


Indeed:

- If $a \sim_R b$, then $b \sim_S a$: Since $\chi_A \subset S \diamond R$, there exists $k \in B$ such that $a \sim_R k$ and $k \sim_S a$, but since $a \sim_R b$ and R is functional, we have $k = b$ and thus $b \sim_S a$.
- If $b \sim_S a$, then $a \sim_R b$: First note that since R is total we have $a \sim_R b'$ for some $b' \in B$. Now, since $R \diamond S \subset \chi_B$, $b \sim_S a$, and $a \sim_R b'$, we have $b = b'$, and thus $a \sim_R b$.

Having shown this, we now have

$$\begin{aligned} f_{R,S}^{-1}(b) &\stackrel{\text{def}}{=} \{a \in A \mid f_{R,S}(a) = b\} \\ &\stackrel{\text{def}}{=} \{a \in A \mid a \sim_R b\} \\ &= \{a \in A \mid b \sim_S a\} \\ &\stackrel{\text{def}}{=} S(b). \end{aligned}$$

for each $b \in B$, showing $f_{R,S}^{-1} = S$.

This finishes the proof. 

00L7 5.3.4 Monads in Rel

Let A be a set.

00L8 PROPOSITION 5.3.4.1 ► MONADS IN Rel

We have a natural identification¹

$$\left\{ \begin{array}{l} \text{Monads in} \\ \mathbf{Rel} \text{ on } A \end{array} \right\} \cong \{\text{Preorders on } A\}.$$

¹See also ?? for an extension of this correspondence to “relative monads in **Rel**”.

PROOF 5.3.4.2 ► PROOF OF PROPOSITION 5.3.4.1

A monad in **Rel** on A consists of a relation $R: A \rightarrow A$ together with maps

$$\begin{aligned} \mu_R: R \diamond R &\subset R, \\ \eta_R: \chi_A &\subset R \end{aligned}$$

making the diagrams

commute. However, since all morphisms involved are inclusions, the commutativity of the above diagrams is automatic, and hence all that is left is the data of the two maps μ_R and η_R , which correspond respectively to the following conditions:

1. For each $a, b, c \in A$, if $a \sim_R b$ and $b \sim_R c$, then $a \sim_R c$.
2. For each $a \in A$, we have $a \sim_R a$.

These are exactly the requirements for R to be a preorder (??). Conversely any preorder \preceq gives rise to a pair of maps μ_{\preceq} and η_{\preceq} , forming a monad on A . ■

00L9 5.3.5 Comonads in Rel

Let A be a set.

00LA PROPOSITION 5.3.5.1 ► COMONADS IN Rel

We have a natural identification

$$\left\{ \begin{array}{l} \text{Comonads in} \\ \mathbf{Rel} \text{ on } A \end{array} \right\} \cong \{\text{Subsets of } A\}.$$

PROOF 5.3.5.2 ► PROOF OF PROPOSITION 5.3.5.1

A comonad in **Rel** on A consists of a relation $R: A \dashv\vdash A$ together with maps

$$\begin{aligned} \Delta_R: R \subset R \diamond R, \\ \epsilon_R: R \subset \chi_A \end{aligned}$$

making the diagrams

commute. However, since all morphisms involved are inclusions, the commutativity of the above diagrams is automatic, and hence all that is left is the data of the two maps Δ_R and ϵ_R , which correspond respectively to the following conditions:

1. For each $a, b \in A$, if $a \sim_R b$, then there exists some $k \in A$ such that $a \sim_R k$ and $k \sim_R b$.
2. For each $a, b \in A$, if $a \sim_R b$, then $a = b$.

Taking $k = b$ in the first condition above shows it to be trivially satisfied, while the second condition implies $R \subset \Delta_A$, i.e. R must be a subset of A . Conversely, any subset U of A satisfies $U \subset \Delta_A$, defining a comonad as above. ▢

00LB 5.3.6 Co/Monoids in Rel

00LC **REMARK 5.3.6.1** ▶ **Co/MONOIDS IN Rel**

The monoids in **Rel** with respect to the Cartesian monoidal structure of **Proposition 5.2.2.10** are called *hypermonoids*, and their theory is explored in ???. Similarly, the comonoids in **Rel** are called *hypercomonoids*, and they are defined and studied in ???.

00LD 5.3.7 Monomorphisms in Rel

In this section we characterise the epimorphisms in the category Rel, following ???.

00LE

PROPOSITION 5.3.7.1 ► CHARACTERISATIONS OF MONOMORPHISMS IN Rel

Let $R: A \rightarrow B$ be a relation. The following conditions are equivalent:

00LF

1. The relation R is a monomorphism in Rel.

00LG

2. The direct image function

$$R_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

associated to R is injective.

00LH

3. The direct image with compact support function

$$R_! : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

associated to R is injective.

Moreover, if R is a monomorphism, then it satisfies the following condition, and the converse holds if R is total:

(★) For each $a, a' \in A$, if there exists some $b \in B$ such that

$$\begin{aligned} a &\sim_R b, \\ a' &\sim_R b, \end{aligned}$$

then $a = a'$.

PROOF 5.3.7.2 ► PROOF OF PROPOSITION 5.3.7.1

Firstly note that **Items 2** and **3** are equivalent by **Item 7** of **Proposition 6.4.1.3**. We then claim that **Items 1** and **2** are also equivalent:

• **Item 1** \implies **Item 2**: Let $U, V \in \mathcal{P}(A)$ and consider the diagram

$$\text{pt} \begin{array}{c} \xrightarrow{U} \\ \xrightarrow{V} \end{array} A \xrightarrow{R} B.$$

By **Remark 6.4.1.2**, we have

$$\begin{aligned} R_*(U) &= R \diamond U, \\ R_*(V) &= R \diamond V. \end{aligned}$$

Now, if $R \diamond U = R \diamond V$, i.e. $R_*(U) = R_*(V)$, then $U = V$ since R is assumed to be a monomorphism, showing R_* to be injective.

- *Item 2* \implies *Item 1*: Conversely, suppose that R_* is injective, consider the diagram

$$X \begin{array}{c} \xrightarrow{S} \\ \dashrightarrow \\ \xrightarrow{T} \end{array} A \xrightarrow{R} B,$$

and suppose that $R \diamond S = R \diamond T$. Note that, since R_* is injective, given a diagram of the form

$$\text{pt} \begin{array}{c} \xrightarrow{U} \\ \dashrightarrow \\ \xrightarrow{V} \end{array} A \xrightarrow{R} B,$$

if $R_*(U) = R \diamond U = R \diamond V = R_*(V)$, then $U = V$. In particular, for each $x \in X$, we may consider the diagram

$$\text{pt} \xrightarrow{[x]} X \begin{array}{c} \xrightarrow{S} \\ \dashrightarrow \\ \xrightarrow{T} \end{array} A \xrightarrow{R} B,$$

for which we have $R \diamond S \diamond [x] = R \diamond T \diamond [x]$, implying that we have

$$S(x) = S \diamond [x] = T \diamond [x] = T(x)$$

for each $x \in X$, implying $S = T$, and thus R is a monomorphism.

We can also prove this in a more abstract way, following [MSE 350788]:

- *Item 1* \implies *Item 2*: Assume that R is a monomorphism.
 - We first notice that the functor $\text{Rel}(\text{pt}, -) : \text{Rel} \rightarrow \text{Sets}$ maps R to R_* by Remark 6.4.1.2.
 - Since $\text{Rel}(\text{pt}, -)$ preserves all limits by ?? of ??, it follows by ?? of ?? that $\text{Rel}(\text{pt}, -)$ also preserves monomorphisms.
 - Since R is a monomorphism and $\text{Rel}(\text{pt}, -)$ maps R to R_* , it follows that R_* is also a monomorphism.
 - Since the monomorphisms in Sets are precisely the injections (?? of ??), it follows that R_* is injective.
- *Item 2* \implies *Item 1*: Assume that R_* is injective.
 - We first notice that the functor $\text{Rel}(\text{pt}, -) : \text{Rel} \rightarrow \text{Sets}$ maps R to R_* by Remark 6.4.1.2.
 - Since the monomorphisms in Sets are precisely the injections (?? of ??), it follows that R_* is a monomorphism.

- Since $\text{Rel}(\text{pt}, -)$ is faithful, it follows by ?? of ?? that $\text{Rel}(\text{pt}, -)$ reflects monomorphisms.
- Since R_* is a monomorphism and $\text{Rel}(\text{pt}, -)$ maps R to R_* , it follows that R is also a monomorphism.

Finally, we prove the second part of the statement. Assume that R is a monomorphism, let $a, a' \in A$ such that $a \sim_R b$ and $a' \sim_R b$ for some $b \in B$, and consider the diagram

$$\begin{array}{ccc} & [a] & \\ \text{pt} & \begin{array}{c} \xrightarrow{\quad} \\ \xrightarrow{\quad} \\ \xrightarrow{\quad} \end{array} & A \xrightarrow{R} B \\ & [a'] & \end{array}$$

Since $\star \sim_{[a]} a$ and $a \sim_R b$, we have $\star \sim_{R \diamond [a]} b$. Similarly, $\star \sim_{R \diamond [a']} b$. Thus $R \diamond [a] = R \diamond [a']$, and since R is a monomorphism, we have $[a] = [a']$, i.e. $a = a'$.

Conversely, assume the condition

- (\star) For each $a, a' \in A$, if there exists some $b \in B$ such that


$$\begin{aligned} a &\sim_R b, \\ a' &\sim_R b, \end{aligned}$$

then $a = a'$.

consider the diagram

$$\begin{array}{ccc} & S & \\ X & \begin{array}{c} \xrightarrow{\quad} \\ \xrightarrow{\quad} \\ \xrightarrow{\quad} \end{array} & A \xrightarrow{R} B \\ & T & \end{array}$$

and let $(x, a) \in S$. Since R is total and $a \in A$, there exists some $b \in B$ such that $a \sim_R b$. In this case, we have $x \sim_{R \diamond S} b$, and since $R \diamond S = R \diamond T$, we have also $x \sim_{R \diamond T} b$. Thus there must exist some $a' \in A$ such that $x \sim_T a'$ and $a' \sim_R b$. However, since $a, a' \sim_R b$, we must have $a = a'$, and thus $(x, a) \in T$ as well.

A similar argument shows that if $(x, a) \in T$, then $(x, a) \in S$, and thus $S = T$ and it follows that R is a monomorphism. 

00LJ 5.3.8 2-Categorical Monomorphisms in Rel

In this section we characterise (for now, some of) the 2-categorical monomorphisms in **Rel**, following [Section 9.1](#).

00LK **PROPOSITION 5.3.8.1 ► 2-CATEGORICAL MONOMORPHISMS IN Rel**

Let $R: A \rightarrow B$ be a relation.

00LL 1. *Representably Faithful Morphisms in Rel.* Every morphism of **Rel** is a representably faithful morphism.

00LM 2. *Representably Full Morphisms in Rel.* The following conditions are equivalent:

00LN (a) The morphism $R: A \rightarrow B$ is a representably full morphism.

00LP (b) For each pair of relations $S, T: X \rightarrow A$, the following condition is satisfied:

(★) If $R \diamond S \subset R \diamond T$, then $S \subset T$.

00LQ (c) The functor

$$R_*: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

is full.

00LR (d) For each $U, V \in \mathcal{P}(A)$, if $R_*(U) \subset R_*(V)$, then $U \subset V$.

00LS (e) The functor

$$R_!: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

is full.

00LT (f) For each $U, V \in \mathcal{P}(A)$, if $R_!(U) \subset R_!(V)$, then $U \subset V$.

00LU 3. *Representably Fully Faithful Morphisms in Rel.* Every representably full morphism in **Rel** is a representably fully faithful morphism.

PROOF 5.3.8.2 ► PROOF OF PROPOSITION 5.3.8.1

Item 1: Representably Faithful Morphisms in Rel

The relation R is a representably faithful morphism in **Rel** iff, for each $X \in \text{Obj}(\mathbf{Rel})$, the functor

$$R_*: \mathbf{Rel}(X, A) \rightarrow \mathbf{Rel}(X, B)$$

is faithful, i.e. iff the morphism

$$R_{*|S,T}: \text{Hom}_{\mathbf{Rel}(X,A)}(S, T) \rightarrow \text{Hom}_{\mathbf{Rel}(X,B)}(R \diamond S, R \diamond T)$$

is injective for each $S, T \in \text{Obj}(\mathbf{Rel}(X, A))$. However, $\text{Hom}_{\mathbf{Rel}(X, A)}(S, T)$ is either empty or a singleton, in either case of which the map $R_{*|S, T}$ is necessarily injective.

Item 2: Representably Full Morphisms in \mathbf{Rel}

We claim **Items 2a** to **2f** are indeed equivalent:

- **Item 2a** \iff **Item 2b**: This is simply a matter of unwinding definitions: The relation R is a representably full morphism in \mathbf{Rel} iff, for each $X \in \text{Obj}(\mathbf{Rel})$, the functor

$$R_* : \mathbf{Rel}(X, A) \rightarrow \mathbf{Rel}(X, B)$$

is full, i.e. iff the morphism

$$R_{*|S, T} : \text{Hom}_{\mathbf{Rel}(X, A)}(S, T) \rightarrow \text{Hom}_{\mathbf{Rel}(X, B)}(R \diamond S, R \diamond T)$$

is surjective for each $S, T \in \text{Obj}(\mathbf{Rel}(X, A))$, i.e. iff, whenever $R \diamond S \subset R \diamond T$, we also have $S \subset T$.

- **Item 2c** \iff **Item 2d**: This is also simply a matter of unwinding definitions: The functor

$$R_* : (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

is full iff, for each $U, V \in \mathcal{P}(A)$, the morphism

$$R_{*|U, V} : \text{Hom}_{\mathcal{P}(A)}(U, V) \rightarrow \text{Hom}_{\mathcal{P}(B)}(R_*(U), R_*(V))$$

is surjective, i.e. iff whenever $R_*(U) \subset R_*(V)$, we also necessarily have $U \subset V$.

- **Item 2e** \iff **Item 2f**: This is once again simply a matter of unwinding definitions, and proceeds exactly in the same way as in the proof of the equivalence between **Items 2c** and **2d** given above.
- **Item 2d** \implies **Item 2f**: Suppose that the following condition is true:

$$(\star) \text{ For each } U, V \in \mathcal{P}(A), \text{ if } R_*(U) \subset R_*(V), \text{ then } U \subset V.$$

We need to show that the condition

$$(\star) \text{ For each } U, V \in \mathcal{P}(A), \text{ if } R_!(U) \subset R_!(V), \text{ then } U \subset V.$$

is also true. We proceed step by step:

1. Suppose we have $U, V \in \mathcal{P}(A)$ with $R_l(U) \subset R_l(V)$.
2. By **Item 7** of **Proposition 6.4.4.4**, we have

$$\begin{aligned} R_l(U) &= B \setminus R_*(A \setminus U), \\ R_l(V) &= B \setminus R_*(A \setminus V). \end{aligned}$$

3. By **Item 1** of **Proposition 2.3.10.2** we have $R_*(A \setminus V) \subset R_*(A \setminus U)$.
4. By assumption, we then have $A \setminus V \subset A \setminus U$.
5. By **Item 1** of **Proposition 2.3.10.2** again, we have $U \subset V$.

· **Item 2f** \implies **Item 2d**: Suppose that the following condition is true:

(★) For each $U, V \in \mathcal{P}(A)$, if $R_l(U) \subset R_l(V)$, then $U \subset V$.

We need to show that the condition

(★) For each $U, V \in \mathcal{P}(A)$, if $R_*(U) \subset R_*(V)$, then $U \subset V$.

is also true. We proceed step by step:

1. Suppose we have $U, V \in \mathcal{P}(A)$ with $R_*(U) \subset R_*(V)$.
2. By **Item 7** of **Proposition 6.4.1.3**, we have

$$\begin{aligned} R_*(U) &= B \setminus R_l(A \setminus U), \\ R_*(V) &= B \setminus R_l(A \setminus V). \end{aligned}$$

3. By **Item 1** of **Proposition 2.3.10.2** we have $R_l(A \setminus V) \subset R_l(A \setminus U)$.
4. By assumption, we then have $A \setminus V \subset A \setminus U$.
5. By **Item 1** of **Proposition 2.3.10.2** again, we have $U \subset V$.

· **Item 2b** \implies **Item 2d**: Consider the diagram

$$X \begin{array}{c} \xrightarrow{S} \\ \dashrightarrow \\ \xrightarrow{T} \end{array} A \xrightarrow{R} B,$$

and suppose that $R \diamond S \subset R \diamond T$. Note that, by assumption, given a diagram of the form

$$\text{pt} \begin{array}{c} \xrightarrow{U} \\ \dashrightarrow \\ \xrightarrow{V} \end{array} A \xrightarrow{R} B,$$

if $R_*(U) = R \diamond U \subset R \diamond V = R_*(V)$, then $U \subset V$. In particular, for each $x \in X$, we may consider the diagram

$$\text{pt} \xrightarrow{[x]} X \begin{array}{c} \xrightarrow{S} \\ \xrightarrow{T} \end{array} A \xrightarrow{R} B,$$

for which we have $R \diamond S \diamond [x] \subset R \diamond T \diamond [x]$, implying that we have

$$S(x) = S \diamond [x] \subset T \diamond [x] = T(x)$$

for each $x \in X$, implying $S \subset T$.

• *Item 2d* \implies *Item 2b*: Let $U, V \in \mathcal{P}(A)$ and consider the diagram

$$\text{pt} \begin{array}{c} \xrightarrow{U} \\ \xrightarrow{V} \end{array} A \xrightarrow{R} B.$$


By [Remark 6.4.1.2](#), we have

$$R_*(U) = R \diamond U,$$

$$R_*(V) = R \diamond V.$$

Now, if $R_*(U) \subset R_*(V)$, i.e. $R \diamond U \subset R \diamond V$, then $U \subset V$ by assumption.

?: Fully Faithful Monomorphisms in **Rel**

This follows from [Items 1](#) and [2](#). 

00LV

QUESTION 5.3.8.3 ► BETTER CHARACTERISATIONS OF REPRESENTABLY FULL MORPHISMS IN **Rel**

[Item 2](#) of [Proposition 5.3.8.1](#) gives a characterisation of the representably full morphisms in **Rel**.

Are there other nice characterisations of these?

This question also appears as [\[MO 467527\]](#).

00LW 5.3.9 Epimorphisms in **Rel**

In this section we characterise the epimorphisms in the category **Rel**, following ??.

00LX **PROPOSITION 5.3.9.1 ► CHARACTERISATIONS OF EPIMORPHISMS IN Rel**

Let $R: A \rightarrow B$ be a relation. The following conditions are equivalent:

00LY 1. The relation R is an epimorphism in Rel.

00LZ 2. The weak inverse image function

$$R^{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

associated to R is injective.

00M0 3. The strong inverse image function

$$R_{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

associated to R is injective.

00M1 4. The function $R: A \rightarrow \mathcal{P}(B)$ is “surjective on singletons”:

(★) For each $b \in B$, there exists some $a \in A$ such that $R(a) = \{b\}$.

Moreover, if R is total and an epimorphism, then it satisfies the following equivalent conditions:

1. For each $b \in B$, there exists some $a \in A$ such that $a \sim_R b$.
2. We have $\text{Im}(R) = B$.

PROOF 5.3.9.2 ► PROOF OF PROPOSITION 5.3.9.1

Firstly note that **Items 2 and 3** are equivalent by **Item 7** of **Proposition 6.4.2.4**. We then claim that **Items 1 and 2** are also equivalent:

• **Item 1** \implies **Item 2**: Let $U, V \in \mathcal{P}(A)$ and consider the diagram

$$A \xrightarrow{R} B \begin{array}{c} \xrightarrow{U} \\ \xrightarrow{V} \end{array} \text{pt.}$$

By **Remark 6.4.1.2**, we have

$$R^{-1}(U) = U \diamond R,$$

$$R^{-1}(V) = V \diamond R.$$

Now, if $U \diamond R = V \diamond R$, i.e. $R^{-1}(U) = R^{-1}(V)$, then $U = V$ since R is assumed to be an epimorphism, showing R^{-1} to be injective.

- *Item 2* \implies *Item 1*: Conversely, suppose that R^{-1} is injective, consider the diagram

$$A \xrightarrow{R} B \begin{array}{c} \xrightarrow{S} \\ \xrightarrow{T} \end{array} X,$$

and suppose that $S \diamond R = T \diamond R$. Note that, since R^{-1} is injective, given a diagram of the form

$$A \xrightarrow{R} B \begin{array}{c} \xrightarrow{U} \\ \xrightarrow{V} \end{array} \text{pt},$$

if $R^{-1}(U) = U \diamond R = V \diamond R = R^{-1}(V)$, then $U = V$. In particular, for each $x \in X$, we may consider the diagram

$$A \xrightarrow{R} B \begin{array}{c} \xrightarrow{S} \\ \xrightarrow{T} \end{array} X \xrightarrow{[x]} \text{pt},$$

for which we have $[x] \diamond S \diamond R = [x] \diamond T \diamond R$, implying that we have

$$S^{-1}(x) = [x] \diamond S = [x] \diamond T = T^{-1}(x)$$

for each $x \in X$, implying $S = T$, and thus R is an epimorphism.

We can also prove this in a more abstract way, following [MSE 350788]:

- *Item 1* \implies *Item 2*: Assume that R is an epimorphism.
 - We first notice that the functor $\text{Rel}(-, \text{pt}) : \text{Rel}^{\text{op}} \rightarrow \text{Sets}$ maps R to R^{-1} by Remark 6.4.3.2.
 - Since $\text{Rel}(-, \text{pt})$ preserves limits by ?? of ??, it follows by ?? of ?? that $\text{Rel}(-, \text{pt})$ also preserves monomorphisms.
 - That is: $\text{Rel}(-, \text{pt})$ sends monomorphisms in Rel^{op} to monomorphisms in Sets.
 - The monomorphisms in Rel^{op} are precisely the epimorphisms in Rel by ?? of ??.
 - Since R is an epimorphism and $\text{Rel}(-, \text{pt})$ maps R to R^{-1} , it follows that R^{-1} is a monomorphism.
 - Since the monomorphisms in Sets are precisely the injections (?? of ??), it follows that R^{-1} is injective.
- *Item 2* \implies *Item 1*: Assume that R^{-1} is injective.

- We first notice that the functor $\text{Rel}(-, \text{pt}) : \text{Rel}^{\text{op}} \rightarrow \text{Sets}$ maps R to R^{-1} by [Remark 6.4.3.2](#).
- Since the monomorphisms in Sets are precisely the injections (?? of ??), it follows that R^{-1} is a monomorphism.
- Since $\text{Rel}(-, \text{pt})$ is faithful, it follows by ?? of ?? that $\text{Rel}(-, \text{pt})$ reflects monomorphisms.
- That is: $\text{Rel}(-, \text{pt})$ reflects monomorphisms in Sets to monomorphisms in Rel^{op} .
- The monomorphisms in Rel^{op} are precisely the epimorphisms in Rel by ?? of ??.
- Since R^{-1} is a monomorphism and $\text{Rel}(-, \text{pt})$ maps R to R^{-1} , it follows that R is an epimorphism.

We also claim that [Items 2](#) and [4](#) are equivalent, following [[MO 350788](#)]:


- [Item 2](#) \implies [Item 4](#): Since $B \setminus \{b\} \subset B$ and R^{-1} is injective, we have $R^{-1}(B \setminus \{b\}) \subseteq R^{-1}(B)$. So taking some $a \in R^{-1}(B) \setminus R^{-1}(B \setminus \{b\})$ we get an element of A such that $R(a) = \{b\}$.
- [Item 4](#) \implies [Item 2](#): Let $U, V \subset B$ with $U \neq V$. Without loss of generality, we can assume $U \setminus V \neq \emptyset$; otherwise just swap U and V . Let then $b \in U \setminus V$. By assumption, there exists an $a \in A$ with $R(a) = \{b\}$. Then $a \in R^{-1}(U)$ but $a \notin R^{-1}(V)$, and thus $R^{-1}(U) \neq R^{-1}(V)$, showing R^{-1} to be injective.

Finally, we prove the second part of the statement. So assume R is a total epimorphism in Rel and consider the diagram

$$A \xrightarrow{R} B \begin{array}{c} \xrightarrow{S} \\ \xrightarrow{T} \end{array} \{0, 1\},$$

where $b \sim_S 0$ for each $b \in B$ and where we have

$$b \sim_T \begin{cases} 0 & \text{if } b \in \text{Im}(R), \\ 1 & \text{otherwise} \end{cases}$$

for each $b \in B$. Since R is total, we have $a \sim_{S \circ R} 0$ and $a \sim_{T \circ R} 0$ for all $a \in A$, and no element of A is related to 1 by $S \circ R$ or $T \circ R$. Thus $S \circ R = T \circ R$, and since R is an epimorphism, we have $S = T$. But by the definition of T , this implies $\text{Im}(R) = B$. 

00M2 **5.3.10 2-Categorical Epimorphisms in Rel**

In this section we characterise (for now, some of) the 2-categorical epimorphisms in **Rel**, following [Section 9.2](#).

00M3 **PROPOSITION 5.3.10.1 ► 2-CATEGORICAL EPIMORPHISMS IN Rel**

Let $R: A \rightarrow B$ be a relation.

00M4 1. *Corepresentably Faithful Morphisms in Rel*. Every morphism of **Rel** is a corepresentably faithful morphism.

00M5 2. *Corepresentably Full Morphisms in Rel*. The following conditions are equivalent:

00M6 (a) The morphism $R: A \rightarrow B$ is a corepresentably full morphism.

00M7 (b) For each pair of relations $S, T: X \rightarrow A$, the following condition is satisfied:

(★) If $S \diamond R \subset T \diamond R$, then $S \subset T$.

00M8 (c) The functor

$$R^{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

is full.

00M9 (d) For each $U, V \in \mathcal{P}(B)$, if $R^{-1}(U) \subset R^{-1}(V)$, then $U \subset V$.

00MA (e) The functor

$$R_{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

is full.

00MB (f) For each $U, V \in \mathcal{P}(B)$, if $R_{-1}(U) \subset R_{-1}(V)$, then $U \subset V$.

00MC 3. *Corepresentably Fully Faithful Morphisms in Rel*. Every corepresentably full morphism of **Rel** is a corepresentably fully faithful morphism.

PROOF 5.3.10.2 ► PROOF OF PROPOSITION 5.3.10.1

Item 1: Corepresentably Faithful Morphisms in Rel

The relation R is a corepresentably faithful morphism in **Rel** iff, for each $X \in \text{Obj}(\mathbf{Rel})$, the functor

$$R^*: \mathbf{Rel}(B, X) \rightarrow \mathbf{Rel}(A, X)$$

is faithful, i.e. iff the morphism

$$R_{S,T}^* : \text{Hom}_{\mathbf{Rel}(B,X)}(S, T) \rightarrow \text{Hom}_{\mathbf{Rel}(A,X)}(S \diamond R, T \diamond R)$$

is injective for each $S, T \in \text{Obj}(\mathbf{Rel}(B, X))$. However, $\text{Hom}_{\mathbf{Rel}(B,X)}(S, T)$ is either empty or a singleton, in either case of which the map $R_{S,T}^*$ is necessarily injective.

Item 2: Corepresentably Full Morphisms in \mathbf{Rel}

We claim **Items 2a** to **2f** are indeed equivalent:

- **Item 2a** \iff **Item 2b**: This is simply a matter of unwinding definitions: The relation R is a corepresentably full morphism in \mathbf{Rel} iff, for each $X \in \text{Obj}(\mathbf{Rel})$, the functor

$$R^* : \mathbf{Rel}(B, X) \rightarrow \mathbf{Rel}(A, X)$$

is full, i.e. iff the morphism

$$R_{S,T}^* : \text{Hom}_{\mathbf{Rel}(B,X)}(S, T) \rightarrow \text{Hom}_{\mathbf{Rel}(A,X)}(S \diamond R, T \diamond R)$$

is surjective for each $S, T \in \text{Obj}(\mathbf{Rel}(B, X))$, i.e. iff, whenever $S \diamond R \subset T \diamond R$, we also have $S \subset T$.

- **Item 2c** \iff **Item 2d**: This is also simply a matter of unwinding definitions: The functor

$$R^{-1} : (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

is full iff, for each $U, V \in \mathcal{P}(A)$, the morphism

$$R_{U,V}^{-1} : \text{Hom}_{\mathcal{P}(B)}(U, V) \rightarrow \text{Hom}_{\mathcal{P}(A)}(R^{-1}(U), R^{-1}(V))$$

is surjective, i.e. iff whenever $R^{-1}(U) \subset R^{-1}(V)$, we also necessarily have $U \subset V$.

- **Item 2e** \iff **Item 2f**: This is once again simply a matter of unwinding definitions, and proceeds exactly in the same way as in the proof of the equivalence between **Items 2c** and **2d** given above.
- **Item 2d** \implies **Item 2f**: Suppose that the following condition is true:

$$(\star) \text{ For each } U, V \in \mathcal{P}(B), \text{ if } R^{-1}(U) \subset R^{-1}(V), \text{ then } U \subset V.$$

We need to show that the condition

(★) For each $U, V \in \mathcal{P}(B)$, if $R_{-1}(U) \subset R_{-1}(V)$, then $U \subset V$.

is also true. We proceed step by step:

1. Suppose we have $U, V \in \mathcal{P}(B)$ with $R_{-1}(U) \subset R_{-1}(V)$.
2. By **Item 7** of **Proposition 6.4.2.4**, we have

$$\begin{aligned} R_{-1}(U) &= B \setminus R^{-1}(A \setminus U), \\ R_{-1}(V) &= B \setminus R^{-1}(A \setminus V). \end{aligned}$$

3. By **Item 1** of **Proposition 2.3.10.2** we have $R^{-1}(A \setminus V) \subset R^{-1}(A \setminus U)$.
4. By assumption, we then have $A \setminus V \subset A \setminus U$.
5. By **Item 1** of **Proposition 2.3.10.2** again, we have $U \subset V$.

· **Item 2f** \implies **Item 2d**: Suppose that the following condition is true:

(★) For each $U, V \in \mathcal{P}(B)$, if $R_{-1}(U) \subset R_{-1}(V)$, then $U \subset V$.

We need to show that the condition

(★) For each $U, V \in \mathcal{P}(B)$, if $R^{-1}(U) \subset R^{-1}(V)$, then $U \subset V$.

is also true. We proceed step by step:

1. Suppose we have $U, V \in \mathcal{P}(B)$ with $R^{-1}(U) \subset R^{-1}(V)$.
2. By **Item 7** of **Proposition 6.4.3.4**, we have

$$\begin{aligned} R^{-1}(U) &= B \setminus R_{-1}(A \setminus U), \\ R^{-1}(V) &= B \setminus R_{-1}(A \setminus V). \end{aligned}$$

3. By **Item 1** of **Proposition 2.3.10.2** we have $R_{-1}(A \setminus V) \subset R_{-1}(A \setminus U)$.
4. By assumption, we then have $A \setminus V \subset A \setminus U$.
5. By **Item 1** of **Proposition 2.3.10.2** again, we have $U \subset V$.

· **Item 2b** \implies **Item 2d**: Consider the diagram

$$A \begin{array}{c} \xrightarrow{R} \\ \dashrightarrow \\ \xrightarrow{T} \end{array} B \begin{array}{c} \xrightarrow{S} \\ \dashrightarrow \\ \xrightarrow{T} \end{array} X,$$

and suppose that $S \diamond R \subset T \diamond R$. Note that, by assumption, given a diagram of the form

$$A \xrightarrow{R} B \begin{array}{c} \xrightarrow{U} \\ \xrightarrow{V} \end{array} \text{pt},$$

if $R^{-1}(U) = R \diamond U \subset R \diamond V = R^{-1}(V)$, then $U \subset V$. In particular, for each $x \in X$, we may consider the diagram

$$A \xrightarrow{R} B \begin{array}{c} \xrightarrow{S} \\ \xrightarrow{T} \end{array} X \xrightarrow{[x]} \text{pt},$$

for which we have $[x] \diamond S \diamond R \subset [x] \diamond T \diamond R$, implying that we have

$$S^{-1}(x) = [x] \diamond S \subset [x] \diamond T = T^{-1}(x)$$

for each $x \in X$, implying $S \subset T$.

- *Item 2d* \implies *Item 2b*: Let $U, V \in \mathcal{P}(B)$ and consider the diagram

$$A \xrightarrow{R} B \begin{array}{c} \xrightarrow{U} \\ \xrightarrow{V} \end{array} \text{pt}.$$

By [Remark 6.4.1.2](#), we have

$$R^{-1}(U) = U \diamond R,$$

$$R^{-1}(V) = V \diamond R.$$

Now, if $R^{-1}(U) \subset R^{-1}(V)$, i.e. $U \diamond R \subset V \diamond R$, then $U \subset V$ by assumption.

Item 3: Corepresentably Fully Faithful Morphisms in Rel

This follows from [Items 1](#) and [2](#). 

00MD

QUESTION 5.3.10.3 ► BETTER CHARACTERISATIONS OF COREPRESENTABLY FULL MORPHISMS IN Rel

[Item 2](#) of [Proposition 5.3.10.1](#) gives a characterisation of the corepresentably full morphisms in **Rel**.

Are there other nice characterisations of these?

This question also appears as [\[MO 467527\]](#).

00ME 5.3.11 Co/Limits in Rel

00MF PROPOSITION 5.3.11.1 ► Co/LIMITS IN Rel

This will be properly written later on.

PROOF 5.3.11.2 ► PROOF OF PROPOSITION 5.3.11.1

Omitted. 

00MG 5.3.12 Kan Extensions and Kan Lifts in Rel

00MH REMARK 5.3.12.1 ► KAN EXTENSIONS AND KAN LIFTS IN Rel

The 2-category **Rel** admits all right Kan extensions and right Kan lifts, though not all left Kan extensions and neither does it admit all left Kan lifts. See [Section 6.2](#) for a detailed discussion of this.

00MJ 5.3.13 Closedness of Rel

00MK PROPOSITION 5.3.13.1 ► CLOSEDNESS OF Rel

The 2-category **Rel** is a closed bicategory, there being, for each $R: A \rightarrow B$ and set X , a pair of adjunctions

$$\begin{aligned} (R^* \dashv \text{Ran}_R) : \text{Rel}(B, X) &\begin{array}{c} \xrightarrow{R^*} \\ \perp \\ \xleftarrow{\text{Ran}_R} \end{array} \text{Rel}(A, X), \\ (R_* \dashv \text{Rift}_R) : \text{Rel}(X, A) &\begin{array}{c} \xrightarrow{R_*} \\ \perp \\ \xleftarrow{\text{Rift}_R} \end{array} \text{Rel}(X, B), \end{aligned}$$

witnessed by bijections

$$\begin{aligned} \mathbf{Rel}(S \diamond R, T) &\cong \mathbf{Rel}(S, \text{Ran}_R(T)), \\ \mathbf{Rel}(R \diamond U, V) &\cong \mathbf{Rel}(U, \text{Rift}_R(V)), \end{aligned}$$

natural in $S \in \text{Rel}(B, X)$, $T \in \text{Rel}(A, X)$, $U \in \text{Rel}(X, A)$, and $V \in \text{Rel}(X, B)$.

PROOF 5.3.13.2 ► PROOF OF PROPOSITION 5.3.13.1

This follows from [Propositions 6.2.3.1](#) and [6.2.4.1](#). 

00ML 5.3.14 Rel as a Category of Free Algebras

00MM PROPOSITION 5.3.14.1 ► **Rel AS A CATEGORY OF FREE ALGEBRAS**

We have an isomorphism of categories

$$\mathbf{Rel} \cong \mathbf{FreeAlg}_{\mathcal{P}_*}(\mathbf{Sets}),$$

where \mathcal{P}_* is the powerset monad of ??.

PROOF 5.3.14.2 ► PROOF OF PROPOSITION 5.3.14.1

Omitted. 

00MN 5.4 The Left Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

00MP 5.4.1 The Left Skew Monoidal Product

Let A and B be sets and let $J: A \rightarrow B$ be a relation.

00MQ DEFINITION 5.4.1.1 ► THE LEFT J -SKEW MONOIDAL PRODUCT OF $\mathbf{REL}(A, B)$

The **left J -skew monoidal product of $\mathbf{Rel}(A, B)$** is the functor

$$\triangleleft_J: \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

where

- *Action on Objects.* For each $R, S \in \mathbf{Obj}(\mathbf{Rel}(A, B))$, we have

$$S \triangleleft_J R \stackrel{\text{def}}{=} S \diamond \mathbf{Rift}_J(R),$$

- *Action on Morphisms.* For each $R, S, R', S' \in \mathbf{Obj}(\mathbf{Rel}(A, B))$, the action on Hom-sets

$$(\triangleleft_J)_{(G,F),(G',F')} : \mathbf{Hom}_{\mathbf{Rel}(A,B)}(S, S') \times \mathbf{Hom}_{\mathbf{Rel}(A,B)}(R, R') \rightarrow \mathbf{Hom}_{\mathbf{Rel}(A,B)}(S \triangleleft_J R, S' \triangleleft_J R')$$

of \triangleleft_J at $((R, S), (R', S'))$ is defined by¹

$$\beta \triangleleft_J \alpha \stackrel{\text{def}}{=} \beta \diamond \text{Rift}_J(\alpha),$$

for each $\beta \in \text{Hom}_{\mathbf{Rel}(A,B)}(S, S')$ and each $\alpha \in \text{Hom}_{\mathbf{Rel}(A,B)}(R, R')$.

¹Since $\mathbf{Rel}(A, B)$ is posetal, this is to say that if $S \subset S'$ and $R \subset R'$, then $S \triangleleft_J R \subset S' \triangleleft_J R'$.

00MR 5.4.2 The Left Skew Monoidal Unit

Let A and B be sets and let $J: A \rightarrow B$ be a relation.

00MS DEFINITION 5.4.2.1 ▶ THE LEFT J -SKEW MONOIDAL UNIT OF $\mathbf{REL}(A, B)$

The **left J -skew monoidal unit** of $\mathbf{Rel}(A, B)$ is the functor

$$\mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A,B)} : \text{pt} \rightarrow \mathbf{Rel}(A, B)$$

picking the object

$$\mathbb{1}_{\mathbf{Rel}(A,B)}^{\triangleleft_J} \stackrel{\text{def}}{=} J$$

of $\mathbf{Rel}(A, B)$.

00MT 5.4.3 The Left Skew Associators


Let A and B be sets and let $J: A \rightarrow B$ be a relation.

00MU DEFINITION 5.4.3.1 ▶ THE LEFT J -SKEW ASSOCIATOR OF $\mathbf{REL}(A, B)$

The **left J -skew associator** of $\mathbf{Rel}(A, B)$ is the natural transformation

$$\alpha^{\mathbf{Rel}(A,B), \triangleleft_J} : \triangleleft_J \circ (\triangleleft_J \times \text{id}) \Longrightarrow \triangleleft_J \circ (\text{id} \times \triangleleft_J) \circ \alpha_{\mathbf{Rel}(A,B), \mathbf{Rel}(A,B), \mathbf{Rel}(A,B)}^{\text{Cats}}$$

as in the diagram



whose component

$$\alpha_{T, S, R}^{\mathbf{Rel}(A, B), \triangleleft_J} : \underbrace{(T \triangleleft_J S) \triangleleft_J R}_{\stackrel{\text{def}}{=} T \diamond \text{Rift}_J(S) \diamond \text{Rift}_J(R)} \hookrightarrow \underbrace{T \triangleleft_J (S \triangleleft_J R)}_{\stackrel{\text{def}}{=} T \diamond \text{Rift}_J(S \diamond \text{Rift}_J(R))}$$

at (T, S, R) is given by

$$\alpha_{T, S, R}^{\mathbf{Rel}(A, B), \triangleleft_J} \stackrel{\text{def}}{=} \text{id}_T \diamond \gamma,$$

where

$$\gamma : \text{Rift}_J(S) \diamond \text{Rift}_J(R) \hookrightarrow \text{Rift}_J(S \diamond \text{Rift}_J(R))$$

is the inclusion adjunct to the inclusion

$$\epsilon_S \star \text{id}_{\text{Rift}_J(R)} : \underbrace{J \diamond \text{Rift}_J(S) \diamond \text{Rift}_J(R)}_{\stackrel{\text{def}}{=} J_* (\text{Rift}_J(S) \diamond \text{Rift}_J(R))} \hookrightarrow S \diamond \text{Rift}_J(R)$$

under the adjunction $J_* \dashv \text{Rift}_J$, where $\epsilon : J \diamond \text{Rift}_J \Longrightarrow \text{id}_{\mathbf{Rel}(A, B)}$ is the counit of the adjunction $J_* \dashv \text{Rift}_J$.

00MV 5.4.4 The Left Skew Left Unitors

Let A and B be sets and let $J : A \rightarrow B$ be a relation.


00MW

DEFINITION 5.4.4.1 ▶ THE LEFT J -SKEW LEFT UNITOR OF $\mathbf{REL}(A, B)$

The **left J -skew left unitor** of $\mathbf{Rel}(A, B)$ is the natural transformation

$$\lambda^{\mathbf{Rel}(A,B), \triangleleft_J} : \triangleleft_J \circ \left(\mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A,B)} \times \text{id} \right) \Longrightarrow \lambda_{\mathbf{Rel}(A,B)}^{\text{Cats}_2}$$

as in the diagram



whose component

$$\lambda_R^{\mathbf{Rel}(A,B), \triangleleft_J} : J \triangleleft_J R \hookrightarrow R$$

$\underbrace{\hspace{2cm}}_{\stackrel{\text{def}}{=} J \circ \text{Rift}_J(R)}$

at R is given by

$$\lambda_R^{\mathbf{Rel}(A,B), \triangleleft_J} \stackrel{\text{def}}{=} \epsilon_R,$$

where $\epsilon : J_* \circ \text{Rift}_J \Longrightarrow \text{id}_{\mathbf{Rel}(A,B)}$ is the counit of the adjunction $J_* \dashv \text{Rift}_J$.

00MX 5.4.5 The Left Skew Right Unitors

Let A and B be sets and let $J : A \rightarrow B$ be a relation.


00MY

DEFINITION 5.4.5.1 ▶ THE LEFT J -SKEW RIGHT UNITOR OF $\mathbf{REL}(A, B)$

The **left J -skew right unitor** of $\mathbf{Rel}(A, B)$ is the natural transformation

$$\rho^{\mathbf{Rel}(A,B), \triangleleft_J} : \rho_{\mathbf{Rel}(A,B)}^{\text{Cats}_2} \Longrightarrow \triangleleft_J \circ \left(\text{id} \times \mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A,B)} \right)$$

as in the diagram



whose component

$$\rho_R^{\mathbf{Rel}(A, B), \triangleleft_J} : R \hookrightarrow R \triangleleft_J J$$

$\underbrace{\hspace{10em}}_{\stackrel{\text{def}}{=} R \diamond \text{Rift}_J(J)}$

at R is given by the composition

$$\begin{aligned} R &\xrightarrow{\sim} R \diamond \chi_A \\ &\xrightarrow{\text{id}_R \diamond \eta_{\chi_A}} R \diamond \text{Rift}_J(J_*(\chi_A)) \\ &\stackrel{\text{def}}{=} R \diamond \text{Rift}_J(J \diamond \chi_A) \\ &\xrightarrow{\sim} R \diamond \text{Rift}_J(J) \\ &\stackrel{\text{def}}{=} R \triangleleft_J J, \end{aligned}$$

where $\eta : \text{id}_{\mathbf{Rel}(A, A)} \implies \text{Rift}_J \circ J_*$ is the unit of the adjunction $J_* \dashv \text{Rift}_J$.

00MZ 5.4.6 The Left Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

00N0 PROPOSITION 5.4.6.1 ► THE LEFT J -SKEW MONOIDAL STRUCTURE ON $\mathbf{REL}(A, B)$

The category $\mathbf{Rel}(A, B)$ admits a left skew monoidal category structure consisting of

- *The Underlying Category.* The posetal category associated to the poset $\mathbf{Rel}(A, B)$ of relations from A to B of **Item 2** of **Definition 5.1.1.3**.
- *The Left Skew Monoidal Product.* The left J -skew monoidal product

$$\triangleleft_J : \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

of Definition 5.4.1.1.

- *The Left Skew Monoidal Unit.* The functor

$$\mathbb{1}^{\mathbf{Rel}(A,B), \triangleleft_J} : \mathbf{pt} \rightarrow \mathbf{Rel}(A, B)$$

of Definition 5.4.2.1.

- *The Left Skew Associators.* The natural transformation

$$\alpha^{\mathbf{Rel}(A,B), \triangleleft_J} : \triangleleft_J \circ (\triangleleft_J \times \mathrm{id}) \Longrightarrow \triangleleft_J \circ (\mathrm{id} \times \triangleleft_J) \circ \alpha_{\mathbf{Rel}(A,B), \mathbf{Rel}(A,B), \mathbf{Rel}(A,B)}^{\mathbf{Cats}}$$

of Definition 5.4.3.1.

- *The Left Skew Left Unitors.* The natural transformation

$$\lambda^{\mathbf{Rel}(A,B), \triangleleft_J} : \triangleleft_J \circ \left(\mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A,B)} \times \mathrm{id} \right) \Longrightarrow \lambda_{\mathbf{Rel}(A,B)}^{\mathbf{Cats}_2}$$


of Definition 5.4.4.1.

- *The Left Skew Right Unitors.* The natural transformation

$$\rho^{\mathbf{Rel}(A,B), \triangleleft_J} : \rho_{\mathbf{Rel}(A,B)}^{\mathbf{Cats}_2} \Longrightarrow \triangleleft_J \circ \left(\mathrm{id} \times \mathbb{1}_{\triangleleft_J}^{\mathbf{Rel}(A,B)} \right)$$

of Definition 5.4.5.1.

PROOF 5.4.6.2 ► PROOF OF PROPOSITION 5.4.6.1

Since $\mathbf{Rel}(A, B)$ is posetal, the commutativity of the pentagon identity, the left skew left triangle identity, the left skew right triangle identity, the left skew middle triangle identity, and the zigzag identity is automatic, and thus $\mathbf{Rel}(A, B)$ together with the data in the statement forms a left skew monoidal category. 

00N1 5.5 The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

Let A and B be sets and let $J : A \rightarrow B$ be a relation.

00N2 5.5.1 The Right Skew Monoidal Product

00N3

DEFINITION 5.5.1.1 ▶ THE RIGHT J -SKEW MONOIDAL PRODUCT OF $\mathbf{REL}(A, B)$

The **right J -skew monoidal product of $\mathbf{Rel}(A, B)$** is the functor

$$\triangleright_J : \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

where

- *Action on Objects.* For each $R, S \in \text{Obj}(\mathbf{Rel}(A, B))$, we have

$$S \triangleright_J R \stackrel{\text{def}}{=} \text{Ran}_J(S) \diamond R,$$

- *Action on Morphisms.* For each $R, S, R', S' \in \text{Obj}(\mathbf{Rel}(A, B))$, the action on Hom-sets

$$(\triangleright_J)_{(S,R),(S',R')} : \text{Hom}_{\mathbf{Rel}(A,B)}(S, S') \times \text{Hom}_{\mathbf{Rel}(A,B)}(R, R') \rightarrow \text{Hom}_{\mathbf{Rel}(A,B)}(S \triangleright_J R, S' \triangleright_J R')$$

of \triangleright_J at $((S, R), (S', R'))$ is defined by¹

$$\beta \triangleright_J \alpha \stackrel{\text{def}}{=} \text{Ran}_J(\beta) \diamond \alpha,$$

for each $\beta \in \text{Hom}_{\mathbf{Rel}(A,B)}(S, S')$ and each $\alpha \in \text{Hom}_{\mathbf{Rel}(A,B)}(R, R')$.

¹Since $\mathbf{Rel}(A, B)$ is posetal, this is to say that if $S \subset S'$ and $R \subset R'$, then $S \triangleright_J R \subset S' \triangleright_J R'$.

00N4 5.5.2 The Right Skew Monoidal Unit

00N5

DEFINITION 5.5.2.1 ▶ THE RIGHT J -SKEW MONOIDAL UNIT OF $\mathbf{REL}(A, B)$

The **right J -skew monoidal unit of $\mathbf{Rel}(A, B)$** is the functor

$$\mathbb{1}_{\triangleright_J}^{\mathbf{Rel}(A,B)} : \text{pt} \rightarrow \mathbf{Rel}(A, B)$$

picking the object

$$\mathbb{1}_{\mathbf{Rel}(A,B)}^{\triangleright J} \stackrel{\text{def}}{=} J$$

of $\mathbf{Rel}(A, B)$.


00N6 **5.5.3 The Right Skew Associators**

00N7 **DEFINITION 5.5.3.1 ▶ THE RIGHT J -SKEW ASSOCIATOR OF $\mathbf{Rel}(A, B)$**

The **right J -skew associator** of $\mathbf{Rel}(A, B)$ is the natural transformation

$$\alpha^{\mathbf{Rel}(A,B),\triangleright J} : \triangleright_J \circ (\text{id} \times \triangleright_J) \implies \triangleright_J \circ (\triangleright_J \times \text{id}) \circ \alpha_{\mathbf{Rel}(A,B),\mathbf{Rel}(A,B),\mathbf{Rel}(A,B)}^{\text{Cats},-1}$$

as in the diagram



whose component

$$\alpha_{T,S,R}^{\mathbf{Rel}(A,B),\triangleright J} : \underbrace{T \triangleright_J (S \triangleright_J R)}_{\stackrel{\text{def}}{=} \text{Ran}_J(T) \diamond \text{Ran}_J(S) \diamond R} \hookrightarrow \underbrace{(T \triangleright_J S) \triangleright_J R}_{\stackrel{\text{def}}{=} \text{Ran}_J(\text{Ran}_J(T) \diamond S) \diamond R}$$

at (T, S, R) is given by

$$\alpha_{T,S,R}^{\mathbf{Rel}(A,B),\triangleright J} \stackrel{\text{def}}{=} \gamma \diamond \text{id}_R,$$

where

$$\gamma : \text{Ran}_J(T) \diamond \text{Ran}_J(S) \hookrightarrow \text{Ran}_J(\text{Ran}_J(T) \diamond S)$$

is the inclusion adjunct to the inclusion

$$\text{id}_{\text{Ran}_J(T)} \diamond \epsilon_S : \underbrace{\text{Ran}_J(T) \diamond \text{Ran}_J(S) \diamond J}_{\stackrel{\text{def}}{=} J^*(\text{Ran}_J(T) \diamond \text{Ran}_J(S))} \hookrightarrow \text{Ran}_J(T) \diamond S$$

under the adjunction $J^* \dashv \text{Ran}_J$, where $\epsilon : \text{Ran}_J \diamond J \implies \text{id}_{\mathbf{Rel}(A,B)}$ is the counit of the adjunction $J^* \dashv \text{Ran}_J$.


00N8 **5.5.4 The Right Skew Left Unitors**

00N9 **DEFINITION 5.5.4.1 ▶ THE RIGHT J -SKEW LEFT UNITOR OF $\mathbf{REL}(A, B)$**

The **right J -skew left unitor** of $\mathbf{Rel}(A, B)$ is the natural transformation

$$\lambda^{\mathbf{Rel}(A,B), \triangleright J} : \lambda_{\mathbf{Rel}(A,B)}^{\mathbf{Cats}_2} \Longrightarrow \triangleright J \circ \left(\mathbb{1}_{\triangleright}^{\mathbf{Rel}(A,B)} \times \text{id} \right),$$

as in the diagram



whose component

$$\lambda_R^{\mathbf{Rel}(A,B), \triangleright J} : R \hookrightarrow \underbrace{J \triangleright_J R}_{\stackrel{\text{def}}{=} \text{Ran}_J(J) \diamond R}$$

at R is given by the composition

$$\begin{aligned} R &\xrightarrow{\sim} \chi_B \diamond R \\ &\xrightarrow{\eta_{\chi_B}} \diamond \text{id}_{\text{Ran}_J(J^*(\chi_A))} \diamond R \\ &\stackrel{\text{def}}{=} \text{Ran}_J(J^* \diamond \chi_A) \diamond R \\ &\xrightarrow{\sim} \text{Ran}_J(J) \diamond R \\ &\stackrel{\text{def}}{=} R \triangleright_J J, \end{aligned}$$

where $\eta : \text{id}_{\mathbf{Rel}(B,B)} \Longrightarrow \text{Ran}_J \circ J^*$ is the unit of the adjunction $J^* \dashv \text{Ran}_J$.

00NA **5.5.5 The Right Skew Right Unitors**


00NB

DEFINITION 5.5.5.1 ▶ THE RIGHT J -SKEW RIGHT UNITOR OF $\mathbf{REL}(A, B)$

The **right J -skew right unitor** of $\mathbf{Rel}(A, B)$ is the natural transformation

$$\rho^{\mathbf{Rel}(A,B), \triangleright J} : \triangleright_J \circ (\text{id} \times \mathbb{1}_{\triangleright}^{\mathbf{Rel}(A,B)}) \implies \rho_{\mathbf{Rel}(A,B)}^{\text{Cats}_2},$$

as in the diagram



whose component

$$\rho_S^{\mathbf{Rel}(A,B), \triangleright J} : S \triangleright_J J \hookrightarrow S$$

$\underbrace{\hspace{1.5cm}}_{\stackrel{\text{def}}{=} \text{Ran}_J(S) \circ J}$

at S is given by

$$\rho_S^{\mathbf{Rel}(A,B), \triangleright J} \stackrel{\text{def}}{=} \epsilon_R,$$

where $\epsilon : J^* \circ \text{Ran}_J \implies \text{id}_{\mathbf{Rel}(A,B)}$ is the counit of the adjunction $J^* \dashv \text{Ran}_J$.

00NC

5.5.6 The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

00ND

PROPOSITION 5.5.6.1 ▶ THE RIGHT J -SKEW MONOIDAL STRUCTURE ON $\mathbf{REL}(A, B)$

The category $\mathbf{Rel}(A, B)$ admits a right skew monoidal category structure consisting of

- *The Underlying Category.* The posetal category associated to the poset $\mathbf{Rel}(A, B)$ of relations from A to B of **Item 2** of **Definition 5.1.1.3**.
- *The Right Skew Monoidal Product.* The right J -skew monoidal product

$$\triangleleft_J : \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

of **Definition 5.5.1.1**.

- *The Right Skew Monoidal Unit.* The functor

$$\mathbb{1}^{\mathbf{Rel}(A,B),\triangleleft_J} : \mathbf{pt} \rightarrow \mathbf{Rel}(A, B)$$

of [Definition 5.5.2.1](#).

- *The Right Skew Associators.* The natural transformation

$$\alpha^{\mathbf{Rel}(A,B),\triangleright_J} : \triangleright_J \circ (\mathrm{id} \times \triangleright_J) \Longrightarrow \triangleright_J \circ (\triangleright_J \times \mathrm{id}) \circ \alpha_{\mathbf{Rel}(A,B),\mathbf{Rel}(A,B),\mathbf{Rel}(A,B)}^{\mathbf{Cats},-1}$$

of [Definition 5.5.3.1](#).

- *The Right Skew Left Unitors.* The natural transformation

$$\lambda^{\mathbf{Rel}(A,B),\triangleright_J} : \lambda_{\mathbf{Rel}(A,B)}^{\mathbf{Cats}_2} \Longrightarrow \triangleright_J \circ \left(\mathbb{1}_{\triangleright}^{\mathbf{Rel}(A,B)} \times \mathrm{id} \right)$$


of [Definition 5.5.4.1](#).

- *The Right Skew Right Unitors.* The natural transformation

$$\rho^{\mathbf{Rel}(A,B),\triangleright_J} : \triangleright_J \circ \left(\mathrm{id} \times \mathbb{1}_{\triangleright}^{\mathbf{Rel}(A,B)} \right) \Longrightarrow \rho_{\mathbf{Rel}(A,B)}^{\mathbf{Cats}_2}$$

of [Definition 5.5.5.1](#).

PROOF 5.5.6.2 ► PROOF OF PROPOSITION 5.5.6.1

Since $\mathbf{Rel}(A, B)$ is posetal, the commutativity of the pentagon identity, the right skew left triangle identity, the right skew right triangle identity, the right skew middle triangle identity, and the zigzag identity is automatic, and thus $\mathbf{Rel}(A, B)$ together with the data in the statement forms a right skew monoidal category. 

Appendices

5.A Other Chapters

Sets

1. [Sets](#)
2. [Constructions With Sets](#)
3. [Pointed Sets](#)
4. [Tensor Products of Pointed Sets](#)

Relations

5. [Relations](#)
6. [Constructions With Relations](#)
7. [Equivalence Relations and Apartness Relations](#)

Category Theory

8. Categories

Bicategories

9. Types of Morphisms in Bicategories

Chapter 6

Constructions With Relations

00NE This chapter contains some material about constructions with relations. Notably, we discuss and explore:

1. The existence or non-existence of Kan extensions and Kan lifts in the 2-category **Rel** (Section 6.2).
2. The various kinds of constructions involving relations, such as graphs, domains, ranges, unions, intersections, products, inverse relations, composition of relations, and collages (Section 6.3).
3. The adjoint pairs

$$\begin{aligned}R_* \dashv R_{-1} &: \mathcal{P}(A) \rightleftarrows \mathcal{P}(B), \\ R^{-1} \dashv R_! &: \mathcal{P}(B) \rightleftarrows \mathcal{P}(A)\end{aligned}$$

of functors (morphisms of posets) between $\mathcal{P}(A)$ and $\mathcal{P}(B)$ induced by a relation $R: A \rightarrow B$, as well as the properties of R_* , R_{-1} , R^{-1} , and $R_!$ (Section 6.4).

Of particular note are the following points:

- (a) These two pairs of adjoint functors are the counterpart for relations of the adjoint triple $f_* \dashv f^{-1} \dashv f_!$ induced by a function $f: A \rightarrow B$ studied in Section 2.4.
- (b) We have $R_{-1} = R^{-1}$ iff R is total and functional (Item 8 of Proposition 6.4.2.4).
- (c) As a consequence of the previous item, when R comes from a function f , the pair of adjunctions

$$R_* \dashv R_{-1} = R^{-1} \dashv R_!$$

reduces to the triple adjunction

$$f_* \dashv f^{-1} \dashv f_!$$

from [Section 2.4](#).

- (d) The pairs $R_* \dashv R_{-1}$ and $R^{-1} \dashv R!$ turn out to be rather important later on, as they appear in the definition and study of continuous, open, and closed relations between topological spaces (??).

Contents

6.1	Co/Limits in the Category of Relations	336
6.2	Kan Extensions and Kan Lifts in the 2-Category of Relations	337
6.2.1	Left Kan Extensions in Rel	337
6.2.2	Left Kan Lifts in Rel	338
6.2.3	Right Kan Extensions in Rel	339
6.2.4	Right Kan Lifts in Rel	340
6.3	More Constructions With Relations	342
6.3.1	The Graph of a Function	342
6.3.2	The Inverse of a Function	346
6.3.3	Representable Relations	348
6.3.4	The Domain and Range of a Relation	348
6.3.5	Binary Unions of Relations	349
6.3.6	Unions of Families of Relations	351
6.3.7	Binary Intersections of Relations	352
6.3.8	Intersections of Families of Relations	353
6.3.9	Binary Products of Relations	354
6.3.10	Products of Families of Relations	356
6.3.11	The Inverse of a Relation	357
6.3.12	Composition of Relations	359
6.3.13	The Collage of a Relation	364
6.4	Functoriality of Powersets	366
6.4.1	Direct Images	366
6.4.2	Strong Inverse Images	372
6.4.3	Weak Inverse Images	378
6.4.4	Direct Images With Compact Support	384
6.4.5	Functoriality of Powersets	391
6.4.6	Functoriality of Powersets: Relations on Powersets ..	392
6.A	Other Chapters	394

00NF 6.1 Co/Limits in the Category of Relations

This section is currently just a stub, and will be properly developed later on.

6.2 Kan Extensions and Kan Lifts in the 2-Category of Relations

00NG

6.2.1 Left Kan Extensions in Rel

00NJ

PROPOSITION 6.2.1.1 ► LEFT KAN EXTENSIONS IN Rel

Let $R: A \rightarrow B$ be a relation.

00NK

1. *Non-Existence of All Left Kan Extensions in Rel.* Not all relations in **Rel** admit left Kan extensions.

00NL

2. *Characterisation of Relations Admitting Left Kan Extensions Along Them.*

The following conditions are equivalent:

(a) The left Kan extension

$$\text{Lan}_R: \mathbf{Rel}(A, X) \rightarrow \mathbf{Rel}(B, X)$$

along R exists.

(b) The relation R admits a left adjoint in **Rel**.

(c) The relation R is of the form f^{-1} (as in [Definition 6.3.2.1](#)) for some function f .

PROOF 6.2.1.2 ► PROOF OF PROPOSITION 6.2.1.1

Item 1: Non-Existence of All Left Kan Extensions in Rel

Omitted, but will eventually follow [Fosco Loregian's comment](#) on [MO 460656].

Item 2: Characterisation of Relations Admitting Left Kan Extensions Along Them

Omitted, but will eventually follow [Tim Champion's answer](#) to [MO 460656].



00NM

QUESTION 6.2.1.3 ► EXISTENCE OF SPECIFIC LEFT KAN EXTENSIONS OF RELATIONS

Given relations $S: A \rightarrow X$ and $R: A \rightarrow B$, is there a characterisation of when the left Kan extension

$$\text{Lan}_S(R): B \rightarrow X$$

exists in terms of properties of R and S ?

This question also appears as [MO 461592].

00NN

QUESTION 6.2.1.4 ► EXPLICIT DESCRIPTION OF LEFT KAN EXTENSIONS ALONG FUNCTIONS

As shown in [Item 2](#) of [Proposition 6.2.1.1](#), the left Kan extension

$$\text{Lan}_R: \mathbf{Rel}(A, X) \rightarrow \mathbf{Rel}(B, X)$$

along a relation of the form $R = f^{-1}$ exists. Is there an explicit description of it, similarly to the explicit description of right Kan extensions given in [Proposition 6.2.3.1](#)?

This question also appears as [\[MO 461592\]](#).

00NP 6.2.2 Left Kan Lifts in Rel

00NQ

PROPOSITION 6.2.2.1 ► LEFT KAN LIFTS IN Rel

Let $R: A \rightarrow B$ be a relation.

00NR

1. *Non-Existence of All Left Kan Lifts in Rel.* Not all relations in **Rel** admit left Kan lifts.

00NS

2. *Characterisation of Relations Admitting Left Kan Lifts Along Them.* The following conditions are equivalent:

- (a) The left Kan lift

$$\text{Lift}_R: \mathbf{Rel}(X, B) \rightarrow \mathbf{Rel}(X, A)$$

along R exists.

- (b) The relation R admits a right adjoint in **Rel**.
- (c) The relation R is of the form $\text{Gr}(f)$ (as in [Definition 6.3.1.1](#)) for some function f .

PROOF 6.2.2.2 ► PROOF OF PROPOSITION 6.2.2.1

Item 1: Non-Existence of All Left Kan Lifts in Rel

Omitted, but will eventually follow (the dual of) [Fosco Loregian's comment](#) on [\[MO 460656\]](#).

Item 2: Characterisation of Relations Admitting Left Kan Lifts Along Them

Omitted, but will eventually follow [Tim Champion's answer](#) to [\[MO 460656\]](#).



00NT

QUESTION 6.2.2.3 ► EXISTENCE OF SPECIFIC LEFT KAN LIFTS OF RELATIONS

Given relations $S: A \rightarrow X$ and $R: A \rightarrow B$, is there a characterisation of when the left Kan lift

$$\text{Lift}_S(R): X \rightarrow A$$

exists in terms of properties of R and S ?

This question also appears as [MO 461592].

00NU

QUESTION 6.2.2.4 ► EXPLICIT DESCRIPTION OF LEFT KAN LIFTS ALONG FUNCTIONS

As shown in Item 2 of Proposition 6.2.2.1, the left Kan lift

$$\text{Lift}_R: \mathbf{Rel}(X, B) \rightarrow \mathbf{Rel}(X, A)$$

along a relation of the form $R = \text{Gr}(f)$ exists. Is there an explicit description of it, similarly to the explicit description of right Kan lifts given in Proposition 6.2.4.1?

This question also appears as [MO 461592].

00NV 6.2.3 Right Kan Extensions in Rel

Let $R: A \rightarrow B$ be a relation.

00NW

PROPOSITION 6.2.3.1 ► EXISTENCE OF RIGHT KAN EXTENSIONS IN Rel

The right Kan extension

$$\text{Ran}_R: \mathbf{Rel}(A, X) \rightarrow \mathbf{Rel}(B, X)$$

along R in \mathbf{Rel} exists and is given by

$$\text{Ran}_R(S) \stackrel{\text{def}}{=} \int_{a \in A} \mathbf{Hom}_{\{t, f\}}(R_a^{-2}, S_a^{-1})$$

for each $S \in \mathbf{Rel}(A, X)$, so that the following conditions are equivalent:

1. We have $b \sim_{\text{Ran}_R(S)} x$.
2. For each $a \in A$, if $a \sim_R b$, then $a \sim_S x$.

PROOF 6.2.3.2 ► PROOF OF PROPOSITION 6.2.3.1

We have


$$\begin{aligned}
\mathrm{Hom}_{\mathbf{Rel}(A,X)}(S \diamond R, T) &\cong \int_{a \in A} \int_{x \in X} \mathbf{Hom}_{\{t,f\}}((S \diamond R)_a^x, T_a^x) \\
&\cong \int_{a \in A} \int_{x \in X} \mathbf{Hom}_{\{t,f\}}\left(\left(\int_{b \in B} S_b^x \times R_a^b\right), T_a^x\right) \\
&\cong \int_{a \in A} \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{t,f\}}(S_b^x \times R_a^b, T_a^x) \\
&\cong \int_{a \in A} \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{t,f\}}(S_b^x, \mathbf{Hom}_{\{t,f\}}(R_a^b, T_a^x)) \\
&\cong \int_{b \in B} \int_{x \in X} \int_{a \in A} \mathbf{Hom}_{\{t,f\}}(S_b^x, \mathbf{Hom}_{\{t,f\}}(R_a^b, T_a^x)) \\
&\cong \int_{b \in B} \int_{x \in X} \mathbf{Hom}_{\{t,f\}}\left(S_b^x, \int_{a \in A} \mathbf{Hom}_{\{t,f\}}(R_a^b, T_a^x)\right) \\
&\cong \mathrm{Hom}_{\mathbf{Rel}(B,X)}\left(S, \int_{a \in A} \mathbf{Hom}_{\{t,f\}}(R_a^{-2}, T_a^{-1})\right)
\end{aligned}$$

naturally in each $S \in \mathbf{Rel}(B, X)$ and each $T \in \mathbf{Rel}(A, X)$, showing that

$$\int_{a \in A} \mathbf{Hom}_{\{t,f\}}(R_a^{-2}, T_a^{-1})$$

is right adjoint to the precomposition functor $- \diamond R$, being thus the right Kan extension along R . Here we have used the following results, respectively (i.e. for each \cong sign):

1. **Item 1** of **Proposition 5.1.1.6**.
2. **Definition 6.3.12.1**.
3. ?? of ??.
4. **Proposition 1.2.2.5**.
5. ?? of ??.
6. ?? of ??.
7. **Item 1** of **Proposition 5.1.1.6**.

This finishes the proof. 

00NX 6.2.4 Right Kan Lifts in Rel

Let $R: A \rightarrow B$ be a relation.

00NY

PROPOSITION 6.2.4.1 ► EXISTENCE OF RIGHT KAN LIFTS IN Rel

The right Kan lift

$$\text{Rift}_R: \text{Rel}(X, B) \rightarrow \text{Rel}(X, A)$$

along R in **Rel** exists and is given by

$$\text{Rift}_R(S) \stackrel{\text{def}}{=} \int_{b \in B} \mathbf{Hom}_{\{t,f\}}(R_{-1}^b, S_{-2}^b)$$

for each $S \in \text{Rel}(X, B)$, so that the following conditions are equivalent:

1. We have $x \sim_{\text{Rift}_R(S)} a$.
2. For each $b \in B$, if $a \sim_R b$, then $x \sim_S b$.

PROOF 6.2.4.2 ► PROOF OF PROPOSITION 6.2.4.1

We have

$$\begin{aligned} \text{Hom}_{\mathbf{Rel}(X,B)}(R \diamond S, T) &\cong \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{t,f\}}((R \diamond S)_x^b, T_x^b) \\ &\cong \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{t,f\}}\left(\left(\int^{a \in A} R_a^b \times S_x^a\right), T_x^b\right) \\ &\cong \int_{x \in X} \int_{b \in B} \int_{a \in A} \mathbf{Hom}_{\{t,f\}}(R_a^b \times S_x^a, T_x^b) \\ &\cong \int_{x \in X} \int_{b \in B} \int_{a \in A} \mathbf{Hom}_{\{t,f\}}(S_x^a, \mathbf{Hom}_{\{t,f\}}(R_a^b, T_x^b)) \\ &\cong \int_{x \in X} \int_{a \in A} \int_{b \in B} \mathbf{Hom}_{\{t,f\}}(S_x^a, \mathbf{Hom}_{\{t,f\}}(R_a^b, T_x^b)) \\ &\cong \int_{x \in X} \int_{a \in A} \mathbf{Hom}_{\{t,f\}}\left(S_x^a, \int_{b \in B} \mathbf{Hom}_{\{t,f\}}(R_a^b, T_x^b)\right) \\ &\cong \text{Hom}_{\mathbf{Rel}(X,A)}\left(S, \int_{b \in B} \mathbf{Hom}_{\{t,f\}}(R_{-1}^b, T_{-2}^b)\right) \end{aligned}$$


naturally in each $S \in \mathbf{Rel}(X, A)$ and each $T \in \mathbf{Rel}(X, B)$, showing that

$$\int_{b \in B} \mathbf{Hom}_{\{t,f\}}(R_{-1}^b, S_{-2}^b)$$

is right adjoint to the postcomposition functor $R \diamond -$, being thus the right Kan lift along R . Here we have used the following results, respectively (i.e. for each \cong sign):

1. **Item 1** of **Proposition 5.1.1.6**.

2. **Definition 6.3.12.1.**
3. ?? of ??.
4. **Proposition 1.2.2.5.**
5. ?? of ??.
6. ?? of ??.
7. **Item 1 of Proposition 5.1.1.6.**

This finishes the proof. 

00NZ 6.3 More Constructions With Relations

00P0 6.3.1 The Graph of a Function

Let $f: A \rightarrow B$ be a function.

00P1 DEFINITION 6.3.1.1 ► THE GRAPH OF A FUNCTION

The **graph of f** is the relation $\text{Gr}(f): A \rightarrow B$ defined as follows:¹

- Viewing relations from A to B as subsets of $A \times B$, we define

$$\text{Gr}(f) \stackrel{\text{def}}{=} \{(a, f(a)) \in A \times B \mid a \in A\}.$$

- Viewing relations from A to B as functions $A \times B \rightarrow \{\text{true}, \text{false}\}$, we define

$$[\text{Gr}(f)](a, b) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } b = f(a), \\ \text{false} & \text{otherwise} \end{cases}$$

for each $(a, b) \in A \times B$.

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$[\text{Gr}(f)](a) \stackrel{\text{def}}{=} \{f(a)\}$$

for each $a \in A$, i.e. we define $\text{Gr}(f)$ as the composition

$$A \xrightarrow{f} B \xrightarrow{\chi_B} \mathcal{P}(B).$$

¹Further Notation: We write $\text{Gr}(A)$ for $\text{Gr}(\text{id}_A)$, and call it the **graph** of A .

00P2

PROPOSITION 6.3.1.2 ► PROPERTIES OF GRAPHS OF FUNCTIONS

Let $f: A \rightarrow B$ be a function.

00P3

1. *Functoriality.* The assignment $A \mapsto \text{Gr}(A)$ defines a functor

$$\text{Gr}: \text{Sets} \rightarrow \text{Rel}$$

where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$\text{Gr}(A) \stackrel{\text{def}}{=} A.$$

- *Action on Morphisms.* For each $A, B \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$\text{Gr}_{A,B}: \text{Sets}(A, B) \rightarrow \underbrace{\text{Rel}(\text{Gr}(A), \text{Gr}(B))}_{\stackrel{\text{def}}{=} \text{Rel}(A, B)}$$

of Gr at (A, B) is defined by

$$\text{Gr}_{A,B}(f) \stackrel{\text{def}}{=} \text{Gr}(f),$$

where $\text{Gr}(f)$ is the graph of f as in [Definition 6.3.1.1](#).

In particular:

- *Preservation of Identities.* We have

$$\text{Gr}(\text{id}_A) = \chi_A$$

for each $A \in \text{Obj}(\text{Sets})$.

- *Preservation of Composition.* We have

$$\text{Gr}(g \circ f) = \text{Gr}(g) \diamond \text{Gr}(f)$$

for each pair of functions $f: A \rightarrow B$ and $g: B \rightarrow C$.

00P4

2. *Adjointness Inside **Rel**.* We have an adjunction

$$(\text{Gr}(f) \dashv f^{-1}): A \begin{array}{c} \xrightarrow{\text{Gr}(f)} \\ \perp \\ \xleftarrow{f^{-1}} \end{array} B$$

in **Rel**, where f^{-1} is the inverse of f of [Definition 6.3.2.1](#).

00P5

3. *Adjointness.* We have an adjunction

$$(\text{Gr} \dashv \mathcal{P}_*): \text{Sets} \begin{array}{c} \xrightarrow{\text{Gr}} \\ \perp \\ \xleftarrow{\mathcal{P}_*} \end{array} \text{Rel},$$

witnessed by a bijection of sets

$$\text{Rel}(\text{Gr}(A), B) \cong \text{Sets}(A, \mathcal{P}(B))$$

natural in $A \in \text{Obj}(\text{Sets})$ and $B \in \text{Obj}(\text{Rel})$.

00P6

4. *Interaction With Inverses.* We have

$$\begin{aligned} \text{Gr}(f)^\dagger &= f^{-1}, \\ (f^{-1})^\dagger &= \text{Gr}(f). \end{aligned}$$

00P7

5. *Cocontinuity.* The functor $\text{Gr}: \text{Sets} \rightarrow \text{Rel}$ of [Item 1](#) preserves colimits.

00P8

6. *Characterisations.* Let $R: A \dashv B$ be a relation. The following conditions are equivalent:

00P9

(a) There exists a function $f: A \rightarrow B$ such that $R = \text{Gr}(f)$.

00PA

(b) The relation R is total and functional.

00PB

(c) The weak and strong inverse images of R agree, i.e. we have $R^{-1} = R_{-1}$.

00PC

(d) The relation R has a right adjoint R^\dagger in Rel .

PROOF 6.3.1.3 ► PROOF OF PROPOSITION 6.3.1.2

Item 1: Functoriality

Clear.

Item 2: Adjointness Inside Rel

We need to check that there are inclusions

$$\begin{aligned} \chi_A &\subset f^{-1} \diamond \text{Gr}(f), \\ \text{Gr}(f) \diamond f^{-1} &\subset \chi_B. \end{aligned}$$

These correspond respectively to the following conditions:

1. For each $a \in A$, there exists some $b \in B$ such that $a \sim_{\text{Gr}(f)} b$ and $b \sim_{f^{-1}} a$.
2. For each $a, b \in A$, if $a \sim_{\text{Gr}(f)} b$ and $b \sim_{f^{-1}} a$, then $a = b$.

In other words, the first condition states that the image of any $a \in A$ by f is nonempty, whereas the second condition states that f is not multivalued. As f is a function, both of these statements are true, and we are done.

Item 3: Adjointness

The stated bijection follows from [Remark 5.1.1.4](#), with naturality being clear.

Item 4: Interaction With Inverses

Clear.

Item 5: Cocontinuity

Omitted.

Item 6: Characterisations

We claim that [Items 6a to 6d](#) are indeed equivalent:

- [Item 6a](#) \iff [Item 6b](#). This is shown in the proof of ?? of ??.
- [Item 6b](#) \implies [Item 6c](#). If R is total and functional, then, for each $a \in A$, the set $R(a)$ is a singleton, implying that

$$R^{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid R(a) \cap V \neq \emptyset\},$$

$$R_{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid R(a) \subset V\}$$

are equal for all $V \in \mathcal{P}(B)$, as the conditions $R(a) \cap V \neq \emptyset$ and $R(a) \subset V$ are equivalent when $R(a)$ is a singleton.

- [Item 6c](#) \implies [Item 6b](#). We claim that R is indeed total and functional:
 - *Totality*. If we had $R(a) = \emptyset$ for some $a \in A$, then we would have $a \in R_{-1}(\emptyset)$, so that $R_{-1}(\emptyset) \neq \emptyset$. But since $R^{-1}(\emptyset) = \emptyset$, this would imply $R_{-1}(\emptyset) \neq R^{-1}(\emptyset)$, a contradiction. Thus $R(a) \neq \emptyset$ for all $a \in A$ and R is total.
 - *Functionality*. If $R^{-1} = R_{-1}$, then we have

$$\begin{aligned} \{a\} &= R^{-1}(\{b\}) \\ &= R_{-1}(\{b\}) \end{aligned}$$

for each $b \in R(a)$ and each $a \in A$, and thus $R(a) \subset \{b\}$. But since R is total, we must have $R(a) = \{b\}$, and thus we see that R is functional.

- *Item 6a* \iff *Item 6d*. This follows from [Proposition 5.3.3.1](#).

This finishes the proof. 

00PD 6.3.2 The Inverse of a Function

Let $f: A \rightarrow B$ be a function.

00PE DEFINITION 6.3.2.1 ► THE INVERSE OF A FUNCTION

The **inverse of f** is the relation $f^{-1}: B \rightarrow A$ defined as follows:

- Viewing relations from B to A as subsets of $B \times A$, we define

$$f^{-1} \stackrel{\text{def}}{=} \{(b, f^{-1}(b)) \in B \times A \mid a \in A\},$$

where

$$f^{-1}(b) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = b\}$$

for each $b \in B$.

- Viewing relations from B to A as functions $B \times A \rightarrow \{\text{true}, \text{false}\}$, we define

$$f^{-1}(b, a) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if there exists } a \in A \text{ with } f(a) = b, \\ \text{false} & \text{otherwise} \end{cases}$$

for each $(b, a) \in B \times A$.

- Viewing relations from B to A as functions $B \rightarrow \mathcal{P}(A)$, we define

$$f^{-1}(b) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = b\}$$

for each $b \in B$.

00PF PROPOSITION 6.3.2.2 ► PROPERTIES OF INVERSES OF FUNCTIONS

Let $f: A \rightarrow B$ be a function.

- 00PG 1. *Functoriality*. The assignment $A \mapsto A, f \mapsto f^{-1}$ defines a functor

$$(-)^{-1}: \text{Sets} \rightarrow \text{Rel}$$

where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$[(-)^{-1}](A) \stackrel{\text{def}}{=} A.$$

- *Action on Morphisms.* For each $A, B \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$(-)_{A,B}^{-1} : \text{Sets}(A, B) \rightarrow \text{Rel}(A, B)$$

of $(-)^{-1}$ at (A, B) is defined by

$$(-)_{A,B}^{-1}(f) \stackrel{\text{def}}{=} [(-)^{-1}](f),$$

where f^{-1} is the inverse of f as in [Definition 6.3.2.1](#).

In particular:

- *Preservation of Identities.* We have

$$\text{id}_A^{-1} = \chi_A$$

for each $A \in \text{Obj}(\text{Sets})$.

- *Preservation of Composition.* We have

$$(g \circ f)^{-1} = g^{-1} \diamond f^{-1}$$

for pair of functions $f: A \rightarrow B$ and $g: B \rightarrow C$.

00PH

2. *Adjointness Inside **Rel**.* We have an adjunction

$$(\text{Gr}(f) \dashv f^{-1}): A \begin{array}{c} \xrightarrow{\text{Gr}(f)} \\ \perp \\ \xleftarrow{f^{-1}} \end{array} B$$

in **Rel**.

00PJ

3. *Interaction With Inverses of Relations.* We have

$$(f^{-1})^\dagger = \text{Gr}(f),$$

$$\text{Gr}(f)^\dagger = f^{-1}.$$

PROOF 6.3.2.3 ► PROOF OF PROPOSITION 6.3.2.2

Item 1: Functoriality

Clear.

Item 2: Adjointness Inside Rel

This is proved in **Item 2** of **Proposition 6.3.1.2**.

Item 3: Interaction With Inverses of Relations

Clear. 

00PK 6.3.3 Representable Relations

Let A and B be sets.

00PL DEFINITION 6.3.3.1 ► REPRESENTABLE RELATIONS

Let $f: A \rightarrow B$ and $g: B \rightarrow A$ be functions.¹

1. The **representable relation associated to f** is the relation $\chi_f: A \rightarrow B$ defined as the composition

$$A \times B \xrightarrow{f \times \text{id}_B} B \times B \xrightarrow{\chi_B} \{\text{true}, \text{false}\},$$

i.e. given by declaring $a \sim_{\chi_f} b$ iff $f(a) = b$.

2. The **corepresentable relation associated to g** is the relation $\chi^g: B \rightarrow A$ defined as the composition

$$B \times A \xrightarrow{g \times \text{id}_A} A \times A \xrightarrow{\chi_A} \{\text{true}, \text{false}\},$$

i.e. given by declaring $b \sim_{\chi^g} a$ iff $g(b) = a$.¹More generally, given functions

$$f: A \rightarrow C,$$

$$g: B \rightarrow D$$

and a relation $B \rightarrow D$, we may consider the composite relation

$$A \times B \xrightarrow{f \times g} C \times D \xrightarrow{R} \{\text{true}, \text{false}\},$$

for which we have $a \sim_{R \circ (f \times g)} b$ iff $f(a) \sim_R g(b)$.

00PM 6.3.4 The Domain and Range of a Relation

Let A and B be sets.

00PN

DEFINITION 6.3.4.1 ► THE DOMAIN AND RANGE OF A RELATION

Let $R \subset A \times B$ be a relation.^{1,2}

1. The **domain of R** is the subset $\text{dom}(R)$ of A defined by

$$\text{dom}(R) \stackrel{\text{def}}{=} \left\{ a \in A \mid \begin{array}{l} \text{there exists some } b \in B \\ \text{such that } a \sim_R b \end{array} \right\}.$$

2. The **range of R** is the subset $\text{range}(R)$ of B defined by

$$\text{range}(R) \stackrel{\text{def}}{=} \left\{ b \in B \mid \begin{array}{l} \text{there exists some } a \in A \\ \text{such that } a \sim_R b \end{array} \right\}.$$

¹Following ??, we may compute the (characteristic functions associated to the) domain and range of a relation using the following colimit formulas:

$$\begin{aligned} \chi_{\text{dom}(R)}(a) &\cong \text{colim}_{b \in B} (R_a^b) & (a \in A) \\ &\cong \bigvee_{b \in B} R_a^b, \\ \chi_{\text{range}(R)}(b) &\cong \text{colim}_{a \in A} (R_a^b) & (b \in B) \\ &\cong \bigvee_{a \in A} R_a^b, \end{aligned}$$

where the join \bigvee is taken in the poset $(\{\text{true}, \text{false}\}, \preceq)$ of Definition 1.2.2.3.

²Viewing R as a function $R: A \rightarrow \mathcal{P}(B)$, we have

$$\begin{aligned} \text{dom}(R) &\cong \text{colim}_{y \in Y} (R(y)) \\ &\cong \bigcup_{y \in Y} R(y), \\ \text{range}(R) &\cong \text{colim}_{x \in X} (R(x)) \\ &\cong \bigcup_{x \in X} R(x), \end{aligned}$$

00PP 6.3.5 Binary Unions of Relations

Let A and B be sets and let R and S be relations from A to B .

00PQ

DEFINITION 6.3.5.1 ► BINARY UNIONS OF RELATIONS

The **union of R and S** ¹ is the relation $R \cup S$ from A to B defined as follows:

- Viewing relations from A to B as subsets of $A \times B$, we define²

$$R \cup S \stackrel{\text{def}}{=} \{(a, b) \in B \times A \mid \text{we have } a \sim_R b \text{ or } a \sim_S b\}.$$

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$[R \cup S](a) \stackrel{\text{def}}{=} R(a) \cup S(a)$$

for each $a \in A$.

¹Further Terminology: Also called the **binary union of R and S** , for emphasis.

²This is the same as the union of R and S as subsets of $A \times B$.

PROPOSITION 6.3.5.2 ► PROPERTIES OF BINARY UNIONS OF RELATIONS

Let R, S, R_1 , and R_2 be relations from A to B , and let S_1 and S_2 be relations from B to C .

- Interaction With Inverses.* We have

$$(R \cup S)^\dagger = R^\dagger \cup S^\dagger.$$

- Interaction With Composition.* We have

$$(S_1 \diamond R_1) \cup (S_2 \diamond R_2) \stackrel{\text{poss.}}{\neq} (S_1 \cup S_2) \diamond (R_1 \cup R_2).$$

PROOF 6.3.5.3 ► PROOF OF PROPOSITION 6.3.5.2

Item 1: Interaction With Inverses

Clear.

Item 2: Interaction With Composition

Unwinding the definitions, we see that:

- The condition for $(S_1 \diamond R_1) \cup (S_2 \diamond R_2)$ is:

(a) There exists some $b \in B$ such that:

i. $a \sim_{R_1} b$ and $b \sim_{S_1} c$;

or

i. $a \sim_{R_2} b$ and $b \sim_{S_2} c$;


- The condition for $(S_1 \cup S_2) \diamond (R_1 \cup R_2)$ is:

(a) There exists some $b \in B$ such that:

i. $a \sim_{R_1} b$ or $a \sim_{R_2} b$;

and

i. $b \sim_{S_1} c$ or $b \sim_{S_2} c$.

These two conditions may fail to agree (counterexample omitted), and thus the two resulting relations on $A \times C$ may differ. 

00PU 6.3.6 Unions of Families of Relations

Let A and B be sets and let $\{R_i\}_{i \in I}$ be a family of relations from A to B .

00PV DEFINITION 6.3.6.1 ► THE UNION OF A FAMILY OF RELATIONS

The **union of the family** $\{R_i\}_{i \in I}$ is the relation $\bigcup_{i \in I} R_i$ from A to B defined as follows:

- Viewing relations from A to B as subsets of $A \times B$, we define¹

$$\bigcup_{i \in I} R_i \stackrel{\text{def}}{=} \left\{ (a, b) \in (A \times B)^{\times I} \mid \begin{array}{l} \text{there exists some } i \in I \\ \text{such that } a \sim_{R_i} b \end{array} \right\}.$$

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$\left[\bigcup_{i \in I} R_i \right] (a) \stackrel{\text{def}}{=} \bigcup_{i \in I} R_i(a)$$

for each $a \in A$.

¹This is the same as the union of $\{R_i\}_{i \in I}$ as a collection of subsets of $A \times B$.

00PW PROPOSITION 6.3.6.2 ► PROPERTIES OF UNIONS OF FAMILIES OF RELATIONS

Let A and B be sets and let $\{R_i\}_{i \in I}$ be a family of relations from A to B .

- 00PX 1. *Interaction With Inverses.* We have

$$\left(\bigcup_{i \in I} R_i \right)^\dagger = \bigcup_{i \in I} R_i^\dagger.$$

PROOF 6.3.6.3 ► PROOF OF PROPOSITION 6.3.6.2

Item 1: Interaction With Inverses

Clear.

00PY 6.3.7 Binary Intersections of Relations

Let A and B be sets and let R and S be relations from A to B .

00PZ DEFINITION 6.3.7.1 ► BINARY INTERSECTIONS OF RELATIONS

The **intersection of R and S** ¹ is the relation $R \cap S$ from A to B defined as follows:

- Viewing relations from A to B as subsets of $A \times B$, we define²

$$R \cap S \stackrel{\text{def}}{=} \{(a, b) \in B \times A \mid \text{we have } a \sim_R b \text{ and } a \sim_S b\}.$$

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$[R \cap S](a) \stackrel{\text{def}}{=} R(a) \cap S(a)$$

for each $a \in A$.¹Further Terminology: Also called the **binary intersection of R and S** , for emphasis.²This is the same as the intersection of R and S as subsets of $A \times B$.

00Q0 PROPOSITION 6.3.7.2 ► PROPERTIES OF BINARY INTERSECTIONS OF RELATIONS

Let R, S, R_1 , and R_2 be relations from A to B , and let S_1 and S_2 be relations from B to C .

- 00Q1 1.
- Interaction With Inverses.*
- We have

$$(R \cap S)^\dagger = R^\dagger \cap S^\dagger.$$

- 00Q2 2.
- Interaction With Composition.*
- We have

$$(S_1 \diamond R_1) \cap (S_2 \diamond R_2) = (S_1 \cap S_2) \diamond (R_1 \cap R_2).$$

PROOF 6.3.7.3 ► PROOF OF PROPOSITION 6.3.7.2

Item 1: Interaction With Inverses

Clear.

Item 2: Interaction With Composition

Unwinding the definitions, we see that:

1. The condition for $(S_1 \diamond R_1) \cap (S_2 \diamond R_2)$ is:

(a) There exists some $b \in B$ such that:

i. $a \sim_{R_1} b$ and $b \sim_{S_1} c$;

and

i. $a \sim_{R_2} b$ and $b \sim_{S_2} c$;


3. The condition for $(S_1 \cap S_2) \diamond (R_1 \cap R_2)$ is:

(a) There exists some $b \in B$ such that:

i. $a \sim_{R_1} b$ and $a \sim_{R_2} b$;

and

i. $b \sim_{S_1} c$ and $b \sim_{S_2} c$.

These two conditions agree, and thus so do the two resulting relations on $A \times C$. 

00Q3 6.3.8 Intersections of Families of Relations

Let A and B be sets and let $\{R_i\}_{i \in I}$ be a family of relations from A to B .

DEFINITION 6.3.8.1 ► THE INTERSECTION OF A FAMILY OF RELATIONS

The **intersection of the family** $\{R_i\}_{i \in I}$ is the relation $\bigcup_{i \in I} R_i$ defined as follows:

- Viewing relations from A to B as subsets of $A \times B$, we define¹

$$\bigcup_{i \in I} R_i \stackrel{\text{def}}{=} \left\{ (a, b) \in (A \times B)^{\times I} \left| \begin{array}{l} \text{for each } i \in I, \\ \text{we have } a \sim_{R_i} b \end{array} \right. \right\}.$$

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$\left[\bigcap_{i \in I} R_i \right] (a) \stackrel{\text{def}}{=} \bigcap_{i \in I} R_i(a)$$

for each $a \in A$.

¹This is the same as the intersection of $\{R_i\}_{i \in I}$ as a collection of subsets of $A \times B$.

00Q5

PROPOSITION 6.3.8.2 ► PROPERTIES OF INTERSECTIONS OF FAMILIES OF RELATIONS

Let A and B be sets and let $\{R_i\}_{i \in I}$ be a family of relations from A to B .

00Q6

1. *Interaction With Inverses.* We have

$$\left(\bigcap_{i \in I} R_i \right)^\dagger = \bigcap_{i \in I} R_i^\dagger.$$

PROOF 6.3.8.3 ► PROOF OF PROPOSITION 6.3.8.2

Item 1: Interaction With Inverses

Clear. 

00Q7 6.3.9 Binary Products of Relations

Let A, B, X , and Y be sets, let $R: A \rightarrow B$ be a relation from A to B , and let $S: X \rightarrow Y$ be a relation from X to Y .

00Q8

DEFINITION 6.3.9.1 ► BINARY PRODUCTS OF RELATIONS

The **product of R and S** ¹ is the relation $R \times S$ from $A \times X$ to $B \times Y$ defined as follows:

- Viewing relations from $A \times X$ to $B \times Y$ as subsets of $(A \times X) \times (B \times Y)$, we define $R \times S$ as the Cartesian product of R and S as subsets of $A \times X$ and $B \times Y$.²
- Viewing relations from $A \times X$ to $B \times Y$ as functions $A \times X \rightarrow \mathcal{P}(B \times Y)$, we define $R \times S$ as the composition

$$A \times X \xrightarrow{R \times S} \mathcal{P}(B) \times \mathcal{P}(Y) \xrightarrow{\mathcal{P}_{B,Y}^\circ} \mathcal{P}(B \times Y)$$

in Sets, i.e. by

$$[R \times S](a, x) \stackrel{\text{def}}{=} R(a) \times S(x)$$

for each $(a, x) \in A \times X$.

¹*Further Terminology:* Also called the **binary product of R and S** for emphasis. That is, $R \times S$ is the relation given by declaring $(u, x) \sim_{R \times S} (v, y)$ iff $u \sim_R v$ and $x \sim_S y$.

00Q9

PROPOSITION 6.3.9.2 ► PROPERTIES OF BINARY PRODUCTS OF RELATIONS

Let A, B, X , and Y be sets.

00QA

1. *Interaction With Inverses.* Let

$$R: A \rightarrow A,$$

$$S: X \rightarrow X$$

We have

$$(R \times S)^\dagger = R^\dagger \times S^\dagger.$$

00QB

2. *Interaction With Composition.* Let

$$R_1: A \rightarrow B,$$

$$S_1: B \rightarrow C,$$

$$R_2: X \rightarrow Y,$$

$$S_2: Y \rightarrow Z$$

be relations. We have

$$(S_1 \circ R_1) \times (S_2 \circ R_2) = (S_1 \times S_2) \circ (R_1 \times R_2).$$

PROOF 6.3.9.3 ► PROOF OF PROPOSITION 6.3.9.2**Item 1: Interaction With Inverses**

Unwinding the definitions, we see that:

1. We have $(a, x) \sim_{(R \times S)^\dagger} (b, y)$ iff:

· We have $(b, y) \sim_{R \times S} (a, x)$, i.e. iff:

– We have $b \sim_R a$;

– We have $y \sim_S x$;

2. We have $(a, x) \sim_{R^\dagger \times S^\dagger} (b, y)$ iff:

· We have $a \sim_{R^\dagger} b$ and $x \sim_{S^\dagger} y$, i.e. iff:

– We have $b \sim_R a$;


– We have $y \sim_S x$.

These two conditions agree, and thus the two resulting relations on $A \times X$ are equal.

Item 2: Interaction With Composition

Unwinding the definitions, we see that:

1. We have $(a, x) \sim_{(S_1 \circ R_1) \times (S_2 \circ R_2)} (c, z)$ iff:
 - (a) We have $a \sim_{S_1 \circ R_1} c$ and $x \sim_{S_2 \circ R_2} z$, i.e. iff:
 - i. There exists some $b \in B$ such that $a \sim_{R_1} b$ and $b \sim_{S_1} c$;
 - ii. There exists some $y \in Y$ such that $x \sim_{R_2} y$ and $y \sim_{S_2} z$;
2. We have $(a, x) \sim_{(S_1 \times S_2) \circ (R_1 \times R_2)} (c, z)$ iff:
 - (a) There exists some $(b, y) \in B \times Y$ such that $(a, x) \sim_{R_1 \times R_2} (b, y)$ and $(b, y) \sim_{S_1 \times S_2} (c, z)$, i.e. such that:
 - i. We have $a \sim_{R_1} b$ and $x \sim_{R_2} y$;
 - ii. We have $b \sim_{S_1} c$ and $y \sim_{S_2} z$.

These two conditions agree, and thus the two resulting relations from $A \times X$ to $C \times Z$ are equal. 

00QC 6.3.10 Products of Families of Relations

Let $\{A_i\}_{i \in I}$ and $\{B_i\}_{i \in I}$ be families of sets, and let $\{R_i : A_i \rightarrow B_i\}_{i \in I}$ be a family of relations.

00QD DEFINITION 6.3.10.1 ► THE PRODUCT OF A FAMILY OF RELATIONS

The **product of the family** $\{R_i\}_{i \in I}$ is the relation $\prod_{i \in I} R_i$ from $\prod_{i \in I} A_i$ to $\prod_{i \in I} B_i$ defined as follows:

- Viewing relations as subsets, we define $\prod_{i \in I} R_i$ as its product as a family of sets, i.e. we have

$$\prod_{i \in I} R_i \stackrel{\text{def}}{=} \left\{ (a_i, b_i)_{i \in I} \in \prod_{i \in I} (A_i \times B_i) \mid \begin{array}{l} \text{for each } i \in I, \\ \text{we have } a_i \sim_{R_i} b_i \end{array} \right\}.$$

- Viewing relations as functions to powersets, we define

$$\left[\prod_{i \in I} R_i \right] ((a_i)_{i \in I}) \stackrel{\text{def}}{=} \prod_{i \in I} R_i(a_i)$$

for each $(a_i)_{i \in I} \in \prod_{i \in I} A_i$.

00QE 6.3.11 The Inverse of a Relation

Let A, B , and C be sets and let $R \subset A \times B$ be a relation.

00QF DEFINITION 6.3.11.1 ► THE INVERSE OF A RELATION

The **inverse of R** ¹ is the relation R^\dagger defined as follows:

- Viewing relations as subsets, we define

$$R^\dagger \stackrel{\text{def}}{=} \{(b, a) \in B \times A \mid \text{we have } b \sim_R a\}.$$

- Viewing relations as functions $A \times B \rightarrow \{\text{true}, \text{false}\}$, we define

$$[R^\dagger]_b^a \stackrel{\text{def}}{=} R_a^b$$

for each $(b, a) \in B \times A$.

- Viewing relations as functions $A \rightarrow \mathcal{P}(B)$, we define

$$\begin{aligned} [R^\dagger](b) &\stackrel{\text{def}}{=} R^\dagger(\{b\}) \\ &\stackrel{\text{def}}{=} \{a \in A \mid b \in R(a)\} \end{aligned}$$

for each $b \in B$, where $R^\dagger(\{b\})$ is the fibre of R over $\{b\}$.

¹Further Terminology: Also called the **opposite of R** , the **transpose of R** , or the **converse of R** .

00QG EXAMPLE 6.3.11.2 ► EXAMPLES OF INVERSES OF RELATIONS

Here are some examples of inverses of relations.

1. *Less Than Equal Signs.* We have $(\leq)^\dagger = \geq$.
2. *Greater Than Equal Signs.* Dually to **Item 1**, we have $(\geq)^\dagger = \leq$.
3. *Functions.* Let $f: A \rightarrow B$ be a function. We have

$$\begin{aligned} \text{Gr}(f)^\dagger &= f^{-1}, \\ (f^{-1})^\dagger &= \text{Gr}(f). \end{aligned}$$

00QL PROPOSITION 6.3.11.3 ► PROPERTIES OF INVERSES OF RELATIONS

Let $R: A \rightarrow B$ and $S: B \rightarrow C$ be relations.

1. *Functoriality.* The assignment $R \mapsto R^\dagger$ defines a functor (i.e. morphism

of posets)

$$(-)^\dagger : \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(B, A).$$

In particular, given relations $R, S : A \rightrightarrows B$, we have:

$$(\star) \text{ If } R \subset S, \text{ then } R^\dagger \subset S^\dagger.$$

00QN

2. *Interaction With Ranges and Domains.* We have

$$\begin{aligned} \text{dom}(R^\dagger) &= \text{range}(R), \\ \text{range}(R^\dagger) &= \text{dom}(R). \end{aligned}$$

00QP

3. *Interaction With Composition I.* We have

$$(S \diamond R)^\dagger = R^\dagger \diamond S^\dagger.$$

00QQ

4. *Interaction With Composition II.* We have

$$\begin{aligned} \chi_B &\subset R \diamond R^\dagger, \\ \chi_A &\subset R^\dagger \diamond R. \end{aligned}$$

00QR

5. *Invertibility.* We have

$$(R^\dagger)^\dagger = R.$$

00QS

6. *Identity.* We have

$$\chi_A^\dagger = \chi_A.$$

PROOF 6.3.11.4 ► PROOF OF PROPOSITION 6.3.11.3

Item 1: Functoriality

Clear.

Item 2: Interaction With Ranges and Domains

Clear.

Item 3: Interaction With Composition I

Clear.

Item 4: Interaction With Composition II

Clear.

Item 5: Invertibility

Clear.

Item 6: Identity

Clear.



00QT 6.3.12 Composition of Relations

Let A, B , and C be sets and let $R: A \rightarrow B$ and $S: B \rightarrow C$ be relations.

00QU DEFINITION 6.3.12.1 ► COMPOSITION OF RELATIONS

The **composition of R and S** is the relation $S \diamond R$ defined as follows:

- Viewing relations from A to C as subsets of $A \times C$, we define

$$S \diamond R \stackrel{\text{def}}{=} \left\{ (a, c) \in A \times C \mid \begin{array}{l} \text{there exists some } b \in B \text{ such} \\ \text{that } a \sim_R b \text{ and } b \sim_S c \end{array} \right\}.$$

- Viewing relations as functions $A \times B \rightarrow \{\text{true}, \text{false}\}$, we define

$$\begin{aligned} (S \diamond R)_{-2}^{-1} &\stackrel{\text{def}}{=} \int^{b \in B} S_b^{-1} \times R_{-2}^b \\ &= \bigvee_{b \in B} S_b^{-1} \times R_{-2}^b, \end{aligned}$$

where the join \bigvee is taken in the poset $(\{\text{true}, \text{false}\}, \preceq)$ of [Definition 1.2.2.3](#).

- Viewing relations as functions $A \rightarrow \mathcal{P}(B)$, we define

$$S \diamond R \stackrel{\text{def}}{=} \text{Lan}_{\chi_B}(S) \circ R,$$

where $\text{Lan}_{\chi_B}(S)$ is computed by the formula

$$\begin{aligned} [\text{Lan}_{\chi_B}(S)](V) &\cong \int^{y \in B} \chi_{\mathcal{P}(B)}(\chi_y, V) \odot S_y \\ &\cong \int^{y \in B} \chi_V(y) \odot S_y \\ &\cong \bigcup_{y \in B} \chi_V(y) \odot S_y \\ &\cong \bigcup_{y \in V} S_y \end{aligned}$$

for each $V \in \mathcal{P}(B)$. In other words, $S \diamond R$ is defined by¹

$$\begin{aligned} [S \diamond R](a) &\stackrel{\text{def}}{=} S(R(a)) \\ &\stackrel{\text{def}}{=} \bigcup_{x \in R(a)} S(x). \end{aligned}$$

for each $a \in A$.

¹That is: the relation R may send $a \in A$ to a number of elements $\{b_i\}_{i \in I}$ in B , and then the relation S may send the image of each of the b_i 's to a number of elements $\{S(b_i)\}_{i \in I} = \{\{c_{ji}\}_{j_i \in J_i}\}_{i \in I}$ in C .

00QV

EXAMPLE 6.3.12.2 ► EXAMPLES OF COMPOSITION OF RELATIONS

Here are some examples of composition of relations.

1. *Composing Less/Greater Than Equal With Greater/Less Than Equal Signs.* We have

$$\begin{aligned} \leq \diamond \geq &= \sim_{\text{triv}}, \\ \geq \diamond \leq &= \sim_{\text{triv}}. \end{aligned}$$

2. *Composing Less/Greater Than Equal Signs With Less/Greater Than Equal Signs.* We have

$$\begin{aligned} \leq \diamond \leq &= \leq, \\ \geq \diamond \geq &= \geq. \end{aligned}$$

00QW

PROPOSITION 6.3.12.3 ► PROPERTIES OF COMPOSITION OF RELATIONS

Let $R: A \rightarrow B$, $S: B \rightarrow C$, and $T: C \rightarrow D$ be relations.

00QX

1. *Interaction With Ranges and Domains.* We have

$$\begin{aligned}\text{dom}(S \diamond R) &\subset \text{dom}(R), \\ \text{range}(S \diamond R) &\subset \text{range}(S).\end{aligned}$$

00QY

2. *Associativity.* We have

$$(T \diamond S) \diamond R = T \diamond (S \diamond R).$$

00QZ

3. *Unitality.* We have

$$\begin{aligned}\chi_B \diamond R &= R, \\ R \diamond \chi_A &= R.\end{aligned}$$

00R0

4. *Interaction With Inverses.* We have

$$(S \diamond R)^\dagger = R^\dagger \diamond S^\dagger.$$

00R1

5. *Interaction With Composition.* We have

$$\begin{aligned}\chi_B &\subset R \diamond R^\dagger, \\ \chi_A &\subset R^\dagger \diamond R.\end{aligned}$$

PROOF 6.3.12.4 ► PROOF OF PROPOSITION 6.3.12.3

Item 1: Interaction With Ranges and Domains

Clear.

Item 2: Associativity

Indeed, we have

$$\begin{aligned}
(T \diamond S) \diamond R &\stackrel{\text{def}}{=} \left(\int^{c \in C} T_c^{-1} \times S_{-2}^c \right) \diamond R \\
&\stackrel{\text{def}}{=} \int^{b \in B} \left(\int^{c \in C} T_c^{-1} \times S_b^c \right) \diamond R_{-2}^b \\
&= \int^{b \in B} \int^{c \in C} (T_c^{-1} \times S_b^c) \diamond R_{-2}^b \\
&= \int^{c \in C} \int^{b \in B} (T_c^{-1} \times S_b^c) \diamond R_{-2}^b \\
&= \int^{c \in C} \int^{b \in B} T_c^{-1} \times (S_b^c \diamond R_{-2}^b) \\
&= \int^{c \in C} T_c^{-1} \times \left(\int^{b \in B} S_b^c \diamond R_{-2}^b \right) \\
&\stackrel{\text{def}}{=} \int^{c \in C} T_c^{-1} \times (S \diamond R)_{-2}^c \\
&\stackrel{\text{def}}{=} T \diamond (S \diamond R).
\end{aligned}$$

In the language of relations, given $a \in A$ and $d \in D$, the stated equality witnesses the equivalence of the following two statements:

1. We have $a \sim_{(T \diamond S) \diamond R} d$, i.e. there exists some $b \in B$ such that:
 - (a) We have $a \sim_R b$;
 - (b) We have $b \sim_{T \diamond S} d$, i.e. there exists some $c \in C$ such that:
 - i. We have $b \sim_S c$;
 - ii. We have $c \sim_T d$;
2. We have $a \sim_{T \diamond (S \diamond R)} d$, i.e. there exists some $c \in C$ such that:
 - (a) We have $a \sim_{S \diamond R} c$, i.e. there exists some $b \in B$ such that:
 - i. We have $a \sim_R b$;
 - ii. We have $b \sim_S c$;
 - (b) We have $c \sim_T d$;

both of which are equivalent to the statement

- There exist $b \in B$ and $c \in C$ such that $a \sim_R b \sim_S c \sim_T d$.

Item 3: Unitality

Indeed, we have

$$\begin{aligned}\chi_B \diamond R &\stackrel{\text{def}}{=} \int^{x \in B} (\chi_B)_x^{-1} \times R_{-2}^x \\ &= \bigvee_{x \in B} (\chi_B)_x^{-1} \times R_{-2}^x \\ &= \bigvee_{\substack{x \in B \\ x = -1}} R_{-2}^x \\ &= R_{-2}^{-1},\end{aligned}$$

and

$$\begin{aligned}R \diamond \chi_A &\stackrel{\text{def}}{=} \int^{x \in A} R_x^{-1} \times (\chi_A)_{-2}^x \\ &= \bigvee_{x \in B} R_x^{-1} \times (\chi_A)_{-2}^x \\ &= \bigvee_{\substack{x \in B \\ x = -2}} R_x^{-1} \\ &= R_{-2}^{-1}.\end{aligned}$$

In the language of relations, given $a \in A$ and $b \in B$:

- The equality

$$\chi_B \diamond R = R$$

witnesses the equivalence of the following two statements:

1. We have $a \sim_b B$.
2. There exists some $b' \in B$ such that:
 - (a) We have $a \sim_R b'$
 - (b) We have $b' \sim_{\chi_B} b$, i.e. $b' = b$.

- The equality

$$R \diamond \chi_A = R$$

witnesses the equivalence of the following two statements:

1. There exists some $a' \in A$ such that:
 - (a) We have $a \sim_{\chi_B} a'$, i.e. $a = a'$.


(b) We have $a' \sim_R b$

2. We have $a \sim_b B$.

Item 4: Interaction With Inverses

Clear.

Item 5: Interaction With Composition

Clear. 

00R2 6.3.13 The Collage of a Relation

Let A and B be sets and let $R: A \rightarrow B$ be a relation from A to B .

00R3 DEFINITION 6.3.13.1 ► THE COLLAGE OF A RELATION

The **collage of R** ¹ is the poset $\mathbf{Coll}(R) \stackrel{\text{def}}{=} (\text{Coll}(R), \preceq_{\mathbf{Coll}(R)})$ consisting of:

- *The Underlying Set.* The set $\text{Coll}(R)$ defined by

$$\text{Coll}(R) \stackrel{\text{def}}{=} A \amalg B.$$

- *The Partial Order.* The partial order

$$\preceq_{\mathbf{Coll}(R)}: \text{Coll}(R) \times \text{Coll}(R) \rightarrow \{\text{true}, \text{false}\}$$

on $\text{Coll}(R)$ defined by

$$\preceq(a, b) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } a = b \text{ or } a \sim_R b, \\ \text{false} & \text{otherwise.} \end{cases}$$

¹Further Terminology: Also called the **cograph** of R .

00R4 PROPOSITION 6.3.13.2 ► PROPERTIES OF COLLAGES OF RELATIONS

Let A and B be sets and let $R: A \rightarrow B$ be a relation from A to B .

- 00R5 1. *Functoriality I.* The assignment $R \mapsto \mathbf{Coll}(R)$ defines a functor¹

$$\mathbf{Coll}: \mathbf{Rel}(A, B) \rightarrow \text{Pos}_{/\Delta^1}(A, B),$$

where

- *Action on Objects.* For each $R \in \text{Obj}(\mathbf{Rel}(A, B))$, we have

$$[\mathbf{Coll}](R) \stackrel{\text{def}}{=} (\mathbf{Coll}(R), \phi_R)$$

for each $R \in \mathbf{Rel}(A, B)$, where

- The poset $\mathbf{Coll}(R)$ is the collage of R of [Definition 6.3.13.1](#).
- The morphism $\phi_R: \mathbf{Coll}(R) \rightarrow \Delta^1$ is given by

$$\phi_R(x) \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } x \in A, \\ 1 & \text{if } x \in B \end{cases}$$

for each $x \in \mathbf{Coll}(R)$.

- *Action on Morphisms.* For each $R, S \in \text{Obj}(\mathbf{Rel}(A, B))$, the action on Hom-sets

$$\mathbf{Coll}_{R,S}: \text{Hom}_{\mathbf{Rel}(A,B)}(R, S) \rightarrow \text{Pos}(\mathbf{Coll}(R), \mathbf{Coll}(S))$$

of \mathbf{Coll} at (R, S) is given by sending an inclusion

$$\iota: R \subset S$$

to the morphism

$$\mathbf{Coll}(\iota): \mathbf{Coll}(R) \rightarrow \mathbf{Coll}(S)$$

of posets over Δ^1 defined by

$$[\mathbf{Coll}(\iota)](x) \stackrel{\text{def}}{=} x$$

for each $x \in \mathbf{Coll}(R)$.²


00R6

2. *Equivalence.* The functor of [Item 1](#) is an equivalence of categories.

¹Here $\text{Pos}_{/\Delta^1}(A, B)$ is the category defined as the pullback

$$\text{Pos}_{/\Delta^1}(A, B) \stackrel{\text{def}}{=} \text{pt} \times_{[A], \text{Pos}, \text{fib}_0} \text{Pos}_{/\Delta^1} \times_{\text{fib}_1, \text{Pos}, [B]} \text{pt},$$

as in the diagram



Explicitly, an object of $\text{Pos}_{/\Delta^1}(A, B)$ is a pair (X, ϕ_X) consisting of

- A poset X ;
- A morphism $\phi_X: X \rightarrow \Delta^1$;

such that $\phi_X^{-1}(0) = A$ and $\phi_X^{-1}(1) = B$, with morphisms between such objects being morphisms of posets over Δ^1 .

²Note that this is indeed a morphism of posets: if $x \preceq_{\mathbf{Coll}(R)} y$, then $x = y$ or $x \sim_R y$, so we have either $x = y$ or $x \sim_S y$ (as $R \subset S$), and thus $x \preceq_{\mathbf{Coll}(S)} y$.

PROOF 6.3.13.3 ► PROOF OF PROPOSITION 6.3.13.2

Item 1: Functoriality

Clear.

Item 2: Equivalence

Omitted. 

00R7 6.4 Functoriality of Powersets

00R8 6.4.1 Direct Images

Let A and B be sets and let $R: A \rightarrow B$ be a relation.

00R9 DEFINITION 6.4.1.1 ► DIRECT IMAGES

The **direct image function associated to R** is the function

$$R_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by^{1,2}

$$\begin{aligned} R_*(U) &\stackrel{\text{def}}{=} R(U) \\ &\stackrel{\text{def}}{=} \bigcup_{a \in U} R(a) \\ &= \left\{ b \in B \mid \begin{array}{l} \text{there exists some } a \in U \\ \text{such that } b \in R(a) \end{array} \right\} \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

¹Further Terminology: The set $R(U)$ is called the **direct image of U by R** .

²We also have

$$R_*(U) = B \setminus R_*(A \setminus U);$$

see Item 7 of Proposition 6.4.1.3.

00RA **REMARK 6.4.1.2 ▶ UNWINDING DEFINITION 6.4.1.1**

Identifying subsets of A with relations from pt to A via **Item 3** of **Proposition 2.4.3.9**, we see that the direct image function associated to R is equivalently the function

$$R_* : \underbrace{\mathcal{P}(A)}_{\cong \text{Rel}(\text{pt}, A)} \rightarrow \underbrace{\mathcal{P}(B)}_{\cong \text{Rel}(\text{pt}, B)}$$

defined by

$$R_*(U) \stackrel{\text{def}}{=} R \diamond U$$

for each $U \in \mathcal{P}(A)$, where $R \diamond U$ is the composition

$$\text{pt} \xrightarrow{U} A \xrightarrow{R} B.$$

00RB **PROPOSITION 6.4.1.3 ▶ PROPERTIES OF DIRECT IMAGE FUNCTIONS**

Let $R: A \rightarrow B$ be a relation.

00RC 1. *Functoriality.* The assignment $U \mapsto R_*(U)$ defines a functor

$$R_* : (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

where

• *Action on Objects.* For each $U \in \mathcal{P}(A)$, we have

$$[R_*](U) \stackrel{\text{def}}{=} R_*(U).$$

• *Action on Morphisms.* For each $U, V \in \mathcal{P}(A)$:

– If $U \subset V$, then $R_*(U) \subset R_*(V)$.

00RD 2. *Adjointness.* We have an adjunction

$$(R_* \dashv R_{-1}) : \mathcal{P}(A) \begin{array}{c} \xrightarrow{R_*} \\ \perp \\ \xleftarrow{R_{-1}} \end{array} \mathcal{P}(B),$$

witnessed by a bijections of sets

$$\text{Hom}_{\mathcal{P}(A)}(R_*(U), V) \cong \text{Hom}_{\mathcal{P}(A)}(U, R_{-1}(V)),$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$, i.e. such that:

(★) The following conditions are equivalent:

- We have $R_*(U) \subset V$.
- We have $U \subset R_{-1}(V)$.

00RE

3. *Preservation of Colimits.* We have an equality of sets

$$R_*\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} R_*(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have equalities

$$\begin{aligned} R_*(U) \cup R_*(V) &= R_*(U \cup V), \\ R_*(\emptyset) &= \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00RF

4. *Oplax Preservation of Limits.* We have an inclusion of sets

$$R_*\left(\bigcap_{i \in I} U_i\right) \subset \bigcap_{i \in I} R_*(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} R_*(U \cap V) &\subset R_*(U) \cap R_*(V), \\ R_*(A) &\subset B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00RG

5. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of [Item 1](#) has a symmetric strict monoidal structure

$$\left(R_*, R_*^\otimes, R_{*|\mathbb{1}}^\otimes\right): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} R_{*|U,V}^\otimes: R_*(U) \cup R_*(V) &\xrightarrow{=} R_*(U \cup V), \\ R_{*|\mathbb{1}}^\otimes: \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00RH

6. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric oplax monoidal structure

$$\left(R_*, R_*^\otimes, R_{*|\mathbb{1}}^\otimes \right): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with inclusions

$$\begin{aligned} R_{*|U,V}^\otimes: R_*(U \cap V) &\subset R_*(U) \cap R_*(V), \\ R_{*|\mathbb{1}}^\otimes: R_*(A) &\subset B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00RJ

7. *Relation to Direct Images With Compact Support.* We have

$$R_*(U) = B \setminus R_!(A \setminus U)$$

for each $U \in \mathcal{P}(A)$.

PROOF 6.4.1.4 ► PROOF OF PROPOSITION 6.4.1.3

Item 1: Functoriality

Clear.

Item 2: Adjointness

This follows from ?? of ??.

Item 3: Preservation of Colimits

This follows from **Item 2** and ?? of ??.

Item 4: Oplax Preservation of Limits

Omitted.

Item 5: Symmetric Strict Monoidality With Respect to Unions

This follows from **Item 3**.

Item 6: Symmetric Oplax Monoidality With Respect to Intersections

This follows from **Item 4**.

Item 7: Relation to Direct Images With Compact Support


The proof proceeds in the same way as in the case of functions (?? of **Proposi-**

tion 2.4.4.4): applying **Item 7** of **Proposition 6.4.4.4** to $A \setminus U$, we have

$$\begin{aligned} R_!(A \setminus U) &= B \setminus R_*(A \setminus (A \setminus U)) \\ &= B \setminus R_*(U). \end{aligned}$$

Taking complements, we then obtain

$$\begin{aligned} R_*(U) &= B \setminus (B \setminus R_!(A \setminus U)), \\ &= B \setminus R_!(A \setminus U), \end{aligned}$$

which finishes the proof. 

00RK

PROPOSITION 6.4.1.5 ► PROPERTIES OF THE DIRECT IMAGE FUNCTION OPERATION

Let $R: A \rightarrow B$ be a relation.

00RL

1. *Functionality I.* The assignment $R \mapsto R_*$ defines a function

$$(-)_*: \text{Rel}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

00RM

2. *Functionality II.* The assignment $R \mapsto R_*$ defines a function

$$(-)_*: \text{Rel}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

00RN

3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have¹

$$(\chi_A)_* = \text{id}_{\mathcal{P}(A)}.$$

00RP

4. *Interaction With Composition.* For each pair of composable relations $R: A \rightarrow B$ and $S: B \rightarrow C$, we have²

$$(S \diamond R)_* = S_* \circ R_*,$$

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{R_*} & \mathcal{P}(B) \\ & \searrow (S \diamond R)_* & \downarrow S_* \\ & & \mathcal{P}(C). \end{array}$$

¹That is, the postcomposition function

$$(\chi_A)_*: \text{Rel}(\text{pt}, A) \rightarrow \text{Rel}(\text{pt}, A)$$

is equal to $\text{id}_{\text{Rel}(\text{pt}, A)}$.

²That is, we have

$$(S \diamond R)_* = S_* \circ R_*$$

$$\begin{array}{ccc} \text{Rel}(\text{pt}, A) & \xrightarrow{R_*} & \text{Rel}(\text{pt}, B) \\ & \searrow (S \diamond R)_* & \downarrow S_* \\ & & \text{Rel}(\text{pt}, C). \end{array}$$

PROOF 6.4.1.6 ► PROOF OF PROPOSITION 6.4.1.5

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

Indeed, we have

$$\begin{aligned} (\chi_A)_*(U) &\stackrel{\text{def}}{=} \bigcup_{a \in U} \chi_A(a) \\ &\stackrel{\text{def}}{=} \bigcup_{a \in U} \{a\} \\ &= U \\ &\stackrel{\text{def}}{=} \text{id}_{\mathcal{P}(A)}(U) \end{aligned}$$

for each $U \in \mathcal{P}(A)$. Thus $(\chi_A)_* = \text{id}_{\mathcal{P}(A)}$.

Item 4: Interaction With Composition

Indeed, we have

$$\begin{aligned} (S \diamond R)_*(U) &\stackrel{\text{def}}{=} \bigcup_{a \in U} [S \diamond R](a) \\ &\stackrel{\text{def}}{=} \bigcup_{a \in U} S(R(a)) \\ &\stackrel{\text{def}}{=} \bigcup_{a \in U} S_*(R(a)) \\ &= S_* \left(\bigcup_{a \in U} R(a) \right) \\ &\stackrel{\text{def}}{=} S_*(R_*(U)) \\ &\stackrel{\text{def}}{=} [S_* \circ R_*](U) \end{aligned}$$

for each $U \in \mathcal{P}(A)$, where we used **Item 3** of **Proposition 6.4.1.3**. Thus $(S \diamond R)_* = S_* \circ R_*$. 

00RQ 6.4.2 Strong Inverse Images

Let A and B be sets and let $R: A \rightarrow B$ be a relation.

00RR DEFINITION 6.4.2.1 ► STRONG INVERSE IMAGES

The **strong inverse image function associated to R** is the function

$$R_{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by¹

$$R_{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid R(a) \subset V\}$$

for each $V \in \mathcal{P}(B)$.


¹*Further Terminology:* The set $R_{-1}(V)$ is called the **strong inverse image of V by R** .

00RS REMARK 6.4.2.2 ► UNWINDING DEFINITION 6.4.2.1

Identifying subsets of B with relations from pt to B via **Item 3** of **Proposition 2.4.3.9**, we see that the inverse image function associated to R is equivalently the function

$$R_{-1}: \underbrace{\mathcal{P}(B)}_{\cong \text{Rel}(\text{pt}, B)} \rightarrow \underbrace{\mathcal{P}(A)}_{\cong \text{Rel}(\text{pt}, A)}$$

defined by

$$R_{-1}(V) \stackrel{\text{def}}{=} \text{Rift}_R(V),$$


and being explicitly computed by


$$R_{-1}(V) \stackrel{\text{def}}{=} \text{Rift}_R(V) \cong \int_{b \in B} \text{Hom}_{\{t, f\}}(R_{-1}^b, V_{-2}^b),$$

where we have used **Proposition 6.2.4.1**.

PROOF 6.4.2.3 ► PROOF OF REMARK 6.4.2.2

We have

$$\begin{aligned}
 \text{Rift}_R(V) &\cong \int_{b \in B} \text{Hom}_{\{t,f\}}(R_{-1}^b, V_{-2}^b) \\
 &= \left\{ a \in A \mid \int_{b \in B} \text{Hom}_{\{t,f\}}(R_a^b, V_{\star}^b) = \text{true} \right\} \\
 &= \left\{ a \in A \mid \begin{array}{l} \text{for each } b \in B, \text{ at least one of the} \\ \text{following conditions hold:} \\ \begin{array}{l} 1. \text{ We have } R_a^b = \text{false} \\ 2. \text{ The following conditions hold:} \\ \quad \text{(a) We have } R_a^b = \text{true} \\ \quad \text{(b) We have } V_{\star}^b = \text{true} \end{array} \end{array} \right\} \\
 &= \left\{ a \in A \mid \begin{array}{l} \text{for each } b \in B, \text{ at least one of the} \\ \text{following conditions hold:} \\ \begin{array}{l} 1. \text{ We have } b \notin R(a) \\ 2. \text{ The following conditions hold:} \\ \quad \text{(a) We have } b \in R(a) \\ \quad \text{(b) We have } b \in V \end{array} \end{array} \right\} \\
 &= \{ a \in A \mid \text{for each } b \in R(a), \text{ we have } b \in V \} \\
 &= \{ a \in A \mid R(a) \subset V \} \\
 &\stackrel{\text{def}}{=} R_{-1}(V).
 \end{aligned}$$

This finishes the proof. 

00RT PROPOSITION 6.4.2.4 ► PROPERTIES OF STRONG INVERSE IMAGES

Let $R: A \rightarrow B$ be a relation.

00RU

1. *Functoriality.* The assignment $V \mapsto R_{-1}(V)$ defines a functor

$$R_{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

where

- *Action on Objects.* For each $V \in \mathcal{P}(B)$, we have

$$[R_{-1}](V) \stackrel{\text{def}}{=} R_{-1}(V).$$

00RV

- *Action on Morphisms.* For each $U, V \in \mathcal{P}(B)$:
 - If $U \subset V$, then $R_{-1}(U) \subset R_{-1}(V)$.

2. *Adjointness.* We have an adjunction

$$(R_* \dashv R_{-1}): \mathcal{P}(A) \begin{array}{c} \xrightarrow{R_*} \\ \perp \\ \xleftarrow{R_{-1}} \end{array} \mathcal{P}(B),$$

witnessed by a bijections of sets

$$\text{Hom}_{\mathcal{P}(A)}(R_*(U), V) \cong \text{Hom}_{\mathcal{P}(A)}(U, R_{-1}(V)),$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$, i.e. such that:

- (★) The following conditions are equivalent:
 - We have $R_*(U) \subset V$.
 - We have $U \subset R_{-1}(V)$.

00RW

3. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} R_{-1}(U_i) \subset R_{-1}\left(\bigcup_{i \in I} U_i\right),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} R_{-1}(U) \cup R_{-1}(V) &\subset R_{-1}(U \cup V), \\ \emptyset &\subset R_{-1}(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00RX

4. *Preservation of Limits.* We have an equality of sets

$$R_{-1}\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} R_{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have equalities

$$\begin{aligned} R_{-1}(U \cap V) &= R_{-1}(U) \cap R_{-1}(V), \\ R_{-1}(B) &= B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00RY

5. *Symmetric Lax Monoidality With Respect to Unions.* The direct image with compact support function of **Item 1** has a symmetric lax monoidal structure

$$\left(R_{-1}, R_{-1}^{\otimes}, R_{-1|\mathbb{1}}^{\otimes}\right): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with inclusions

$$\begin{aligned} R_{-1|U,V}^{\otimes}: R_{-1}(U) \cup R_{-1}(V) &\subset R_{-1}(U \cup V), \\ R_{-1|\mathbb{1}}^{\otimes}: \emptyset &\subset R_{-1}(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00RZ

6. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$\left(R_{-1}, R_{-1}^{\otimes}, R_{-1|\mathbb{1}}^{\otimes}\right): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with equalities

$$\begin{aligned} R_{-1|U,V}^{\otimes}: R_{-1}(U \cap V) &\xrightarrow{=} R_{-1}(U) \cap R_{-1}(V), \\ R_{-1|\mathbb{1}}^{\otimes}: R_{-1}(A) &\xrightarrow{=} B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00S0

7. *Interaction With Weak Inverse Images I.* We have

$$R_{-1}(V) = A \setminus R^{-1}(B \setminus V)$$

for each $V \in \mathcal{P}(B)$.

00S1

8. *Interaction With Weak Inverse Images II.* Let $R: A \dashrightarrow B$ be a relation from A to B .

00S2

(a) If R is a total relation, then we have an inclusion of sets

$$R_{-1}(V) \subset R^{-1}(V)$$

natural in $V \in \mathcal{P}(B)$.

00S3

(b) If R is total and functional, then the above inclusion is in fact an equality.

00S4

(c) Conversely, if we have $R_{-1} = R^{-1}$, then R is total and functional.

PROOF 6.4.2.5 ► PROOF OF PROPOSITION 6.4.2.4

Item 1: Functoriality

Clear.

Item 2: Adjointness

This follows from ?? of ??.

Item 3: Lax Preservation of Colimits

Omitted.

Item 4: Preservation of Limits

This follows from Item 2 and ?? of ??.

Item 5: Symmetric Lax Monoidality With Respect to Unions

This follows from Item 3.

Item 6: Symmetric Strict Monoidality With Respect to Intersections

This follows from Item 4.

Item 7: Interaction With Weak Inverse Images I

We claim we have an equality

$$R_{-1}(B \setminus V) = A \setminus R^{-1}(V).$$

Indeed, we have

$$\begin{aligned} R_{-1}(B \setminus V) &= \{a \in A \mid R(a) \subset B \setminus V\}, \\ A \setminus R^{-1}(V) &= \{a \in A \mid R(a) \cap V = \emptyset\}. \end{aligned}$$

Taking $V = B \setminus V$ then implies the original statement.

Item 8: Interaction With Weak Inverse Images II

Item 8a is clear, while Items 8b and 8c follow from Item 6 of Proposition 6.3.1.2.



PROPOSITION 6.4.2.6 ► PROPERTIES OF THE STRONG INVERSE IMAGE FUNCTION OPERATION

00S5

Let $R: A \rightarrow B$ be a relation.

00S6

1. *Functionality I*. The assignment $R \mapsto R_{-1}$ defines a function

$$(-)_{-1}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

00S7

2. *Functionality II.* The assignment $R \mapsto R_{-1}$ defines a function

$$(-)_{-1}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

00S8

3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$(\text{id}_A)_{-1} = \text{id}_{\mathcal{P}(A)}.$$

00S9

4. *Interaction With Composition.* For each pair of composable relations $R: A \rightarrow B$ and $S: B \rightarrow C$, we have

$$(S \circ R)_{-1} = R_{-1} \circ S_{-1},$$

$$\begin{array}{ccc} \mathcal{P}(C) & \xrightarrow{S_{-1}} & \mathcal{P}(B) \\ & \searrow (S \circ R)_{-1} & \downarrow R_{-1} \\ & & \mathcal{P}(A). \end{array}$$

PROOF 6.4.2.7 ► PROOF OF PROPOSITION 6.4.2.6

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

Indeed, we have

$$\begin{aligned} (\chi_A)_{-1}(U) &\stackrel{\text{def}}{=} \{a \in A \mid \chi_A(a) \subset U\} \\ &\stackrel{\text{def}}{=} \{a \in A \mid \{a\} \subset U\} \\ &= U \end{aligned}$$

for each $U \in \mathcal{P}(A)$. Thus $(\chi_A)_{-1} = \text{id}_{\mathcal{P}(A)}$.


Item 4: Interaction With Composition

Indeed, we have

$$\begin{aligned}
 (S \diamond R)_{-1}(U) &\stackrel{\text{def}}{=} \{a \in A \mid [S \diamond R](a) \subset U\} \\
 &\stackrel{\text{def}}{=} \{a \in A \mid S(R(a)) \subset U\} \\
 &\stackrel{\text{def}}{=} \{a \in A \mid S_*(R(a)) \subset U\} \\
 &= \{a \in A \mid R(a) \subset S_{-1}(U)\} \\
 &\stackrel{\text{def}}{=} R_{-1}(S_{-1}(U)) \\
 &\stackrel{\text{def}}{=} [R_{-1} \circ S_{-1}](U)
 \end{aligned}$$

for each $U \in \mathcal{P}(C)$, where we used [Item 2 of Proposition 6.4.2.4](#), which implies that the conditions

- We have $S_*(R(a)) \subset U$.
- We have $R(a) \subset S_{-1}(U)$.

are equivalent. Thus $(S \diamond R)_{-1} = R_{-1} \circ S_{-1}$. 

00SA 6.4.3 Weak Inverse Images

Let A and B be sets and let $R: A \rightarrow B$ be a relation.

00SB DEFINITION 6.4.3.1 ► WEAK INVERSE IMAGES

The **weak inverse image function associated to R^1** is the function

$$R^{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by²

$$R^{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid R(a) \cap V \neq \emptyset\}$$

for each $V \in \mathcal{P}(B)$.

¹*Further Terminology:* Also called simply the **inverse image function associated to R** .

²*Further Terminology:* The set $R^{-1}(V)$ is called the **weak inverse image of V by R** or simply the **inverse image of V by R** .

00SC REMARK 6.4.3.2 ► UNWINDING DEFINITION 6.4.3.1

Identifying subsets of B with relations from B to pt via [Item 3 of Proposition 2.4.3.9](#), we see that the weak inverse image function associated to R is

equivalently the function

$$R^{-1}: \underbrace{\mathcal{P}(B)}_{\cong \text{Rel}(B, \text{pt})} \rightarrow \underbrace{\mathcal{P}(A)}_{\cong \text{Rel}(A, \text{pt})}$$

defined by

$$R^{-1}(V) \stackrel{\text{def}}{=} V \diamond R$$

for each $V \in \mathcal{P}(A)$, where $R \diamond V$ is the composition

$$A \xrightarrow{R} B \xrightarrow{V} \text{pt}.$$


Explicitly, we have

$$\begin{aligned} R^{-1}(V) &\stackrel{\text{def}}{=} V \diamond R \\ &\stackrel{\text{def}}{=} \int^{b \in B} V_b^{-1} \times R_{-2}^b. \end{aligned}$$

PROOF 6.4.3.3 ► PROOF OF REMARK 6.4.3.2

We have

$$\begin{aligned}
 V \diamond R &\stackrel{\text{def}}{=} \int^{b \in B} V_b^{-1} \times R_{-2}^b \\
 &= \left\{ a \in A \mid \int^{b \in B} V_b^* \times R_a^b = \text{true} \right\} \\
 &= \left\{ a \in A \mid \begin{array}{l} \text{there exists } b \in B \text{ such that the} \\ \text{following conditions hold:} \\ 1. \text{ We have } V_b^* = \text{true} \\ 2. \text{ We have } R_a^b = \text{true} \end{array} \right\} \\
 &= \left\{ a \in A \mid \begin{array}{l} \text{there exists } b \in B \text{ such that the} \\ \text{following conditions hold:} \\ 1. \text{ We have } b \in V \\ 2. \text{ We have } b \in R(a) \end{array} \right\} \\
 &= \{ a \in A \mid \text{there exists } b \in V \text{ such that } b \in R(a) \} \\
 &= \{ a \in A \mid R(a) \cap V \neq \emptyset \} \\
 &\stackrel{\text{def}}{=} R^{-1}(V)
 \end{aligned}$$

This finishes the proof. 

00SD

PROPOSITION 6.4.3.4 ► PROPERTIES OF WEAK INVERSE IMAGE FUNCTIONS

Let $R: A \rightarrow B$ be a relation.

00SE

1. *Functoriality.* The assignment $V \mapsto R^{-1}(V)$ defines a functor

$$R^{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

where

- *Action on Objects.* For each $V \in \mathcal{P}(B)$, we have

$$[R^{-1}](V) \stackrel{\text{def}}{=} R^{-1}(V).$$

- *Action on Morphisms.* For each $U, V \in \mathcal{P}(B)$:

– If $U \subset V$, then $R^{-1}(U) \subset R^{-1}(V)$.

00SF

2. *Adjointness.* We have an adjunction

$$(R^{-1} \dashv R_!) : \mathcal{P}(B) \begin{array}{c} \xrightarrow{R^{-1}} \\ \perp \\ \xleftarrow{R_!} \end{array} \mathcal{P}(A),$$

witnessed by a bijections of sets

$$\mathrm{Hom}_{\mathcal{P}(A)}(R^{-1}(U), V) \cong \mathrm{Hom}_{\mathcal{P}(A)}(U, R_!(V)),$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$, i.e. such that:

(★) The following conditions are equivalent:

- We have $R^{-1}(U) \subset V$.
- We have $U \subset R_!(V)$.

00SG

3. *Preservation of Colimits.* We have an equality of sets

$$R^{-1}\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} R^{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have equalities

$$\begin{aligned} R^{-1}(U) \cup R^{-1}(V) &= R^{-1}(U \cup V), \\ R^{-1}(\emptyset) &= \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00SH

4. *Oplax Preservation of Limits.* We have an inclusion of sets

$$R^{-1}\left(\bigcap_{i \in I} U_i\right) \subset \bigcap_{i \in I} R^{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} R^{-1}(U \cap V) &\subset R^{-1}(U) \cap R^{-1}(V), \\ R^{-1}(A) &\subset B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00SJ

5. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$\left(R^{-1}, R^{-1, \otimes}, R_{\perp}^{-1, \otimes}\right): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} R_{U, V}^{-1, \otimes}: R^{-1}(U) \cup R^{-1}(V) &\xrightarrow{=} R^{-1}(U \cup V), \\ R_{\perp}^{-1, \otimes}: \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00SK

6. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric oplax monoidal structure

$$\left(R^{-1}, R^{-1, \otimes}, R_{\perp}^{-1, \otimes}\right): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with inclusions

$$\begin{aligned} R_{U, V}^{-1, \otimes}: R^{-1}(U \cap V) &\subset R^{-1}(U) \cap R^{-1}(V), \\ R_{\perp}^{-1, \otimes}: R^{-1}(A) &\subset B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00SL

7. *Interaction With Strong Inverse Images I.* We have

$$R^{-1}(V) = A \setminus R_{-1}(B \setminus V)$$

for each $V \in \mathcal{P}(B)$.

00SM

8. *Interaction With Strong Inverse Images II.* Let $R: A \rightarrow B$ be a relation from A to B .

00SN

(a) If R is a total relation, then we have an inclusion of sets

$$R_{-1}(V) \subset R^{-1}(V)$$

natural in $V \in \mathcal{P}(B)$.

00SP

(b) If R is total and functional, then the above inclusion is in fact an equality.

00SQ

(c) Conversely, if we have $R_{-1} = R^{-1}$, then R is total and functional.

PROOF 6.4.3.5 ► PROOF OF PROPOSITION 6.4.3.4

Item 1: Functoriality

Clear.

Item 2: Adjointness

This follows from ?? of ??.

Item 3: Preservation of Colimits

This follows from Item 2 and ?? of ??.

Item 4: Oplax Preservation of Limits

Omitted.

Item 5: Symmetric Strict Monoidality With Respect to Unions

This follows from Item 3.

Item 6: Symmetric Oplax Monoidality With Respect to Intersections

This follows from Item 4.

Item 7: Interaction With Strong Inverse Images I

This follows from Item 7 of Proposition 6.4.2.4.

Item 8: Interaction With Strong Inverse Images II

This was proved in Item 8 of Proposition 6.4.2.4. 

PROPOSITION 6.4.3.6 ► PROPERTIES OF THE WEAK INVERSE IMAGE FUNCTION OPERATION

00SR

Let $R: A \dashrightarrow B$ be a relation.

00SS

1. *Functionality I.* The assignment $R \mapsto R^{-1}$ defines a function

$$(-)^{-1}: \text{Rel}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

00ST

2. *Functionality II.* The assignment $R \mapsto R^{-1}$ defines a function

$$(-)^{-1}: \text{Rel}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

00SU

3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have¹

$$(\chi_A)^{-1} = \text{id}_{\mathcal{P}(A)}.$$

00SV

4. *Interaction With Composition.* For each pair of composable relations $R: A \rightarrow B$ and $S: B \rightarrow C$, we have²

$$(S \circ R)^{-1} = R^{-1} \circ S^{-1},$$

$$\begin{array}{ccc} \mathcal{P}(C) & \xrightarrow{S^{-1}} & \mathcal{P}(B) \\ & \searrow (S \circ R)^{-1} & \downarrow R^{-1} \\ & & \mathcal{P}(A). \end{array}$$

¹That is, the postcomposition

$$(\chi_A)^{-1}: \text{Rel}(\text{pt}, A) \rightarrow \text{Rel}(\text{pt}, A)$$

is equal to $\text{id}_{\text{Rel}(\text{pt}, A)}$.

²That is, we have

$$(S \circ R)^{-1} = R^{-1} \circ S^{-1},$$

$$\begin{array}{ccc} \text{Rel}(\text{pt}, C) & \xrightarrow{R^{-1}} & \text{Rel}(\text{pt}, B) \\ & \searrow (S \circ R)^{-1} & \downarrow S^{-1} \\ & & \text{Rel}(\text{pt}, A). \end{array}$$

PROOF 6.4.3.7 ► PROOF OF PROPOSITION 6.4.3.6

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

This follows from **Item 5** of **Proposition 8.1.6.2**.

Item 4: Interaction With Composition

This follows from **Item 2** of **Proposition 8.1.6.2**. 

00SW 6.4.4 Direct Images With Compact Support

Let A and B be sets and let $R: A \rightarrow B$ be a relation.

00SX

DEFINITION 6.4.4.1 ► DIRECT IMAGES WITH COMPACT SUPPORT

The **direct image with compact support function associated to R** is the function

$$R_! : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by^{1,2}

$$\begin{aligned} R_!(U) &\stackrel{\text{def}}{=} \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ if we have} \\ b \in R(a), \text{ then } a \in U \end{array} \right\} \\ &= \{ b \in B \mid R^{-1}(b) \subset U \} \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

¹Further Terminology: The set $R_!(U)$ is called the **direct image with compact support of U by R** .

²We also have

$$R_!(U) = B \setminus R_*(A \setminus U);$$

see Item 7 of Proposition 6.4.4.4.

00SY

REMARK 6.4.4.2 ► UNWINDING DEFINITION 6.4.4.1

Identifying subsets of B with relations from pt to B via Item 3 of Proposition 2.4.3.9, we see that the direct image with compact support function associated to R is equivalently the function

$$R_! : \underbrace{\mathcal{P}(A)}_{\cong \text{Rel}(A, \text{pt})} \rightarrow \underbrace{\mathcal{P}(B)}_{\cong \text{Rel}(B, \text{pt})}$$

defined by

$$R_!(U) \stackrel{\text{def}}{=} \text{Ran}_R(U),$$

being explicitly computed by


$$\begin{aligned} R^*(U) &\stackrel{\text{def}}{=} \text{Ran}_R(U) \\ &\cong \int_{a \in A} \text{Hom}_{\{\text{t}, \text{f}\}}(R_a^{-2}, U_a^{-1}), \end{aligned}$$

where we have used Proposition 6.2.3.1.

PROOF 6.4.4.3 ► PROOF OF REMARK 6.4.4.2

We have

$$\begin{aligned}
 \text{Ran}_R(V) &\cong \int_{a \in A} \text{Hom}_{\{t,f\}}(R_a^{-2}, U_a^{-1}) \\
 &= \left\{ b \in B \mid \int_{a \in A} \text{Hom}_{\{t,f\}}(R_a^b, U_a^*) = \text{true} \right\} \\
 &= \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ at least one of the} \\ \text{following conditions hold:} \\ \quad 1. \text{ We have } R_a^b = \text{false} \\ \quad 2. \text{ The following conditions hold:} \\ \qquad (a) \text{ We have } R_a^b = \text{true} \\ \qquad (b) \text{ We have } U_a^* = \text{true} \end{array} \right\} \\
 &= \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ at least one of the} \\ \text{following conditions hold:} \\ \quad 1. \text{ We have } b \notin R(A) \\ \quad 2. \text{ The following conditions hold:} \\ \qquad (a) \text{ We have } b \in R(a) \\ \qquad (b) \text{ We have } a \in U \end{array} \right\} \\
 &= \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ if we have} \\ b \in R(a), \text{ then } a \in U \end{array} \right\} \\
 &= \{ b \in B \mid R^{-1}(b) \subset U \} \\
 &\stackrel{\text{def}}{=} R^{-1}(U).
 \end{aligned}$$

This finishes the proof. 

00SZ

PROPOSITION 6.4.4.4 ► PROPERTIES OF DIRECT IMAGES WITH COMPACT SUPPORT

Let $R: A \rightarrow B$ be a relation.

00T0

1. *Functoriality.* The assignment $U \mapsto R_!(U)$ defines a functor

$$R_!: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

where

· *Action on Objects.* For each $U \in \mathcal{P}(A)$, we have

$$[R_!](U) \stackrel{\text{def}}{=} R_!(U).$$

· *Action on Morphisms.* For each $U, V \in \mathcal{P}(A)$:

– If $U \subset V$, then $R_!(U) \subset R_!(V)$.

00T1

2. *Adjointness.* We have an adjunction

$$(R^{-1} \dashv R_!): \mathcal{P}(B) \begin{array}{c} \xrightarrow{R^{-1}} \\ \perp \\ \xleftarrow{R_!} \end{array} \mathcal{P}(A),$$

witnessed by a bijections of sets

$$\text{Hom}_{\mathcal{P}(A)}(R^{-1}(U), V) \cong \text{Hom}_{\mathcal{P}(A)}(U, R_!(V)),$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$, i.e. such that:

(★) The following conditions are equivalent:

- We have $R^{-1}(U) \subset V$.
- We have $U \subset R_!(V)$.

00T2

3. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} R_!(U_i) \subset R_!\left(\bigcup_{i \in I} U_i\right),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} R_!(U) \cup R_!(V) &\subset R_!(U \cup V), \\ \emptyset &\subset R_!(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00T3

4. *Preservation of Limits.* We have an equality of sets

$$R_!\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} R_!(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have equalities

$$\begin{aligned} R_!(U \cap V) &= R_!(U) \cap R_!(V), \\ R_!(A) &= B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00T4

5. *Symmetric Lax Monoidality With Respect to Unions.* The direct image with compact support function of **Item 1** has a symmetric lax monoidal structure

$$(R_!, R_!^\otimes, R_{!1}^\otimes): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with inclusions

$$R_{!U,V}^\otimes: R_!(U) \cup R_!(V) \subset R_!(U \cup V),$$

$$R_{!1}^\otimes: \emptyset \subset R_!(\emptyset),$$

natural in $U, V \in \mathcal{P}(A)$.

00T5

6. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$(R_!, R_!^\otimes, R_{!1}^\otimes): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with equalities

$$R_{!U,V}^\otimes: R_!(U \cap V) \xrightarrow{=} R_!(U) \cap R_!(V),$$

$$R_{!1}^\otimes: R_!(A) \xrightarrow{=} B,$$

natural in $U, V \in \mathcal{P}(A)$.

00T6

7. *Relation to Direct Images.* We have

$$R_!(U) = B \setminus R_*(A \setminus U)$$

for each $U \in \mathcal{P}(A)$.

PROOF 6.4.4.5 ► PROOF OF PROPOSITION 6.4.4.4

Item 1: Functoriality

Clear.

Item 2: Adjointness

This follows from ?? of ??.

Item 3: Lax Preservation of Colimits

Omitted.

Item 4: Preservation of Limits

This follows from [Item 2](#) and ?? of ??.

Item 5: Symmetric Lax Monoidality With Respect to Unions

This follows from [Item 3](#).

Item 6: Symmetric Strict Monoidality With Respect to Intersections

This follows from [Item 4](#).

Item 7: Relation to Direct Images

This follows from [Item 7](#) of [Proposition 6.4.1.3](#). Alternatively, we may prove it directly as follows, with the proof proceeding in the same way as in the case of functions ([Item 9](#) of [Proposition 2.4.6.6](#)).

We claim that $R_!(U) = B \setminus R_*(A \setminus U)$:

- *The First Implication.* We claim that

$$R_!(U) \subset B \setminus R_*(A \setminus U).$$

Let $b \in R_!(U)$. We need to show that $b \notin R_*(A \setminus U)$, i.e. that there is no $a \in A \setminus U$ such that $b \in R(a)$.

This is indeed the case, as otherwise we would have $a \in R^{-1}(b)$ and $a \notin U$, contradicting $R^{-1}(b) \subset U$ (which holds since $b \in R_!(U)$).

Thus $b \in B \setminus R_*(A \setminus U)$.


- *The Second Implication.* We claim that

$$B \setminus R_*(A \setminus U) \subset R_!(U).$$

Let $b \in B \setminus R_*(A \setminus U)$. We need to show that $b \in R_!(U)$, i.e. that $R^{-1}(b) \subset U$.

Since $b \notin R_*(A \setminus U)$, there exists no $a \in A \setminus U$ such that $b \in R(a)$, and hence $R^{-1}(b) \subset U$.

Thus $b \in R_!(U)$.

This finishes the proof. 

PROPOSITION 6.4.4.6 ► PROPERTIES OF THE DIRECT IMAGE WITH COMPACT SUPPORT FUNCTION OPERATION

00T7

Let $R: A \rightarrow B$ be a relation.

00T8

1. *Functionality I.* The assignment $R \mapsto R_!$ defines a function

$$(-)_!: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

00T9

2. *Functionality II.* The assignment $R \mapsto R_!$ defines a function

$$(-)_!: \text{Sets}(A, B) \rightarrow \text{Hom}_{\text{Pos}}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

00TA

3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$(\text{id}_A)_! = \text{id}_{\mathcal{P}(A)}.$$

00TB

4. *Interaction With Composition.* For each pair of composable relations $R: A \rightarrow B$ and $S: B \rightarrow C$, we have

$$(S \diamond R)_! = S_! \circ R_!, \quad \begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{R_!} & \mathcal{P}(B) \\ & \searrow (S \diamond R)_! & \downarrow S_! \\ & & \mathcal{P}(C). \end{array}$$

PROOF 6.4.4.7 ► PROOF OF PROPOSITION 6.4.4.6

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

Indeed, we have

$$\begin{aligned} (\chi_A)_!(U) &\stackrel{\text{def}}{=} \{a \in A \mid \chi_A^{-1}(a) \subset U\} \\ &\stackrel{\text{def}}{=} \{a \in A \mid \{a\} \subset U\} \\ &= U \end{aligned}$$

for each $U \in \mathcal{P}(A)$. Thus $(\chi_A)_! = \text{id}_{\mathcal{P}(A)}$.


Item 4: Interaction With Composition

Indeed, we have

$$\begin{aligned}
 (S \diamond R)_! (U) &\stackrel{\text{def}}{=} \{c \in C \mid [S \diamond R]^{-1}(c) \subset U\} \\
 &\stackrel{\text{def}}{=} \{c \in C \mid S^{-1}(R^{-1}(c)) \subset U\} \\
 &= \{c \in C \mid R^{-1}(c) \subset S_!(U)\} \\
 &\stackrel{\text{def}}{=} R_!(S_!(U)) \\
 &\stackrel{\text{def}}{=} [R_! \circ S_!](U)
 \end{aligned}$$

for each $U \in \mathcal{P}(C)$, where we used [Item 2](#) of [Proposition 6.4.4.4](#), which implies that the conditions

- We have $S^{-1}(R^{-1}(c)) \subset U$.
- We have $R^{-1}(c) \subset S_!(U)$.

are equivalent. Thus $(S \diamond R)_! = S_! \circ R_!$. 

00TC 6.4.5 Functoriality of Powersets

00TD PROPOSITION 6.4.5.1 ► FUNCTORIALITY OF POWERSETS I

The assignment $X \mapsto \mathcal{P}(X)$ defines functors¹

$$\begin{aligned}
 \mathcal{P}_* &: \text{Rel} \rightarrow \text{Sets}, \\
 \mathcal{P}_{-1} &: \text{Rel}^{\text{op}} \rightarrow \text{Sets}, \\
 \mathcal{P}^{-1} &: \text{Rel}^{\text{op}} \rightarrow \text{Sets}, \\
 \mathcal{P}_! &: \text{Rel} \rightarrow \text{Sets}
 \end{aligned}$$

where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Rel})$, we have

$$\begin{aligned}
 \mathcal{P}_*(A) &\stackrel{\text{def}}{=} \mathcal{P}(A), \\
 \mathcal{P}_{-1}(A) &\stackrel{\text{def}}{=} \mathcal{P}(A), \\
 \mathcal{P}^{-1}(A) &\stackrel{\text{def}}{=} \mathcal{P}(A), \\
 \mathcal{P}_!(A) &\stackrel{\text{def}}{=} \mathcal{P}(A).
 \end{aligned}$$

· *Action on Morphisms.* For each morphism $R: A \rightarrow B$ of Rel, the images

$$\begin{aligned}\mathcal{P}_*(R) &: \mathcal{P}(A) \rightarrow \mathcal{P}(B), \\ \mathcal{P}_{-1}(R) &: \mathcal{P}(B) \rightarrow \mathcal{P}(A), \\ \mathcal{P}^{-1}(R) &: \mathcal{P}(B) \rightarrow \mathcal{P}(A), \\ \mathcal{P}_! (R) &: \mathcal{P}(A) \rightarrow \mathcal{P}(B)\end{aligned}$$

of R by \mathcal{P}_* , \mathcal{P}_{-1} , \mathcal{P}^{-1} , and $\mathcal{P}_!$ are defined by

$$\begin{aligned}\mathcal{P}_*(R) &\stackrel{\text{def}}{=} R_*, \\ \mathcal{P}_{-1}(R) &\stackrel{\text{def}}{=} R_{-1}, \\ \mathcal{P}^{-1}(R) &\stackrel{\text{def}}{=} R^{-1}, \\ \mathcal{P}_!(R) &\stackrel{\text{def}}{=} R_!\end{aligned}$$

as in [Definitions 6.4.1.1](#), [6.4.2.1](#), [6.4.3.1](#) and [6.4.4.1](#).

¹The functor $\mathcal{P}_*: \text{Rel} \rightarrow \text{Sets}$ admits a left adjoint; see [Item 3](#) of [Proposition 6.3.1.2](#).

PROOF 6.4.5.2 ► PROOF OF PROPOSITION 6.4.5.1

This follows from [Items 3](#) and [4](#) of [Proposition 6.4.1.5](#), [Items 3](#) and [4](#) of [Proposition 6.4.2.6](#), [Items 3](#) and [4](#) of [Proposition 6.4.3.6](#), and [Items 3](#) and [4](#) of [Proposition 6.4.4.6](#). ▢

00TE 6.4.6 Functoriality of Powersets: Relations on Powersets

Let A and B be sets and let $R: A \rightarrow B$ be a relation.

00TF DEFINITION 6.4.6.1 ► THE RELATION ON POWERSETS ASSOCIATED TO A RELATION

The **relation on powersets associated to** R is the relation


$$\mathcal{P}(R): \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by¹

$$\mathcal{P}(R)_U^V \stackrel{\text{def}}{=} \mathbf{Rel}(\chi_{\text{pt}}, V \diamond R \diamond U)$$

for each $U \in \mathcal{P}(A)$ and each $V \in \mathcal{P}(B)$.

¹Illustration:



00TG

REMARK 6.4.6.2 ► UNWINDING DEFINITION 6.4.6.1

In detail, we have $U \sim_{\mathcal{P}(R)} V$ iff the following equivalent conditions hold:

- We have $\chi_{\text{pt}} \subset V \diamond R \diamond U$.
- We have $(V \diamond R \diamond U)_{\star}^{\star} = \text{true}$, i.e. we have

$$\int^{a \in A} \int^{b \in B} V_b^{\star} \times R_a^b \times U_{\star}^a = \text{true}.$$

- There exists some $a \in A$ and some $b \in B$ such that:
 - We have $U_{\star}^a = \text{true}$.
 - We have $R_a^b = \text{true}$.
 - We have $V_b^{\star} = \text{true}$.
- There exists some $a \in A$ and some $b \in B$ such that:
 - We have $a \in U$.
 - We have $a \sim_R b$.
 - We have $b \in V$.

00TH

PROPOSITION 6.4.6.3 ► FUNCTORIALITY OF POWERSETS II

The assignment $R \mapsto \mathcal{P}(R)$ defines a functor

$$\mathcal{P}: \text{Rel} \rightarrow \text{Rel}.$$

PROOF 6.4.6.4 ► PROOF OF PROPOSITION 6.4.6.3

Omitted. 

Appendices

6.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets

Relations

5. Relations

6. Constructions With Relations

7. Equivalence Relations and Apartness Relations

Category Theory

8. Categories

Bicategories

9. Types of Morphisms in Bicategories

Chapter 7

Equivalence Relations and Apartness Relations

00TJ This chapter contains some material about reflexive, symmetric, transitive, equivalence, and apartness relations.

Contents

7.1	Reflexive Relations	395
7.1.1	Foundations.....	395
7.1.2	The Reflexive Closure of a Relation.....	397
7.2	Symmetric Relations	399
7.2.1	Foundations.....	399
7.2.2	The Symmetric Closure of a Relation.....	399
7.3	Transitive Relations	401
7.3.1	Foundations.....	401
7.3.2	The Transitive Closure of a Relation.....	403
7.4	Equivalence Relations	406
7.4.1	Foundations.....	406
7.4.2	The Equivalence Closure of a Relation.....	407
7.5	Quotients by Equivalence Relations	408
7.5.1	Equivalence Classes.....	408
7.5.2	Quotients of Sets by Equivalence Relations.....	409
7.A	Other Chapters	414

00TK 7.1 Reflexive Relations

00TL 7.1.1 Foundations

Let A be a set.

00TM DEFINITION 7.1.1.1 ► REFLEXIVE RELATIONS

A **reflexive relation** is equivalently:¹

- An \mathbb{E}_0 -monoid in $(\mathbf{N}_\bullet(\mathbf{Rel}(A, A)), \chi_A)$.
- A pointed object in $(\mathbf{Rel}(A, A), \chi_A)$.

¹Note that since $\mathbf{Rel}(A, A)$ is posetal, reflexivity is a property of a relation, rather than extra structure.

00TN REMARK 7.1.1.2 ► UNWINDING DEFINITION 7.1.1.1

In detail, a relation R on A is **reflexive** if we have an inclusion

$$\eta_R: \chi_A \subset R$$

of relations in $\mathbf{Rel}(A, A)$, i.e. if, for each $a \in A$, we have $a \sim_R a$.

00TP DEFINITION 7.1.1.3 ► THE PO/SET OF REFLEXIVE RELATIONS ON A SET

Let A be a set.

1. The **set of reflexive relations on A** is the subset $\mathbf{Rel}^{\text{refl}}(A, A)$ of $\mathbf{Rel}(A, A)$ spanned by the reflexive relations.
2. The **poset of relations on A** is the subposet $\mathbf{Rel}^{\text{refl}}(A, A)$ of $\mathbf{Rel}(A, A)$ spanned by the reflexive relations.

00TS PROPOSITION 7.1.1.4 ► PROPERTIES OF REFLEXIVE RELATIONS

Let R and S be relations on A .

1. *Interaction With Inverses.* If R is reflexive, then so is R^\dagger .
2. *Interaction With Composition.* If R and S are reflexive, then so is $S \diamond R$.

PROOF 7.1.1.5 ► PROOF OF PROPOSITION 7.1.1.4

Item 1: Interaction With Inverses

Clear.

Item 2: Interaction With Composition

Clear.



00TV **7.1.2 The Reflexive Closure of a Relation**

Let R be a relation on A .

00TW **DEFINITION 7.1.2.1 ► THE REFLEXIVE CLOSURE OF A RELATION**

The **reflexive closure** of \sim_R is the relation \sim_R^{refl} satisfying the following universal property:²

- (★) Given another reflexive relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{refl}} \subset \sim_S$.

¹Further Notation: Also written R^{refl} .

²Slogan: The reflexive closure of R is the smallest reflexive relation containing R .

00TX **CONSTRUCTION 7.1.2.2 ► THE REFLEXIVE CLOSURE OF A RELATION**

Concretely, \sim_R^{refl} is the free pointed object on R in $(\mathbf{Rel}(A, A), \chi_A)$ ¹, being given by

$$\begin{aligned} R^{\text{refl}} &\stackrel{\text{def}}{=} R \amalg^{\mathbf{Rel}(A, A)} \Delta_A \\ &= R \cup \Delta_A \\ &= \{(a, b) \in A \times A \mid \text{we have } a \sim_R b \text{ or } a = b\}. \end{aligned}$$

¹Or, equivalently, the free \mathbb{E}_0 -monoid on R in $(\mathbf{N}_\bullet(\mathbf{Rel}(A, A)), \chi_A)$.

PROOF 7.1.2.3 ► PROOF OF CONSTRUCTION 7.1.2.2

Clear. 

00TY **PROPOSITION 7.1.2.4 ► PROPERTIES OF THE REFLEXIVE CLOSURE OF A RELATION**

Let R be a relation on A .

- 00TZ 1. *Adjointness.* We have an adjunction

$$\left((-)^{\text{refl}} \dashv \overset{\circlearrowleft}{\text{忘}} \right): \mathbf{Rel}(A, A) \begin{matrix} \xrightarrow{(-)^{\text{refl}}} \\ \perp \\ \xleftarrow{\overset{\circlearrowleft}{\text{忘}}} \end{matrix} \mathbf{Rel}^{\text{refl}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{refl}}(R^{\text{refl}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{refl}}(A, A))$ and $S \in \text{Obj}(\mathbf{Rel}(A, A))$.

00U0 2. *The Reflexive Closure of a Reflexive Relation.* If R is reflexive, then $R^{\text{refl}} = R$.

00U1 3. *Idempotency.* We have

$$\left(R^{\text{refl}}\right)^{\text{refl}} = R^{\text{refl}}.$$

00U2 4. *Interaction With Inverses.* We have

$$\left(R^\dagger\right)^{\text{refl}} = \left(R^{\text{refl}}\right)^\dagger, \quad \begin{array}{ccc} \text{Rel}(A, A) & \xrightarrow{(-)^{\text{refl}}} & \text{Rel}(A, A) \\ (-)^\dagger \downarrow & & \downarrow (-)^\dagger \\ \text{Rel}(A, A) & \xrightarrow{(-)^{\text{refl}}} & \text{Rel}(A, A). \end{array}$$

00U3 5. *Interaction With Composition.* We have

$$\left(S \diamond R\right)^{\text{refl}} = S^{\text{refl}} \diamond R^{\text{refl}}, \quad \begin{array}{ccc} \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A) \\ (-)^{\text{refl}} \times (-)^{\text{refl}} \downarrow & & \downarrow (-)^{\text{refl}} \\ \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A). \end{array}$$

PROOF 7.1.2.5 ► PROOF OF PROPOSITION 7.1.2.4

Item 1: Adjointness

This is a rephrasing of the universal property of the reflexive closure of a relation, stated in [Definition 7.1.2.1](#).

Item 2: The Reflexive Closure of a Reflexive Relation

Clear.

Item 3: Idempotency

This follows from [Item 2](#).

Item 4: Interaction With Inverses

Clear.

Item 5: Interaction With Composition

This follows from [Item 2](#) of [Proposition 7.1.1.4](#). 

00U4 7.2 Symmetric Relations

00U5 7.2.1 Foundations

Let A be a set.

00U6 DEFINITION 7.2.1.1 ► SYMMETRIC RELATIONS

A relation R on A is **symmetric** if we have $R^\dagger = R$.

00U7 REMARK 7.2.1.2 ► UNWINDING DEFINITION 7.2.1.1

In detail, a relation R is symmetric if it satisfies the following condition:

(★) For each $a, b \in A$, if $a \sim_R b$, then $b \sim_R a$.

00U8 DEFINITION 7.2.1.3 ► THE PO/SET OF SYMMETRIC RELATIONS ON A SET

Let A be a set.

- 00U9 1. The **set of symmetric relations on A** is the subset $\text{Rel}^{\text{symm}}(A, A)$ of $\text{Rel}(A, A)$ spanned by the symmetric relations.
- 00UA 2. The **poset of relations on A** is the subposet $\mathbf{Rel}^{\text{symm}}(A, A)$ of $\mathbf{Rel}(A, A)$ spanned by the symmetric relations.

00UB PROPOSITION 7.2.1.4 ► PROPERTIES OF SYMMETRIC RELATIONS

Let R and S be relations on A .

- 00UC 1. *Interaction With Inverses.* If R is symmetric, then so is R^\dagger .
- 00UD 2. *Interaction With Composition.* If R and S are symmetric, then so is $S \diamond R$.

PROOF 7.2.1.5 ► PROOF OF PROPOSITION 7.2.1.4

Item 1: Interaction With Inverses

Clear.

Item 2: Interaction With Composition

Clear. 

00UE 7.2.2 The Symmetric Closure of a Relation

Let R be a relation on A .

00UF

DEFINITION 7.2.2.1 ► THE SYMMETRIC CLOSURE OF A RELATION

The **symmetric closure** of \sim_R is the relation \sim_R^{symm} ¹ satisfying the following universal property:²

- (★) Given another symmetric relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{symm}} \subset \sim_S$.

¹Further Notation: Also written R^{symm} .

²Slogan: The symmetric closure of R is the smallest symmetric relation containing R .

00UG

CONSTRUCTION 7.2.2.2 ► THE SYMMETRIC CLOSURE OF A RELATION

Concretely, \sim_R^{symm} is the symmetric relation on A defined by

$$\begin{aligned} R^{\text{symm}} &\stackrel{\text{def}}{=} R \cup R^\dagger \\ &= \{(a, b) \in A \times A \mid \text{we have } a \sim_R b \text{ or } b \sim_R a\}. \end{aligned}$$

PROOF 7.2.2.3 ► PROOF OF CONSTRUCTION 7.2.2.2

Clear. 

00UH

PROPOSITION 7.2.2.4 ► PROPERTIES OF THE SYMMETRIC CLOSURE OF A RELATION

Let R be a relation on A .

00UJ

1. *Adjointness.* We have an adjunction

$$\left((-)^{\text{symm}} \dashv \overset{\circlearrowleft}{\sim} \right): \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{symm}}} \\ \perp \\ \xleftarrow{\overset{\circlearrowleft}{\sim}} \end{array} \mathbf{Rel}^{\text{symm}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{symm}}(R^{\text{symm}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{symm}}(A, A))$ and $S \in \text{Obj}(\mathbf{Rel}(A, A))$.

00UK

2. *The Symmetric Closure of a Symmetric Relation.* If R is symmetric, then $R^{\text{symm}} = R$.

00UL

3. *Idempotency.* We have

$$(R^{\text{symm}})^{\text{symm}} = R^{\text{symm}}.$$

00UM

4. *Interaction With Inverses.* We have

$$\begin{array}{ccc} (R^\dagger)^{\text{symm}} = (R^{\text{symm}})^\dagger, & \text{Rel}(A, A) \xrightarrow{(-)^{\text{symm}}} \text{Rel}(A, A) & \\ & \downarrow (-)^\dagger & \downarrow (-)^\dagger \\ & \text{Rel}(A, A) \xrightarrow{(-)^{\text{symm}}} \text{Rel}(A, A). & \end{array}$$

00UN

5. *Interaction With Composition.* We have

$$\begin{array}{ccc} (S \diamond R)^{\text{symm}} = S^{\text{symm}} \diamond R^{\text{symm}}, & \text{Rel}(A, A) \times \text{Rel}(A, A) \xrightarrow{\diamond} \text{Rel}(A, A) & \\ & \downarrow (-)^{\text{symm}} \times (-)^{\text{symm}} & \downarrow (-)^{\text{symm}} \\ & \text{Rel}(A, A) \times \text{Rel}(A, A) \xrightarrow{\diamond} \text{Rel}(A, A). & \end{array}$$

PROOF 7.2.2.5 ► PROOF OF PROPOSITION 7.2.2.4**Item 1: Adjointness**

This is a rephrasing of the universal property of the symmetric closure of a relation, stated in [Definition 7.2.2.1](#).

Item 2: The Symmetric Closure of a Symmetric Relation

Clear.

Item 3: Idempotency

This follows from [Item 2](#).

Item 4: Interaction With Inverses

Clear.

Item 5: Interaction With Composition

This follows from [Item 2](#) of [Proposition 7.2.1.4](#). 

00UP 7.3 Transitive Relations**00UQ 7.3.1 Foundations**

Let A be a set.

00UR

DEFINITION 7.3.1.1 ► TRANSITIVE RELATIONS

A **transitive relation** is equivalently:¹

- A non-unital \mathbb{E}_1 -monoid in $(\mathbf{N} \cdot (\mathbf{Rel}(A, A)), \diamond)$.
- A non-unital monoid in $(\mathbf{Rel}(A, A), \diamond)$.

¹Note that since $\mathbf{Rel}(A, A)$ is posetal, transitivity is a property of a relation, rather than extra structure.

00US

REMARK 7.3.1.2 ► UNWINDING DEFINITION 7.3.1.1

In detail, a relation R on A is **transitive** if we have an inclusion

$$\mu_R: R \diamond R \subset R$$

of relations in $\mathbf{Rel}(A, A)$, i.e. if, for each $a, c \in A$, the following condition is satisfied:

- (★) If there exists some $b \in A$ such that $a \sim_R b$ and $b \sim_R c$, then $a \sim_R c$.

00UT

DEFINITION 7.3.1.3 ► THE PO/SET OF TRANSITIVE RELATIONS ON A SET

Let A be a set.

00UU

1. The **set of transitive relations from A to B** is the subset $\mathbf{Rel}^{\text{trans}}(A)$ of $\mathbf{Rel}(A, A)$ spanned by the transitive relations.

00UV

2. The **poset of relations from A to B** is the subposet $\mathbf{Rel}^{\text{trans}}(A)$ of $\mathbf{Rel}(A, A)$ spanned by the transitive relations.

00UW

PROPOSITION 7.3.1.4 ► PROPERTIES OF TRANSITIVE RELATIONS

Let R and S be relations on A .

00UX

1. *Interaction With Inverses.* If R is transitive, then so is R^\dagger .

00UY

2. *Interaction With Composition.* If R and S are transitive, then $S \diamond R$ **may fail to be transitive**.

PROOF 7.3.1.5 ► PROOF OF PROPOSITION 7.3.1.4

Item 1: Interaction With Inverses

Clear.

Item 2: Interaction With Composition

See [MSE 2096272].¹

¹*Intuition:* Transitivity for R and S fails to imply that of $S \circ R$ because the composition operation for relations intertwines R and S in an incompatible way:

1. If $a \sim_{S \circ R} c$ and $c \sim_{S \circ R} e$, then:
 - (a) There is some $b \in A$ such that:
 - i. $a \sim_R b$;
 - ii. $b \sim_S c$;
 - (b) There is some $d \in A$ such that:
 - i. $c \sim_R d$;
 - ii. $d \sim_S e$.

00UZ 7.3.2 The Transitive Closure of a Relation

Let R be a relation on A .

00V0 DEFINITION 7.3.2.1 ► THE TRANSITIVE CLOSURE OF A RELATION

The **transitive closure** of \sim_R is the relation \sim_R^{trans} ¹ satisfying the following universal property:²

- (★) Given another transitive relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{trans}} \subset \sim_S$.

¹*Further Notation:* Also written R^{trans} .

²*Slogan:* The transitive closure of R is the smallest transitive relation containing R .

00V1 CONSTRUCTION 7.3.2.2 ► THE TRANSITIVE CLOSURE OF A RELATION

Concretely, \sim_R^{trans} is the free non-unital monoid on R in $(\mathbf{Rel}(A, A), \diamond)$ ¹, being given by

$$\begin{aligned} R^{\text{trans}} &\stackrel{\text{def}}{=} \prod_{n=1}^{\infty} R^{\diamond n} \\ &\stackrel{\text{def}}{=} \bigcup_{n=1}^{\infty} R^{\diamond n} \\ &\stackrel{\text{def}}{=} \left\{ (a, b) \in A \times B \mid \begin{array}{l} \text{there exists some } (x_1, \dots, x_n) \in R^{\times n} \\ \text{such that } a \sim_R x_1 \sim_R \dots \sim_R x_n \sim_R b \end{array} \right\}. \end{aligned}$$

¹Or, equivalently, the free non-unital \mathbb{B}_1 -monoid on R in $(\mathbf{N}_\bullet(\mathbf{Rel}(A, A)), \diamond)$.

PROOF 7.3.2.3 ▶ PROOF OF CONSTRUCTION 7.3.2.2

Clear. 

00V2 PROPOSITION 7.3.2.4 ▶ PROPERTIES OF THE TRANSITIVE CLOSURE OF A RELATION

Let R be a relation on A .

00V3 1. *Adjointness.* We have an adjunction

$$\left((-)^{\text{trans}} \dashv \overline{\text{忘}} \right): \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{trans}}} \\ \perp \\ \xleftarrow{\overline{\text{忘}}} \end{array} \mathbf{Rel}^{\text{trans}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{trans}}(R^{\text{trans}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{trans}}(A, A))$ and $S \in \text{Obj}(\mathbf{Rel}(A, B))$.

00V4 2. *The Transitive Closure of a Transitive Relation.* If R is transitive, then $R^{\text{trans}} = R$.

00V5 3. *Idempotency.* We have

$$(R^{\text{trans}})^{\text{trans}} = R^{\text{trans}}.$$

00V6

4. *Interaction With Inverses.* We have

$$\begin{array}{ccc} \text{Rel}(A, A) & \xrightarrow{(-)^{\text{trans}}} & \text{Rel}(A, A) \\ (-)^{\dagger} \downarrow & & \downarrow (-)^{\dagger} \\ \text{Rel}(A, A) & \xrightarrow{(-)^{\text{trans}}} & \text{Rel}(A, A). \end{array}$$

$$(R^{\dagger})^{\text{trans}} = (R^{\text{trans}})^{\dagger},$$

00V7

5. *Interaction With Composition.* We have

$$\begin{array}{ccc} \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A) \\ (-)^{\text{trans}} \times (-)^{\text{trans}} \downarrow & \times & \downarrow (-)^{\text{trans}} \\ \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A). \end{array}$$

$$(S \diamond R)^{\text{trans}} \stackrel{\text{poss.}}{\neq} S^{\text{trans}} \diamond R^{\text{trans}},$$

PROOF 7.3.2.5 ► PROOF OF PROPOSITION 7.3.2.4**Item 1: Adjointness**

This is a rephrasing of the universal property of the transitive closure of a relation, stated in [Definition 7.3.2.1](#).

Item 2: The Transitive Closure of a Transitive Relation

Clear.

Item 3: Idempotency

This follows from [Item 2](#).

Item 4: Interaction With Inverses

We have

$$\begin{aligned} (R^{\dagger})^{\text{trans}} &= \bigcup_{n=1}^{\infty} (R^{\dagger})^{\diamond n} \\ &= \bigcup_{n=1}^{\infty} (R^{\diamond n})^{\dagger} \\ &= \left(\bigcup_{n=1}^{\infty} R^{\diamond n} \right)^{\dagger} \\ &= (R^{\text{trans}})^{\dagger}, \end{aligned}$$

where we have used, respectively:

1. [Construction 7.3.2.2.](#)
2. [Item 4 of Proposition 6.3.12.3.](#)
3. [Item 1 of Proposition 6.3.6.2.](#)
4. [Construction 7.3.2.2.](#)

Item 5: Interaction With Composition

This follows from [Item 2 of Proposition 7.3.1.4.](#)



00V8 7.4 Equivalence Relations

00V9 7.4.1 Foundations

Let A be a set.

00VA DEFINITION 7.4.1.1 ► EQUIVALENCE RELATIONS

A relation R is an **equivalence relation** if it is reflexive, symmetric, and transitive.¹

¹*Further Terminology:* If instead R is just symmetric and transitive, then it is called a **partial equivalence relation**.

00VB EXAMPLE 7.4.1.2 ► THE KERNEL OF A FUNCTION

The **kernel of a function** $f: A \rightarrow B$ is the equivalence relation $\sim_{\text{Ker}(f)}$ on A obtained by declaring $a \sim_{\text{Ker}(f)} b$ iff $f(a) = f(b)$.¹

¹The kernel $\text{Ker}(f): A \dashv\vdash A$ of f is the underlying functor of the monad induced by the adjunction $\text{Gr}(f) \dashv f^{-1}: A \rightleftarrows B$ in **Rel** of [Item 2 of Proposition 6.3.1.2.](#)

00VC DEFINITION 7.4.1.3 ► THE PO/SET OF EQUIVALENCE RELATIONS ON A SET

Let A and B be sets.

- 00VD 1. The **set of equivalence relations from A to B** is the subset $\text{Rel}^{\text{eq}}(A, B)$ of $\text{Rel}(A, B)$ spanned by the equivalence relations.
- 00VE 2. The **poset of relations from A to B** is the subset $\mathbf{Rel}^{\text{eq}}(A, B)$ of $\mathbf{Rel}(A, B)$ spanned by the equivalence relations.

00VF 7.4.2 The Equivalence Closure of a Relation

Let R be a relation on A .

00VG DEFINITION 7.4.2.1 ► THE EQUIVALENCE CLOSURE OF A RELATION

The **equivalence closure**¹ of \sim_R is the relation \sim_R^{eq} ² satisfying the following universal property:³

- (★) Given another equivalence relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{eq}} \subset \sim_S$.

¹Further Terminology: Also called the **equivalence relation associated to** \sim_R .

²Further Notation: Also written R^{eq} .

³Slogan: The equivalence closure of R is the smallest equivalence relation containing R .

00VH CONSTRUCTION 7.4.2.2 ► THE EQUIVALENCE CLOSURE OF A RELATION

Concretely, \sim_R^{eq} is the equivalence relation on A defined by

$$R^{\text{eq}} \stackrel{\text{def}}{=} \left((R^{\text{refl}})^{\text{symm}} \right)^{\text{trans}}$$

$$= \left((R^{\text{symm}})^{\text{trans}} \right)^{\text{refl}}$$


$$= \left\{ (a, b) \in A \times B \mid \left. \begin{array}{l} \text{there exists } (x_1, \dots, x_n) \in R^{\times n} \text{ satisfying at} \\ \text{least one of the following conditions:} \\ \\ \text{1. The following conditions are satisfied:} \\ \\ \text{(a) We have } a \sim_R x_1 \text{ or } x_1 \sim_R a; \\ \text{(b) We have } x_i \sim_R x_{i+1} \text{ or } x_{i+1} \sim_R x_i \\ \text{for each } 1 \leq i \leq n-1; \\ \text{(c) We have } b \sim_R x_n \text{ or } x_n \sim_R b; \\ \\ \text{2. We have } a = b. \end{array} \right\}.$$

PROOF 7.4.2.3 ► PROOF OF CONSTRUCTION 7.4.2.2

From the universal properties of the reflexive, symmetric, and transitive closures of a relation (Definitions 7.1.2.1, 7.2.2.1 and 7.3.2.1), we see that it suffices to prove that:

00VJ 1. The symmetric closure of a reflexive relation is still reflexive.

00VK 2. The transitive closure of a symmetric relation is still symmetric.

which are both clear. 

00VL **PROPOSITION 7.4.2.4 ► PROPERTIES OF EQUIVALENCE RELATIONS**

Let R be a relation on A .

00VM 1. *Adjointness.* We have an adjunction

$$((-)^{\text{eq}} \dashv \text{忘}) : \mathbf{Rel}(A, B) \begin{array}{c} \xrightarrow{(-)^{\text{eq}}} \\ \perp \\ \xleftarrow{\text{忘}} \end{array} \mathbf{Rel}^{\text{eq}}(A, B),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{eq}}(R^{\text{eq}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{eq}}(A, B))$ and $S \in \text{Obj}(\mathbf{Rel}(A, B))$.

00VN 2. *The Equivalence Closure of an Equivalence Relation.* If R is an equivalence relation, then $R^{\text{eq}} = R$.

00VP 3. *Idempotency.* We have

$$(R^{\text{eq}})^{\text{eq}} = R^{\text{eq}}.$$

PROOF 7.4.2.5 ► PROOF OF PROPOSITION 7.4.2.4

Item 1: Adjointness

This is a rephrasing of the universal property of the equivalence closure of a relation, stated in [Definition 7.4.2.1](#).

Item 2: The Equivalence Closure of an Equivalence Relation

Clear.

Item 3: Idempotency

This follows from [Item 2](#). 

00VQ **7.5 Quotients by Equivalence Relations**

00VR **7.5.1 Equivalence Classes**

Let A be a set, let R be a relation on A , and let $a \in A$.

00VS

DEFINITION 7.5.1.1 ► EQUIVALENCE CLASSES

The **equivalence class associated to** a is the set $[a]$ defined by

$$\begin{aligned} [a] &\stackrel{\text{def}}{=} \{x \in X \mid x \sim_R a\} \\ &= \{x \in X \mid a \sim_R x\}. \end{aligned} \quad (\text{since } R \text{ is symmetric})$$

00VT 7.5.2 Quotients of Sets by Equivalence Relations

Let A be a set and let R be a relation on A .

00VU

DEFINITION 7.5.2.1 ► QUOTIENTS OF SETS BY EQUIVALENCE RELATIONS

The **quotient of** X **by** R is the set X/\sim_R defined by

$$X/\sim_R \stackrel{\text{def}}{=} \{[a] \in \mathcal{P}(X) \mid a \in X\}.$$

00VV

REMARK 7.5.2.2 ► WHY USE “EQUIVALENCE” RELATIONS FOR QUOTIENT SETS

The reason we define quotient sets for equivalence relations only is that each of the properties of being an equivalence relation—reflexivity, symmetry, and transitivity—ensures that the equivalence classes $[a]$ of X under R are well-behaved:

- *Reflexivity.* If R is reflexive, then, for each $a \in X$, we have $a \in [a]$.
- *Symmetry.* The equivalence class $[a]$ of an element a of X is defined by

$$[a] \stackrel{\text{def}}{=} \{x \in X \mid x \sim_R a\},$$

but we could equally well define

$$[a]' \stackrel{\text{def}}{=} \{x \in X \mid a \sim_R x\}$$

instead. This is not a problem when R is symmetric, as we then have $[a] = [a]'$.¹

- *Transitivity.* If R is transitive, then $[a]$ and $[b]$ are disjoint iff $a \not\sim_R b$, and equal otherwise.

¹When categorifying equivalence relations, one finds that $[a]$ and $[a]'$ correspond to presheaves and copresheaves; see ??.

00VW **PROPOSITION 7.5.2.3 ► PROPERTIES OF QUOTIENT SETS**

Let $f: X \rightarrow Y$ be a function and let R be a relation on X .

00VX 1. *As a Coequaliser.* We have an isomorphism of sets

$$X/\sim_R^{\text{eq}} \cong \text{CoEq} \left(R \hookrightarrow X \times X \begin{array}{c} \xrightarrow{\text{pr}_1} \\ \xrightarrow{\text{pr}_2} X \end{array} \right),$$

where \sim_R^{eq} is the equivalence relation generated by \sim_R .

00VY 2. *As a Pushout.* We have an isomorphism of sets¹

$$X/\sim_R^{\text{eq}} \cong X \amalg_{\text{Eq}(\text{pr}_1, \text{pr}_2)} X, \quad \begin{array}{ccc} X/\sim_R^{\text{eq}} & \longleftarrow & X \\ \uparrow & \ulcorner & \uparrow \\ X & \longleftarrow & \text{Eq}(\text{pr}_1, \text{pr}_2). \end{array}$$

where \sim_R^{eq} is the equivalence relation generated by \sim_R .

00VZ 3. *The First Isomorphism Theorem for Sets.* We have an isomorphism of sets^{2,3}

$$X/\sim_{\text{Ker}(f)} \cong \text{Im}(f).$$

00W0 4. *Descending Functions to Quotient Sets, I.* Let R be an equivalence relation on X . The following conditions are equivalent:

(a) There exists a map

$$\bar{f}: X/\sim_R \rightarrow Y$$

making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \exists \nearrow \bar{f} & \\ X/\sim_R & & \end{array}$$

commute.

(b) We have $R \subset \text{Ker}(f)$.

(c) For each $x, y \in X$, if $x \sim_R y$, then $f(x) = f(y)$.

00W1

5. *Descending Functions to Quotient Sets, II.* Let R be an equivalence relation on X . If the conditions of **Item 4** hold, then \bar{f} is the *unique* map making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \exists! \nearrow \bar{f} & \\ X/\sim_R & & \end{array}$$

commute.

00W2

6. *Descending Functions to Quotient Sets, III.* Let R be an equivalence relation on X . We have a bijection

$$\text{Hom}_{\text{Sets}}(X/\sim_R, Y) \cong \text{Hom}_{\text{Sets}}^R(X, Y),$$

natural in $X, Y \in \text{Obj}(\text{Sets})$, given by the assignment $f \mapsto \bar{f}$ of **Items 4** and **5**, where $\text{Hom}_{\text{Sets}}^R(X, Y)$ is the set defined by

$$\text{Hom}_{\text{Sets}}^R(X, Y) \stackrel{\text{def}}{=} \left\{ f \in \text{Hom}_{\text{Sets}}(X, Y) \left| \begin{array}{l} \text{for each } x, y \in X, \\ \text{if } x \sim_R y, \text{ then} \\ f(x) = f(y) \end{array} \right. \right\}.$$

00W3

7. *Descending Functions to Quotient Sets, IV.* Let R be an equivalence relation on X . If the conditions of **Item 4** hold, then the following conditions are equivalent:

- The map \bar{f} is an injection.
- We have $R = \text{Ker}(f)$.
- For each $x, y \in X$, we have $x \sim_R y$ iff $f(x) = f(y)$.

00W4

8. *Descending Functions to Quotient Sets, V.* Let R be an equivalence relation on X . If the conditions of **Item 4** hold, then the following conditions are equivalent:

- The map $f: X \rightarrow Y$ is surjective.
- The map $\bar{f}: X/\sim_R \rightarrow Y$ is surjective.

00W5

9. *Descending Functions to Quotient Sets, VI.* Let R be a relation on X and let \sim_R^{eq} be the equivalence relation associated to R . The following conditions are equivalent:

00W6

(a) The map f satisfies the equivalent conditions of **Item 4**:

- There exists a map

$$\bar{f}: X/\sim_R^{\text{eq}} \rightarrow Y$$

making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \exists \nearrow \bar{f} & \\ X/\sim_R^{\text{eq}} & & \end{array}$$

commute.

- For each $x, y \in X$, if $x \sim_R^{\text{eq}} y$, then $f(x) = f(y)$.

00W7

(b) For each $x, y \in X$, if $x \sim_R y$, then $f(x) = f(y)$.

¹Dually, we also have an isomorphism of sets

$$\begin{array}{ccc} \text{Eq}(pr_1, pr_2) & \longrightarrow & X \\ \downarrow \lrcorner & & \downarrow \\ X & \longrightarrow & X/\sim_R^{\text{eq}} \end{array}$$

$\text{Eq}(pr_1, pr_2) \cong X \times_{X/\sim_R^{\text{eq}}} X$,

²*Further Terminology:* The set $X/\sim_{\text{Ker}(f)}$ is often called the **coimage of f** , and denoted by $\text{Coim}(f)$.

³In a sense this is a result relating the monad in **Rel** induced by f with the comonad in **Rel** induced by f , as the kernel and image

$$\begin{aligned} \text{Ker}(f) &: X \dashv\vdash X, \\ \text{Im}(f) &\subset Y \end{aligned}$$

of f are the underlying functors of (respectively) the induced monad and comonad of the adjunction

$$(\text{Gr}(f) \dashv f^{-1}): \begin{array}{ccc} & \text{Gr}(f) & \\ & \uparrow & \\ A & \xrightarrow{\quad} & B \\ & \downarrow & \\ & f^{-1} & \end{array}$$

of **Item 2 of Proposition 6.3.1.2**.

PROOF 7.5.2.4 ► PROOF OF PROPOSITION 7.5.2.3

Item 1: As a Coequaliser

Omitted.

Item 2: As a Pushout

Omitted.

Item 3: The First Isomorphism Theorem for Sets

Clear.

Item 4: Descending Functions to Quotient Sets, I

See [Pro240].

Item 5: Descending Functions to Quotient Sets, II

See [Pro24aa].

Item 6: Descending Functions to Quotient Sets, III

This follows from [Items 5](#) and [6](#).

Item 7: Descending Functions to Quotient Sets, IV

See [Pro24n].

Item 8: Descending Functions to Quotient Sets, V

See [Pro24m].

Item 9: Descending Functions to Quotient Sets, VI


The implication [Item 9a](#) \implies [Item 9b](#) is clear.

Conversely, suppose that, for each $x, y \in X$, if $x \sim_R y$, then $f(x) = f(y)$. Spelling out the definition of the equivalence closure of R , we see that the condition $x \sim_R^{\text{eq}} y$ unwinds to the following:

- (★) There exist $(x_1, \dots, x_n) \in R^{\times n}$ satisfying at least one of the following conditions:
1. The following conditions are satisfied:
 - (a) We have $x \sim_R x_1$ or $x_1 \sim_R x$;
 - (b) We have $x_i \sim_R x_{i+1}$ or $x_{i+1} \sim_R x_i$ for each $1 \leq i \leq n - 1$;
 - (c) We have $y \sim_R x_n$ or $x_n \sim_R y$;
 2. We have $x = y$.

Now, if $x = y$, then $f(x) = f(y)$ trivially; otherwise, we have

$$\begin{aligned}f(x) &= f(x_1), \\f(x_1) &= f(x_2), \\&\vdots \\f(x_{n-1}) &= f(x_n), \\f(x_n) &= f(y),\end{aligned}$$

and $f(x) = f(y)$, as we wanted to show. 

Appendices

7.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets

Relations

5. Relations

6. Constructions With Relations

7. Equivalence Relations and Apartness Relations

Category Theory

8. Categories

Bicategories

9. Types of Morphisms in Bicategories

Part III

Category Theory

Chapter 8

Categories

00W8 This chapter contains some elementary material about categories, functors, and natural transformations. Notably, we discuss and explore:

1. Categories ([Section 8.1](#)).
2. The quadruple adjunction $\pi_0 \dashv (-)_{\text{disc}} \dashv \text{Obj} \dashv (-)_{\text{indisc}}$ between the category of categories and the category of sets ([Section 8.2](#)).
3. Groupoids, categories in which all morphisms admit inverses ([Section 8.3](#)).
4. Functors ([Section 8.4](#)).
5. The conditions one may impose on functors in decreasing order of importance:
 - (a) [Section 8.5](#) introduces the foundationally important conditions one may impose on functors, such as faithfulness, conservativity, essential surjectivity, etc.
 - (b) [Section 8.6](#) introduces more conditions one may impose on functors that are still important but less omni-present than those of [Section 8.5](#), such as being dominant, being a monomorphism, being pseudomonic, etc.
 - (c) [Section 8.7](#) introduces some rather rare or uncommon conditions one may impose on functors that are nevertheless still useful to explicit record in this chapter.
6. Natural transformations ([Section 8.8](#)).
7. The various categorical and 2-categorical structures formed by categories, functors, and natural transformations ([Section 8.9](#)).

Contents

8.1	Categories	418
8.1.1	Foundations	418
8.1.2	Examples of Categories	420
8.1.3	Posetal Categories	424
8.1.4	Subcategories	425
8.1.5	Skeletons of Categories	426
8.1.6	Precomposition and Postcomposition	428
8.2	The Quadruple Adjunction With Sets	431
8.2.1	Statement	431
8.2.2	Connected Components and Connected Categories	432
8.2.3	Discrete Categories	435
8.2.4	Indiscrete Categories	437
8.3	Groupoids	438
8.3.1	Foundations	439
8.3.2	The Groupoid Completion of a Category	439
8.3.3	The Core of a Category	443
8.4	Functors	446
8.4.1	Foundations	446
8.4.2	Contravariant Functors	450
8.4.3	Forgetful Functors	452
8.4.4	The Natural Transformation Associated to a Func-	
tor		454
8.5	Conditions on Functors	456
8.5.1	Faithful Functors	456
8.5.2	Full Functors	459
8.5.3	Fully Faithful Functors	462
8.5.4	Conservative Functors	466
8.5.5	Essentially Injective Functors	468
8.5.6	Essentially Surjective Functors	469
8.5.7	Equivalences of Categories	469
8.5.8	Isomorphisms of Categories	472
8.6	More Conditions on Functors	473
8.6.1	Dominant Functors	473
8.6.2	Monomorphisms of Categories	475
8.6.3	Epimorphisms of Categories	476
8.6.4	Pseudomonic Functors	478
8.6.5	Pseudoepic Functors	480
8.7	Even More Conditions on Functors	483
8.7.1	Injective on Objects Functors	483
8.7.2	Surjective on Objects Functors	484

8.7.3	Bijjective on Objects Functors	484
8.7.4	Functors Representably Faithful on Cores	484
8.7.5	Functors Representably Full on Cores	485
8.7.6	Functors Representably Fully Faithful on Cores	486
8.7.7	Functors Corepresentably Faithful on Cores	487
8.7.8	Functors Corepresentably Full on Cores	489
8.7.9	Functors Corepresentably Fully Faithful on Cores	490
8.8	Natural Transformations	492
8.8.1	Transformations	492
8.8.2	Natural Transformations	492
8.8.3	Vertical Composition of Natural Transformations	493
8.8.4	Horizontal Composition of Natural Transformations	
497		
8.8.5	Properties of Natural Transformations	503
8.8.6	Natural Isomorphisms	504
8.9	Categories of Categories	506
8.9.1	Functor Categories	506
8.9.2	The Category of Categories and Functors	510
8.9.3	The 2-Category of Categories, Functors, and Natu- ral Transformations	511
8.9.4	The Category of Groupoids	512
8.9.5	The 2-Category of Groupoids	512
8.A	Other Chapters	513

00W9 8.1 Categories

00WA 8.1.1 Foundations

00WB

DEFINITION 8.1.1.1 ► CATEGORIES

A **category** $(C, \circ^C, \mathbb{1}^C)$ consists of:

- *Objects.* A class $\text{Obj}(C)$ of **objects**.
- *Morphisms.* For each $A, B \in \text{Obj}(C)$, a class $\text{Hom}_C(A, B)$, called the **class of morphisms of C from A to B** .
- *Identities.* For each $A \in \text{Obj}(C)$, a map of sets

$$\mathbb{1}_A^C : \text{pt} \rightarrow \text{Hom}_C(A, A),$$

called the **unit map of C at A** , determining a morphism

$$\text{id}_A: A \rightarrow A$$

of C , called the **identity morphism of A** .


- *Composition.* For each $A, B, C \in \text{Obj}(C)$, a map of sets

$$\circ_{A,B,C}^C: \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) \rightarrow \text{Hom}_C(A, C),$$

called the **composition map of C at (A, B, C)** .

such that the following conditions are satisfied:


1. *Associativity.* The diagram



commutes, i.e. for each composable triple (f, g, h) of morphisms of C , we have

$$(f \circ g) \circ h = f \circ (g \circ h).$$

2. *Left Unitality.* The diagram



commutes, i.e. for each morphism $f: A \rightarrow B$ of C , we have

$$\text{id}_B \circ f = f.$$

3. *Right Unitality*. The diagram

$$\begin{array}{ccc}
 \text{Hom}_C(A, B) \times \text{pt} & & \\
 \downarrow \text{id}_{\text{Hom}_C(A, B)} \times \mathbb{1}_A^C & \searrow \rho_{\text{Hom}_C(A, B)}^{\text{Sets}} & \\
 \text{Hom}_C(A, B) \times \text{Hom}_C(A, A) & \xrightarrow{\circ_{A, A, B}^C} & \text{Hom}_C(A, B)
 \end{array}$$

commutes, i.e. for each morphism $f: A \rightarrow B$ of C , we have

$$f \circ \text{id}_A = f.$$

00WC NOTATION 8.1.1.2 ► FURTHER NOTATION FOR MORPHISMS IN CATEGORIES

Let C be a category.

- 00WD 1. We also write $C(A, B)$ for $\text{Hom}_C(A, B)$.
- 00WE 2. We write $\text{Mor}(C)$ for the class of all morphisms of C .

00WF DEFINITION 8.1.1.3 ► SIZE CONDITIONS ON CATEGORIES

Let κ be a regular cardinal. A category C is

- 00WG 1. **Locally small** if, for each $A, B \in \text{Obj}(C)$, the class $\text{Hom}_C(A, B)$ is a set.
- 00WH 2. **Locally essentially small** if, for each $A, B \in \text{Obj}(C)$, the class

$$\text{Hom}_C(A, B) / \{\text{isomorphisms}\}$$
 is a set.
- 00WJ 3. **Small** if C is locally small and $\text{Obj}(C)$ is a set.
- 00WK 4. **κ -Small** if C is locally small, $\text{Obj}(C)$ is a set, and we have $\#\text{Obj}(C) < \kappa$.

00WL 8.1.2 Examples of Categories

00WM

EXAMPLE 8.1.2.1 ► THE PUNCTUAL CATEGORY

The **punctual category**¹ is the category pt where

- *Objects.* We have

$$\text{Obj}(\text{pt}) \stackrel{\text{def}}{=} \{\star\}.$$

- *Morphisms.* The unique Hom-set of pt is defined by

$$\text{Hom}_{\text{pt}}(\star, \star) \stackrel{\text{def}}{=} \{\text{id}_{\star}\}.$$

- *Identities.* The unit map

$$\mathbb{1}_{\star}^{\text{pt}} : \text{pt} \rightarrow \text{Hom}_{\text{pt}}(\star, \star)$$

of pt at \star is defined by

$$\text{id}_{\star}^{\text{pt}} \stackrel{\text{def}}{=} \text{id}_{\star}.$$

- *Composition.* The composition map

$$\circ_{\star, \star, \star}^{\text{pt}} : \text{Hom}_{\text{pt}}(\star, \star) \times \text{Hom}_{\text{pt}}(\star, \star) \rightarrow \text{Hom}_{\text{pt}}(\star, \star)$$

of pt at (\star, \star, \star) is given by the bijection $\text{pt} \times \text{pt} \cong \text{pt}$.

¹Further Terminology: Also called the **singleton category**.

00WN

EXAMPLE 8.1.2.2 ► MONOIDS AS ONE-OBJECT CATEGORIES

We have an isomorphism of categories¹

$$\text{Mon} \cong_{\text{Sets}} \text{pt} \times \text{Cats},$$

$$\begin{array}{ccc} \text{Mon} & \longrightarrow & \text{Cats} \\ \downarrow \lrcorner & & \downarrow \text{Obj} \\ \text{pt} & \xrightarrow{[\text{pt}]} & \text{Sets} \end{array}$$

via the delooping functor $B : \text{Mon} \rightarrow \text{Cats}$ of ?? of ??, exhibiting monoids as exactly those categories having a single object.

¹This can be enhanced to an isomorphism of 2-categories

$$\text{Mon}_{2\text{disc}} \cong \text{pt}_{\text{bi}} \times_{\text{Sets}_{2\text{disc}}} \text{Cats}_{2,*}$$

$$\begin{array}{ccc} \text{Mon}_{2\text{disc}} & \longrightarrow & \text{Cats}_{2,*} \\ \downarrow & \lrcorner & \downarrow \text{Obj} \\ \text{pt}_{\text{bi}} & \xrightarrow{[\text{pt}]} & \text{Sets}_{2\text{disc}} \end{array}$$

between the discrete 2-category $\text{Mon}_{2\text{disc}}$ on Mon and the 2-category of pointed categories with one object.

PROOF 8.1.2.3 ▶ PROOF OF EXAMPLE 8.1.2.2
 Omitted. 

00WP

EXAMPLE 8.1.2.4 ▶ THE EMPTY CATEGORY

The **empty category** is the category \emptyset_{cat} where

- *Objects.* We have $\text{Obj}(\emptyset_{\text{cat}}) \stackrel{\text{def}}{=} \emptyset$.
- *Morphisms.* We have $\text{Mor}(\emptyset_{\text{cat}}) \stackrel{\text{def}}{=} \emptyset$.
- *Identities and Composition.* Having no objects, \emptyset_{cat} has no unit nor composition maps.

00WQ

EXAMPLE 8.1.2.5 ▶ ORDINAL CATEGORIES

The **n th ordinal category** is the category \mathfrak{n} where¹

- *Objects.* We have $\text{Obj}(\mathfrak{n}) \stackrel{\text{def}}{=} \{[0], \dots, [n]\}$.
- *Morphisms.* For each $[i], [j] \in \text{Obj}(\mathfrak{n})$, we have

$$\text{Hom}_{\mathfrak{n}}([i], [j]) \stackrel{\text{def}}{=} \begin{cases} \{\text{id}_{[i]}\} & \text{if } [i] = [j], \\ \{[i] \rightarrow [j]\} & \text{if } [j] < [i], \\ \emptyset & \text{if } [j] > [i]. \end{cases}$$

- *Identities.* For each $[i] \in \text{Obj}(\mathfrak{n})$, the unit map

$$\mathbb{1}_{[i]}^{\mathfrak{n}} : \text{pt} \rightarrow \text{Hom}_{\mathfrak{n}}([i], [i])$$

of \mathfrak{n} at $[i]$ is defined by

$$\text{id}_{[i]}^{\mathfrak{n}} \stackrel{\text{def}}{=} \mathbb{1}_{[i]}^{\mathfrak{n}}.$$

- *Composition.* For each $[i], [j], [k] \in \text{Obj}(\mathfrak{n})$, the composition map

$$\circ_{[i],[j],[k]}^{\mathfrak{n}} : \text{Hom}_{\mathfrak{n}}([j], [k]) \times \text{Hom}_{\mathfrak{n}}([i], [j]) \rightarrow \text{Hom}_{\mathfrak{n}}([i], [k])$$

of \mathfrak{n} at $([i], [j], [k])$ is defined by

$$\begin{aligned} \text{id}_{[i]} \circ \text{id}_{[i]} &= \text{id}_{[i]}, \\ ([j] \rightarrow [k]) \circ ([i] \rightarrow [j]) &= ([i] \rightarrow [k]). \end{aligned}$$

¹In other words, \mathfrak{n} is the category associated to the poset

$$[0] \rightarrow [1] \rightarrow \cdots \rightarrow [n-1] \rightarrow [n].$$

The category \mathfrak{n} for $n \geq 2$ may also be defined in terms of $\mathfrak{0}$ and joins (\star): we have isomorphisms of categories

$$\begin{aligned} \mathbb{1} &\cong \mathfrak{0} \star \mathfrak{0}, \\ \mathbb{2} &\cong \mathbb{1} \star \mathfrak{0} \\ &\cong (\mathfrak{0} \star \mathfrak{0}) \star \mathfrak{0}, \\ \mathbb{3} &\cong \mathbb{2} \star \mathfrak{0} \\ &\cong (\mathbb{1} \star \mathfrak{0}) \star \mathfrak{0} \\ &\cong ((\mathfrak{0} \star \mathfrak{0}) \star \mathfrak{0}) \star \mathfrak{0}, \\ \mathbb{4} &\cong \mathbb{3} \star \mathfrak{0} \\ &\cong (\mathbb{2} \star \mathfrak{0}) \star \mathfrak{0} \\ &\cong ((\mathbb{1} \star \mathfrak{0}) \star \mathfrak{0}) \star \mathfrak{0} \\ &\cong (((\mathfrak{0} \star \mathfrak{0}) \star \mathfrak{0}) \star \mathfrak{0}) \star \mathfrak{0}, \end{aligned}$$

and so on.

EXAMPLE 8.1.2.6 ► MORE EXAMPLES OF CATEGORIES

Here we list some of the other categories appearing throughout this work.

1. The category Sets_* of pointed sets of [Definition 3.1.3.1](#).
2. The category Rel of sets and relations of [Definition 5.2.1.1](#).
3. The category $\text{Span}(A, B)$ of spans from a set A to a set B of [??](#).

- 00WV 4. The category $\mathbf{ISets}(K)$ of K -indexed sets of ??.
- 00WW 5. The category \mathbf{ISets} of indexed sets of ??.
- 00WX 6. The category $\mathbf{FibSets}(K)$ of K -fibred sets of ??.
- 00WY 7. The category $\mathbf{FibSets}$ of fibred sets of ??.
- 00WZ 8. Categories of functors $\mathbf{Fun}(C, \mathcal{D})$ as in [Definition 8.9.1.1](#).
- 00X0 9. The category of categories \mathbf{Cats} of [Definition 8.9.2.1](#).
- 00X1 10. The category of groupoids \mathbf{Grpd} of [Definition 8.9.4.1](#).

00X2 8.1.3 Posetal Categories

00X3 DEFINITION 8.1.3.1 ► POSETAL CATEGORIES

Let (X, \preceq_X) be a poset.

1. The **posetal category associated to** (X, \preceq_X) is the category X_{pos} where

00X4

- *Objects.* We have

$$\text{Obj}(X_{\text{pos}}) \stackrel{\text{def}}{=} X.$$

- *Morphisms.* For each $a, b \in \text{Obj}(X_{\text{pos}})$, we have

$$\text{Hom}_{X_{\text{pos}}}(a, b) \stackrel{\text{def}}{=} \begin{cases} \text{pt} & \text{if } a \preceq_X b, \\ \emptyset & \text{otherwise.} \end{cases}$$

- *Identities.* For each $a \in \text{Obj}(X_{\text{pos}})$, the unit map

$$\mathbb{1}_a^{X_{\text{pos}}} : \text{pt} \rightarrow \text{Hom}_{X_{\text{pos}}}(a, a)$$

of X_{pos} at a is given by the identity map.

- *Composition.* For each $a, b, c \in \text{Obj}(X_{\text{pos}})$, the composition map

$$\circ_{a,b,c}^{X_{\text{pos}}} : \text{Hom}_{X_{\text{pos}}}(b, c) \times \text{Hom}_{X_{\text{pos}}}(a, b) \rightarrow \text{Hom}_{X_{\text{pos}}}(a, c)$$

of X_{pos} at (a, b, c) is defined as either the inclusion $\emptyset \hookrightarrow \text{pt}$ or the identity map of pt , depending on whether we have $a \preceq_X b$, $b \preceq_X c$, and $a \preceq_X c$.

00X5 2. A category C is **posetal**¹ if C is equivalent to X_{pos} for some poset (X, \preceq_X) .

¹Further Terminology: Also called a **thin** category or a **(0, 1)-category**.

00X6 **PROPOSITION 8.1.3.2 ► PROPERTIES OF POSETAL CATEGORIES**

Let (X, \preceq_X) be a poset and let C be a category.

00X7 1. *Functoriality.* The assignment $(X, \preceq_X) \mapsto X_{\text{pos}}$ defines a functor

$$(-)_{\text{pos}}: \text{Pos} \rightarrow \text{Cats}.$$

00X8 2. *Fully Faithfulness.* The functor $(-)_{\text{pos}}$ of **Item 1** is fully faithful.

00X9 3. *Characterisations.* The following conditions are equivalent:

00XA (a) The category C is posetal.

00XB (b) For each $A, B \in \text{Obj}(C)$ and each $f, g \in \text{Hom}_C(A, B)$, we have $f = g$.

PROOF 8.1.3.3 ► PROOF OF PROPOSITION 8.1.3.2

Item 1: Functoriality

Omitted.

Item 2: Fully Faithfulness

Omitted.

Item 3: Characterisations

Clear. 

00XC **8.1.4 Subcategories**

Let C be a category.

00XD **DEFINITION 8.1.4.1 ► SUBCATEGORIES**

A **subcategory** of C is a category \mathcal{A} satisfying the following conditions:

1. *Objects.* We have $\text{Obj}(\mathcal{A}) \subset \text{Obj}(C)$.

2. *Morphisms.* For each $A, B \in \text{Obj}(\mathcal{A})$, we have

$$\text{Hom}_{\mathcal{A}}(A, B) \subset \text{Hom}_C(A, B).$$

3. *Identities.* For each $A \in \text{Obj}(\mathcal{A})$, we have

$$\mathbb{1}_A^{\mathcal{A}} = \mathbb{1}_A^C.$$

4. *Composition.* For each $A, B, C \in \text{Obj}(\mathcal{A})$, we have

$$\circ_{A,B,C}^{\mathcal{A}} = \circ_{A,B,C}^C.$$

00XE DEFINITION 8.1.4.2 ► FULL SUBCATEGORIES

A subcategory \mathcal{A} of C is **full** if the canonical inclusion functor $\mathcal{A} \rightarrow C$ is full, i.e. if, for each $A, B \in \text{Obj}(\mathcal{A})$, the inclusion

$$\iota_{A,B}: \text{Hom}_{\mathcal{A}}(A, B) \hookrightarrow \text{Hom}_C(A, B)$$

is surjective (and thus bijective).

00XF DEFINITION 8.1.4.3 ► STRICTLY FULL SUBCATEGORIES

A subcategory \mathcal{A} of a category C is **strictly full** if it satisfies the following conditions:

1. *Fullness.* The subcategory \mathcal{A} is full.
2. *Closedness Under Isomorphisms.* The class $\text{Obj}(\mathcal{A})$ is closed under isomorphisms.¹

¹That is, given $A \in \text{Obj}(\mathcal{A})$ and $C \in \text{Obj}(C)$, if $C \cong A$, then $C \in \text{Obj}(\mathcal{A})$.

00XG DEFINITION 8.1.4.4 ► WIDE SUBCATEGORIES

A subcategory \mathcal{A} of C is **wide**¹ if $\text{Obj}(\mathcal{A}) = \text{Obj}(C)$.

¹*Further Terminology:* Also called **lluf**.

00XH 8.1.5 Skeletons of Categories

00XJ

DEFINITION 8.1.5.1 ► SKELETONS OF CATEGORIES

A¹ **skeleton** of a category C is a full subcategory $\text{Sk}(C)$ with one object from each isomorphism class of objects of C .

¹Due to [Item 3](#) of [Proposition 8.1.5.3](#), we often refer to any such full subcategory $\text{Sk}(C)$ of C as *the* skeleton of C .

00XK

DEFINITION 8.1.5.2 ► SKELETAL CATEGORIES

A category C is **skeletal** if $C \cong \text{Sk}(C)$.¹

¹That is, C is **skeletal** if isomorphic objects of C are equal.

00XL

PROPOSITION 8.1.5.3 ► PROPERTIES OF SKELETONS OF CATEGORIES

Let C be a category.

00XM

1. *Existence.* Assuming the axiom of choice, $\text{Sk}(C)$ always exists.

00XN

2. *Pseudofunctoriality.* The assignment $C \mapsto \text{Sk}(C)$ defines a pseudo-functor

$$\text{Sk}: \text{Cats}_2 \rightarrow \text{Cats}_2.$$

00XP

3. *Uniqueness Up to Equivalence.* Any two skeletons of C are equivalent.

00XQ

4. *Inclusions of Skeletons Are Equivalences.* The inclusion

$$\iota_C: \text{Sk}(C) \hookrightarrow C$$

of a skeleton of C into C is an equivalence of categories.

PROOF 8.1.5.4 ► PROOF OF PROPOSITION 8.1.5.3

Item 1: Existence

See [\[nLab23, Section “Existence of Skeletons of Categories”\]](#).

Item 2: Pseudofunctoriality

See [\[nLab23, Section “Skeletons as an Endo-Pseudofunctor on \$\mathcal{C}at\$ ”\]](#).

Item 3: Uniqueness Up to Equivalence

Clear.

Item 4: Inclusions of Skeletons Are Equivalences

Clear.



ØØXR 8.1.6 Precomposition and Postcomposition

Let C be a category and let $A, B, C \in \text{Obj}(C)$.

00XS

DEFINITION 8.1.6.1 ► PRECOMPOSITION AND POSTCOMPOSITION FUNCTIONS

Let $f: A \rightarrow B$ and $g: B \rightarrow C$ be morphisms of \mathcal{C} .

00XT

1. The **precomposition function associated to f** is the function

$$f^*: \text{Hom}_{\mathcal{C}}(B, C) \rightarrow \text{Hom}_{\mathcal{C}}(A, C)$$

defined by

$$f^*(\phi) \stackrel{\text{def}}{=} \phi \circ f$$

for each $\phi \in \text{Hom}_{\mathcal{C}}(B, C)$.

00XU

2. The **postcomposition function associated to g** is the function

$$g_*: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{C}}(A, C)$$

defined by

$$g_*(\phi) \stackrel{\text{def}}{=} g \circ \phi$$

for each $\phi \in \text{Hom}_{\mathcal{C}}(A, B)$.

00XV

PROPOSITION 8.1.6.2 ► PROPERTIES OF PRE/POSTCOMPOSITION

Let $A, B, C, D \in \text{Obj}(\mathcal{C})$ and let $f: A \rightarrow B$ and $g: B \rightarrow C$ be morphisms of \mathcal{C} .

00XW

1. *Interaction Between Precomposition and Postcomposition.* We have

$$g_* \circ f^* = f^* \circ g_*$$

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(B, C) & \xrightarrow{g_*} & \text{Hom}_{\mathcal{C}}(B, D) \\ f^* \downarrow & & \downarrow f^* \\ \text{Hom}_{\mathcal{C}}(A, C) & \xrightarrow{g_*} & \text{Hom}_{\mathcal{C}}(A, D). \end{array}$$

00XX

2. *Interaction With Composition I.* We have

$$\begin{array}{ccc}
 \text{Hom}_C(X, A) & \xrightarrow{f_*} & \text{Hom}_C(X, B) \\
 & \searrow (g \circ f)_* & \downarrow g_* \\
 & & \text{Hom}_C(X, C), \\
 \\
 \text{Hom}_C(C, X) & \xrightarrow{g^*} & \text{Hom}_C(B, X) \\
 & \searrow (g \circ f)^* & \downarrow f^* \\
 & & \text{Hom}_C(A, X).
 \end{array}$$

$(g \circ f)^* = f^* \circ g^*$,
 $(g \circ f)_* = g_* \circ f_*$

00XY

3. *Interaction With Composition II.* We have

$$\begin{array}{ccc}
 \text{pt} \xrightarrow{[f]} \text{Hom}_C(A, B) & & \text{pt} \xrightarrow{[g]} \text{Hom}_C(B, C) \\
 \downarrow [g \circ f] & \downarrow g_* & \downarrow f^* \\
 \text{Hom}_C(A, C) & & \text{Hom}_C(A, C).
 \end{array}$$

$[g \circ f] = g_* \circ [f]$,
 $[g \circ f] = f^* \circ [g]$

00XZ

4. *Interaction With Composition III.* We have

$$\begin{array}{ccc}
 \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,C}^C} & \text{Hom}_C(A, C) \\
 \downarrow \text{id} \times f_* & & \downarrow f^* \\
 \text{Hom}_C(B, C) \times \text{Hom}_C(X, B) & \xrightarrow{\circ_{X,B,C}^C} & \text{Hom}_C(X, C), \\
 \\
 \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,C}^C} & \text{Hom}_C(A, C) \\
 \downarrow g_* \times \text{id} & & \downarrow g^* \\
 \text{Hom}_C(B, D) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,D}^C} & \text{Hom}_C(A, D).
 \end{array}$$

$f^* \circ \circ_{A,B,C}^C = \circ_{X,B,C}^C \circ (f^* \times \text{id})$,
 $g^* \circ \circ_{A,B,C}^C = \circ_{A,B,D}^C \circ (\text{id} \times g^*)$

00Y0

5. *Interaction With Identities.* We have

$$\begin{aligned}
 (\text{id}_A)^* &= \text{id}_{\text{Hom}_C(A,B)}, \\
 (\text{id}_B)_* &= \text{id}_{\text{Hom}_C(A,B)}.
 \end{aligned}$$

PROOF 8.1.6.3 ► PROOF OF PROPOSITION 8.1.6.2

Item 1: Interaction Between Precomposition and Postcomposition
Clear.

Item 2: Interaction With Composition I
Clear.

Item 3: Interaction With Composition II
Clear.

Item 4: Interaction With Composition III
Clear.

Item 5: Interaction With Identities
Clear.


00Y1 **8.2 The Quadruple Adjunction With Sets**

00Y2 **8.2.1 Statement**

Let C be a category.

00Y3 **PROPOSITION 8.2.1.1 ► THE QUADRUPLE ADJUNCTION BETWEEN Sets AND Cats**

We have a quadruple adjunction



witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\text{Sets}}(\pi_0(C), X) &\cong \text{Hom}_{\text{Cats}}(C, X_{disc}), \\ \text{Hom}_{\text{Cats}}(X_{disc}, C) &\cong \text{Hom}_{\text{Sets}}(X, \text{Obj}(C)), \\ \text{Hom}_{\text{Sets}}(\text{Obj}(C), X) &\cong \text{Hom}_{\text{Cats}}(C, X_{indisc}), \end{aligned}$$

natural in $C \in \text{Obj}(\text{Cats})$ and $X \in \text{Obj}(\text{Sets})$, where

- The functor

$$\pi_0: \text{Cats} \rightarrow \text{Sets},$$

the **connected components functor**, is the functor sending a category to its set of connected components of [Definition 8.2.2.2](#).

- The functor

$$(-)_{\text{disc}} : \text{Sets} \rightarrow \text{Cats},$$

the **discrete category functor**, is the functor sending a set to its associated discrete category of [Item 1](#).

- The functor

$$\text{Obj} : \text{Cats} \rightarrow \text{Sets},$$

the **object functor**, is the functor sending a category to its set of objects.

- The functor

$$(-)_{\text{indisc}} : \text{Sets} \rightarrow \text{Cats},$$

the **indiscrete category functor**, is the functor sending a set to its associated indiscrete category of [Item 1](#).

PROOF 8.2.1.2 ► PROOF OF PROPOSITION 8.2.1.1

Omitted.



00Y4 8.2.2 Connected Components and Connected Categories

00Y5 8.2.2.1 Connected Components of Categories

Let C be a category.

00Y6 DEFINITION 8.2.2.1 ► CONNECTED COMPONENTS OF CATEGORIES

A **connected component** of C is a full subcategory I of C satisfying the following conditions:¹

1. *Non-Emptiness.* We have $\text{Obj}(I) \neq \emptyset$.
2. *Connectedness.* There exists a zigzag of arrows between any two objects of I .

¹In other words, a **connected component** of C is an element of the set $\text{Obj}(C)/\sim$ with \sim the equivalence relation generated by the relation \sim' obtained by declaring $A \sim' B$ iff there exists a morphism of C from A to B .

00Y7 8.2.2.2 Sets of Connected Components of Categories

Let C be a category.

00Y8

DEFINITION 8.2.2.2 ▶ SETS OF CONNECTED COMPONENTS OF CATEGORIES

The **set of connected components of C** is the set $\pi_0(C)$ whose elements are the connected components of C .

00Y9

PROPOSITION 8.2.2.3 ▶ PROPERTIES OF SETS OF CONNECTED COMPONENTS

Let C be a category.


00YA

1. *Functoriality.* The assignment $C \mapsto \pi_0(C)$ defines a functor

$$\pi_0: \text{Cats} \rightarrow \text{Sets}.$$

00YB

2. *Adjointness.* We have a quadruple adjunction



00YC

3. *Interaction With Groupoids.* If C is a groupoid, then we have an isomorphism of categories

$$\pi_0(C) \cong \text{K}(C),$$

where $\text{K}(C)$ is the set of isomorphism classes of C of ??.

00YD

4. *Preservation of Colimits.* The functor π_0 of **Item 1** preserves colimits. In particular, we have bijections of sets

$$\begin{aligned} \pi_0(C \amalg \mathcal{D}) &\cong \pi_0(C) \amalg \pi_0(\mathcal{D}), \\ \pi_0(C \amalg_{\mathcal{E}} \mathcal{D}) &\cong \pi_0(C) \amalg_{\pi_0(\mathcal{E})} \pi_0(\mathcal{D}), \\ \pi_0\left(\text{CoEq}\left(C \begin{array}{c} \xrightarrow{F} \\ \xrightarrow{G} \end{array} \mathcal{D}\right)\right) &\cong \text{CoEq}\left(\pi_0(C) \begin{array}{c} \xrightarrow{\pi_0(F)} \\ \xrightarrow{\pi_0(G)} \end{array} \pi_0(\mathcal{D})\right), \end{aligned}$$

natural in $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats})$.

00YE

5. *Symmetric Strong Monoidality With Respect to Coproducts.* The connected components functor of **Item 1** has a symmetric strong monoidal structure

$$\left(\pi_0, \pi_0 \amalg, \pi_0|_{\perp}\right): (\text{Cats}, \amalg, \emptyset_{\text{cat}}) \rightarrow (\text{Sets}, \amalg, \emptyset),$$

being equipped with isomorphisms

$$\begin{aligned}\prod_{0|C, \mathcal{D}}: \pi_0(C) \amalg \pi_0(\mathcal{D}) &\xrightarrow{\cong} \pi_0(C \amalg \mathcal{D}), \\ \prod_{0|\mathbb{1}}: \emptyset &\xrightarrow{\cong} \pi_0(\emptyset_{\text{cat}}),\end{aligned}$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$.

00YF

6. *Symmetric Strong Monoidality With Respect to Products.* The connected components functor of [Item 1](#) has a symmetric strong monoidal structure

$$\left(\pi_0, \pi_0^\times, \pi_{0|\mathbb{1}}^\times\right): (\text{Cats}, \times, \text{pt}) \rightarrow (\text{Sets}, \times, \text{pt}),$$

being equipped with isomorphisms

$$\begin{aligned}\pi_{0|C, \mathcal{D}}^\times: \pi_0(C) \times \pi_0(\mathcal{D}) &\xrightarrow{\cong} \pi_0(C \times \mathcal{D}), \\ \pi_{0|\mathbb{1}}^\times: \text{pt} &\xrightarrow{\cong} \pi_0(\text{pt}),\end{aligned}$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$.

PROOF 8.2.2.4 ► PROOF OF PROPOSITION 8.2.2.3

Item 1: Functoriality

Clear.

Item 2: Adjointness

This is proved in [Proposition 8.2.1.1](#).

Item 3: Interaction With Groupoids

Clear.

Item 4: Preservation of Colimits

This follows from [Item 2](#) and ?? of ??.

Item 5: Symmetric Strong Monoidality With Respect to Coproducts

Clear.

Item 6: Symmetric Strong Monoidality With Respect to Products

Clear. 

00YG 8.2.2.3 Connected Categories

00YH DEFINITION 8.2.2.5 ► CONNECTED CATEGORIES

A category C is **connected** if $\pi_0(C) \cong \text{pt.}$ ^{1,2}

¹*Further Terminology:* A category is **disconnected** if it is not connected.

²*Example:* A groupoid is connected iff any two of its objects are isomorphic.

00YJ 8.2.3 Discrete Categories

00YK DEFINITION 8.2.3.1 ► DISCRETE CATEGORIES

Let X be a set.

00YL 1. The **discrete category on X** is the category X_{disc} where

• *Objects.* We have

$$\text{Obj}(X_{\text{disc}}) \stackrel{\text{def}}{=} X.$$

• *Morphisms.* For each $A, B \in \text{Obj}(X_{\text{disc}})$, we have

$$\text{Hom}_{X_{\text{disc}}}(A, B) \stackrel{\text{def}}{=} \begin{cases} \text{id}_A & \text{if } A = B, \\ \emptyset & \text{if } A \neq B. \end{cases}$$

• *Identities.* For each $A \in \text{Obj}(X_{\text{disc}})$, the unit map

$$\mathbb{1}_A^{X_{\text{disc}}} : \text{pt} \rightarrow \text{Hom}_{X_{\text{disc}}}(A, A)$$

of X_{disc} at A is defined by

$$\text{id}_A^{X_{\text{disc}}} \stackrel{\text{def}}{=} \text{id}_A.$$

• *Composition.* For each $A, B, C \in \text{Obj}(X_{\text{disc}})$, the composition map

$$\circ_{A,B,C}^{X_{\text{disc}}} : \text{Hom}_{X_{\text{disc}}}(B, C) \times \text{Hom}_{X_{\text{disc}}}(A, B) \rightarrow \text{Hom}_{X_{\text{disc}}}(A, C)$$

of X_{disc} at (A, B, C) is defined by

$$\text{id}_A \circ \text{id}_A \stackrel{\text{def}}{=} \text{id}_A.$$

00YM 2. A category C is **discrete** if it is equivalent to X_{disc} for some set X .

00YN **PROPOSITION 8.2.3.2 ► PROPERTIES OF DISCRETE CATEGORIES ON SETS**

Let X be a set.

- 00YP 1. *Functoriality.* The assignment $X \mapsto X_{\text{disc}}$ defines a functor

$$(-)_{\text{disc}}: \text{Sets} \rightarrow \text{Cats}.$$

- 00YQ 2. *Adjointness.* We have a quadruple adjunction

$$(\pi_0 \dashv (-)_{\text{disc}} \dashv \text{Obj} \dashv (-)_{\text{indisc}}): \text{Sets} \begin{array}{c} \xrightarrow{\pi_0} \\ \dashv \text{disc} \\ \dashv \text{Obj} \\ \xrightarrow{(-)_{\text{indisc}}} \end{array} \text{Cats}.$$

- 00YR 3. *Symmetric Strong Monoidality With Respect to Coproducts.* The functor of **Item 1** has a symmetric strong monoidal structure

$$\left((-)_{\text{disc}}, (-)_{\text{disc}}, (-)_{\text{disc}|\mathbb{1}} \right): (\text{Sets}, \amalg, \emptyset) \rightarrow (\text{Cats}, \amalg, \emptyset_{\text{cat}}),$$

being equipped with isomorphisms

$$\begin{aligned} (-)_{\text{disc}|\mathbb{1}}: X_{\text{disc}} \amalg Y_{\text{disc}} &\xrightarrow{\cong} (X \amalg Y)_{\text{disc}}, \\ (-)_{\text{disc}|\mathbb{1}}: \emptyset_{\text{cat}} &\xrightarrow{\cong} \emptyset_{\text{disc}}, \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

- 00YS 4. *Symmetric Strong Monoidality With Respect to Products.* The functor of **Item 1** has a symmetric strong monoidal structure

$$\left((-)_{\text{disc}}, (-)_{\text{disc}}^{\times}, (-)_{\text{disc}|\mathbb{1}}^{\times} \right): (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Cats}, \times, \text{pt}),$$

being equipped with isomorphisms

$$\begin{aligned} (-)_{\text{disc}|\mathbb{1}}^{\times}: X_{\text{disc}} \times Y_{\text{disc}} &\xrightarrow{\cong} (X \times Y)_{\text{disc}}, \\ (-)_{\text{disc}|\mathbb{1}}^{\times}: \text{pt} &\xrightarrow{\cong} \text{pt}_{\text{disc}}, \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

PROOF 8.2.3.3 ► PROOF OF PROPOSITION 8.2.3.2

Item 1: Functoriality

Clear.

Item 2: Adjointness

This is proved in [Proposition 8.2.1.1](#).

Item 3: Symmetric Strong Monoidality With Respect to Coproducts

Clear.

Item 4: Symmetric Strong Monoidality With Respect to Products

Clear. 00YT **8.2.4 Indiscrete Categories**

00YU DEFINITION 8.2.4.1 ► INDISCRETE CATEGORIES

Let X be a set.00YV 1. The **indiscrete category on X** ¹ is the category X_{indisc} where· *Objects.* We have

$$\text{Obj}(X_{\text{indisc}}) \stackrel{\text{def}}{=} X.$$

· *Morphisms.* For each $A, B \in \text{Obj}(X_{\text{indisc}})$, we have

$$\begin{aligned} \text{Hom}_{X_{\text{disc}}}(A, B) &\stackrel{\text{def}}{=} \{[A] \rightarrow [B]\} \\ &\cong \text{pt}. \end{aligned}$$

· *Identities.* For each $A \in \text{Obj}(X_{\text{indisc}})$, the unit map

$$\mathbb{1}_A^{X_{\text{indisc}}} : \text{pt} \rightarrow \text{Hom}_{X_{\text{indisc}}}(A, A)$$

of X_{indisc} at A is defined by

$$\text{id}_A^{X_{\text{indisc}}} \stackrel{\text{def}}{=} \{[A] \rightarrow [A]\}.$$

· *Composition.* For each $A, B, C \in \text{Obj}(X_{\text{indisc}})$, the composition map

$$\circ_{A,B,C}^{X_{\text{indisc}}} : \text{Hom}_{X_{\text{indisc}}}(B, C) \times \text{Hom}_{X_{\text{indisc}}}(A, B) \rightarrow \text{Hom}_{X_{\text{indisc}}}(A, C)$$

of X_{disc} at (A, B, C) is defined by

$$([B] \rightarrow [C]) \circ ([A] \rightarrow [B]) \stackrel{\text{def}}{=} ([A] \rightarrow [C]).$$

00YW 2. A category C is **indiscrete** if it is equivalent to X_{indisc} for some set X .¹Further Terminology: Sometimes called the **chaotic category on X** .


00YX **PROPOSITION 8.2.4.2 ► PROPERTIES OF INDISCRETE CATEGORIES ON SETS**

Let X be a set.

00YY 1. *Functoriality.* The assignment $X \mapsto X_{\text{indisc}}$ defines a functor

$$(-)_{\text{indisc}} : \text{Sets} \rightarrow \text{Cats}.$$

00YZ 2. *Adjointness.* We have a quadruple adjunction



00Z0 3. *Symmetric Strong Monoidality With Respect to Products.* The functor of **Item 1** has a symmetric strong monoidal structure

$$\left((-)_{\text{indisc}}, (-)_{\text{indisc}}^{\times}, (-)_{\text{indisc}|\mathbb{1}}^{\times} \right) : (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Cats}, \times, \text{pt}),$$

being equipped with isomorphisms

$$\begin{aligned} (-)_{\text{indisc}|X,Y}^{\times} : X_{\text{indisc}} \times Y_{\text{indisc}} &\xrightarrow{\cong} (X \times Y)_{\text{indisc}}, \\ (-)_{\text{indisc}|\mathbb{1}}^{\times} : \text{pt} &\xrightarrow{\cong} \text{pt}_{\text{indisc}}, \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

PROOF 8.2.4.3 ► PROOF OF PROPOSITION 8.2.4.2

Item 1: Functoriality

Clear.

Item 2: Adjointness

This is proved in **Proposition 8.2.1.1**.

Item 3: Symmetric Strong Monoidality With Respect to Products

Clear. 

00Z1 **8.3 Groupoids**

00Z2 **8.3.1 Foundations**

Let C be a category.

00Z3 **DEFINITION 8.3.1.1 ► ISOMORPHISMS**

A morphism $f: A \rightarrow B$ of C is an **isomorphism** if there exists a morphism $f^{-1}: B \rightarrow A$ of C such that

$$\begin{aligned} f \circ f^{-1} &= \text{id}_B, \\ f^{-1} \circ f &= \text{id}_A. \end{aligned}$$

00Z4 **NOTATION 8.3.1.2 ► THE SET OF ISOMORPHISMS BETWEEN TWO OBJECTS IN A CATEGORY**

We write $\text{Iso}_C(A, B)$ for the set of all isomorphisms in C from A to B .

00Z5 **DEFINITION 8.3.1.3 ► GROUPOIDS**

A **groupoid** is a category in which every morphism is an isomorphism.

00Z6 **8.3.2 The Groupoid Completion of a Category**

Let C be a category.

00Z7 **DEFINITION 8.3.2.1 ► THE GROUPOID COMPLETION OF A CATEGORY**

The **groupoid completion of C^1** is the pair $(K_0(C), \iota_C)$ consisting of

- A groupoid $K_0(C)$;
- A functor $\iota_C: C \rightarrow K_0(C)$;

satisfying the following universal property:²

(UP) Given another such pair (\mathcal{G}, i) , there exists a unique functor $K_0(C) \xrightarrow{\exists!} \mathcal{G}$ making the diagram

$$\begin{array}{ccc} & & K_0(C) \\ & \nearrow \iota_C & \downarrow \exists! \\ C & \xrightarrow{i} & \mathcal{G} \end{array}$$

commute.

¹*Further Terminology:* Also called the **Grothendieck groupoid of C** or the **Grothendieck groupoid completion of C** ; see [Item 3 of Proposition 8.3.2.4](#) for an explicit construction.

00Z8

CONSTRUCTION 8.3.2.2 ▶ CONSTRUCTION OF THE GROUPOID COMPLETION OF A CATEGORY

Concretely, the groupoid completion of C is the Gabriel–Zisman localisation $\text{Mor}(C)^{-1}C$ of C at the set $\text{Mor}(C)$ of all morphisms of C ; see ??.
(To be expanded upon later on.)

PROOF 8.3.2.3 ▶ PROOF OF CONSTRUCTION 8.3.2.2

Omitted.

00Z9

PROPOSITION 8.3.2.4 ▶ PROPERTIES OF GROUPOID COMPLETION

Let C be a category.

00ZA

1. *Functoriality.* The assignment $C \mapsto K_0(C)$ defines a functor

$$K_0: \text{Cats} \rightarrow \text{Grpd}.$$

00ZB

2. *2-Functoriality.* The assignment $C \mapsto K_0(C)$ defines a 2-functor

$$K_0: \text{Cats}_2 \rightarrow \text{Grpd}_2.$$

00ZC

3. *Adjointness.* We have an adjunction

$$(K_0 \dashv \iota): \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp \\ \xleftarrow{\iota} \end{array} \text{Grpd},$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Grpd}}(K_0(C), \mathcal{G}) \cong \text{Hom}_{\text{Cats}}(C, \mathcal{G}),$$

natural in $C \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$, forming, together with the functor Core of [Item 1 of Proposition 8.3.3.5](#), a triple adjunction

$$(K_0 \dashv \iota \dashv \text{Core}): \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp \\ \xleftarrow{\iota} \\ \perp \\ \xrightarrow{\text{Core}} \end{array} \text{Grpd},$$

witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\text{Grpd}}(K_0(C), \mathcal{G}) &\cong \text{Hom}_{\text{Cats}}(C, \mathcal{G}), \\ \text{Hom}_{\text{Cats}}(\mathcal{G}, \mathcal{D}) &\cong \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{D})), \end{aligned}$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$.

00ZD

4. *2-Adjointness*. We have a 2-adjunction

$$(K_0 \dashv \iota): \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \xleftarrow{\iota} \\ \xrightarrow{\perp_2} \end{array} \text{Grpd},$$

witnessed by an isomorphism of categories

$$\text{Fun}(K_0(C), \mathcal{G}) \cong \text{Fun}(C, \mathcal{G}),$$

natural in $C \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$, forming, together with the 2-functor Core of **Item 2** of **Proposition 8.3.3.5**, a triple 2-adjunction

$$(K_0 \dashv \iota \dashv \text{Core}): \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \xleftarrow{\iota} \\ \xrightarrow{\perp_2} \\ \xleftarrow{\text{Core}} \end{array} \text{Grpd},$$

witnessed by isomorphisms of categories

$$\begin{aligned} \text{Fun}(K_0(C), \mathcal{G}) &\cong \text{Fun}(C, \mathcal{G}), \\ \text{Fun}(\mathcal{G}, \mathcal{D}) &\cong \text{Fun}(\mathcal{G}, \text{Core}(\mathcal{D})), \end{aligned}$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$.

00ZE

5. *Interaction With Classifying Spaces*. We have an isomorphism of groupoids

$$K_0(C) \cong \Pi_{\leq 1}(|N_\bullet(C)|),$$

natural in $C \in \text{Obj}(\text{Cats})$; i.e. the diagram

$$\begin{array}{ccc} \text{Cats} & \xrightarrow{K_0} & \text{Grp} \\ \downarrow N_\bullet & \hat{\cong} & \uparrow \Pi_{\leq 1} \\ \text{sSets} & \xrightarrow{|\cdot|} & \text{Top} \end{array}$$

commutes up to natural isomorphism.

00ZF

6. *Symmetric Strong Monoidality With Respect to Coproducts.* The groupoid completion functor of **Item 1** has a symmetric strong monoidal structure

$$\left(K_0, K_0^{\amalg}, K_{0|\mathbb{1}}^{\amalg} \right) : (\text{Cats}, \amalg, \emptyset_{\text{cat}}) \rightarrow (\text{Grpd}, \amalg, \emptyset_{\text{cat}})$$

being equipped with isomorphisms

$$\begin{aligned} K_{0|C, \mathcal{D}}^{\amalg} : K_0(C) \amalg K_0(\mathcal{D}) &\xrightarrow{\cong} K_0(C \amalg \mathcal{D}), \\ K_{0|\mathbb{1}}^{\amalg} : \emptyset_{\text{cat}} &\xrightarrow{\cong} K_0(\emptyset_{\text{cat}}), \end{aligned}$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$.

00ZG

7. *Symmetric Strong Monoidality With Respect to Products.* The groupoid completion functor of **Item 1** has a symmetric strong monoidal structure

$$\left(K_0, K_0^{\times}, K_{0|\mathbb{1}}^{\times} \right) : (\text{Cats}, \times, \text{pt}) \rightarrow (\text{Grpd}, \times, \text{pt})$$

being equipped with isomorphisms

$$\begin{aligned} K_{0|C, \mathcal{D}}^{\times} : K_0(C) \times K_0(\mathcal{D}) &\xrightarrow{\cong} K_0(C \times \mathcal{D}), \\ K_{0|\mathbb{1}}^{\times} : \text{pt} &\xrightarrow{\cong} K_0(\text{pt}), \end{aligned}$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$.

PROOF 8.3.2.5 ► PROOF OF PROPOSITION 8.3.2.4

Item 1: Functoriality

Omitted.

Item 2: 2-Functoriality

Omitted.

Item 3: Adjointness

Omitted.

Item 4: 2-Adjointness

Omitted.

Item 5: Interaction With Classifying Spaces

See Corollary 18.33 of <https://web.ma.utexas.edu/users/dafr/M392C-2012/Notes/lecture18.pdf>.

Item 6: Symmetric Strong Monoidality With Respect to Coproducts

Omitted.

Item 7: Symmetric Strong Monoidality With Respect to Products

Omitted.



00ZH 8.3.3 The Core of a Category

Let C be a category.

00ZJ DEFINITION 8.3.3.1 ► THE CORE OF A CATEGORY

The **core** of C is the pair $(\text{Core}(C), \iota_C)$ consisting of

- A groupoid $\text{Core}(C)$;
- A functor $\iota_C: \text{Core}(C) \hookrightarrow C$;

satisfying the following universal property:

(UP) Given another such pair (\mathcal{G}, i) , there exists a unique functor $\mathcal{G} \xrightarrow{\exists!} \text{Core}(C)$ making the diagram

$$\begin{array}{ccc}
 & \text{Core}(C) & \\
 \exists! \nearrow & \uparrow \iota_C & \\
 \mathcal{G} & \xrightarrow{i} & C
 \end{array}$$

commute.

00ZK NOTATION 8.3.3.2 ► ALTERNATIVE NOTATION FOR THE CORE OF A CATEGORY

We also write C^\simeq for $\text{Core}(C)$.

00ZL CONSTRUCTION 8.3.3.3 ► CONSTRUCTION OF THE CORE OF A CATEGORY

The core of C is the wide subcategory of C spanned by the isomorphisms of C , i.e. the category $\text{Core}(C)$ where¹


1. *Objects.* We have

$$\text{Obj}(\text{Core}(C)) \stackrel{\text{def}}{=} \text{Obj}(C).$$

2. *Morphisms.* The morphisms of $\text{Core}(C)$ are the isomorphisms of C .

¹*Slogan:* The groupoid $\text{Core}(C)$ is the maximal subgroupoid of C .

PROOF 8.3.3.4 ► PROOF OF CONSTRUCTION 8.3.3.3

This follows from the fact that functors preserve isomorphisms (Item 1 of Proposition 8.4.1.8). 

00ZM PROPOSITION 8.3.3.5 ► PROPERTIES OF THE CORE OF A CATEGORY

Let C be a category.

00ZN 1. *Functoriality.* The assignment $C \mapsto \text{Core}(C)$ defines a functor

$$\text{Core}: \text{Cats} \rightarrow \text{Grpd}.$$

00ZP 2. *2-Functoriality.* The assignment $C \mapsto \text{Core}(C)$ defines a 2-functor

$$\text{Core}: \text{Cats}_2 \rightarrow \text{Grpd}_2.$$

00ZQ 3. *Adjointness.* We have an adjunction

$$(\iota \dashv \text{Core}): \text{Grpd} \begin{array}{c} \xrightarrow{\iota} \\ \perp \\ \xleftarrow{\text{Core}} \end{array} \text{Cats},$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Cats}}(\mathcal{G}, \mathcal{D}) \cong \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in $\mathcal{G} \in \text{Obj}(\text{Grpd})$ and $\mathcal{D} \in \text{Obj}(\text{Cats})$, forming, together with the functor K_0 of Item 1 of Proposition 8.3.2.4, a triple adjunction

$$(K_0 \dashv \iota \dashv \text{Core}): \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp \\ \xleftarrow{\iota} \\ \perp \\ \xrightarrow{\text{Core}} \end{array} \text{Grpd},$$

witnessed by bijections of sets

$$\text{Hom}_{\text{Grpd}}(K_0(C), \mathcal{G}) \cong \text{Hom}_{\text{Cats}}(C, \mathcal{G}),$$

$$\text{Hom}_{\text{Cats}}(\mathcal{G}, \mathcal{D}) \cong \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$.

00ZR

4. *2-Adjointness.* We have an adjunction

$$(\iota \dashv \text{Core}): \text{Grpd} \begin{array}{c} \xrightarrow{\iota} \\ \dashv_2 \\ \xleftarrow{\text{Core}} \end{array} \text{Cats},$$

witnessed by an isomorphism of categories

$$\text{Fun}(\mathcal{G}, \mathcal{D}) \cong \text{Fun}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in $\mathcal{G} \in \text{Obj}(\text{Grpd})$ and $\mathcal{D} \in \text{Obj}(\text{Cats})$, forming, together with the 2-functor K_0 of [Item 2 of Proposition 8.3.2.4](#), a triple 2-adjunction

$$(\text{K}_0 \dashv \iota \dashv \text{Core}): \text{Cats} \begin{array}{c} \xrightarrow{\text{K}_0} \\ \dashv_2 \\ \xleftarrow{\iota} \\ \dashv_2 \\ \xrightarrow{\text{Core}} \end{array} \text{Grpd},$$

witnessed by isomorphisms of categories

$$\text{Fun}(\text{K}_0(\mathcal{C}), \mathcal{G}) \cong \text{Fun}(\mathcal{C}, \mathcal{G}),$$

$$\text{Fun}(\mathcal{G}, \mathcal{D}) \cong \text{Fun}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in $\mathcal{C}, \mathcal{D} \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$.

00ZS

5. *Symmetric Strong Monoidality With Respect to Products.* The core functor of [Item 1](#) has a symmetric strong monoidal structure

$$(\text{Core}, \text{Core}^\times, \text{Core}_{\mathbb{1}}^\times): (\text{Cats}, \times, \text{pt}) \rightarrow (\text{Grpd}, \times, \text{pt})$$

being equipped with isomorphisms

$$\text{Core}_{\mathcal{C}, \mathcal{D}}^\times: \text{Core}(\mathcal{C}) \times \text{Core}(\mathcal{D}) \xrightarrow{\cong} \text{Core}(\mathcal{C} \times \mathcal{D}),$$

$$\text{Core}_{\mathbb{1}}^\times: \text{pt} \xrightarrow{\cong} \text{Core}(\text{pt}),$$

natural in $\mathcal{C}, \mathcal{D} \in \text{Obj}(\text{Cats})$.

00ZT

6. *Symmetric Strong Monoidality With Respect to Coproducts.* The core functor of [Item 1](#) has a symmetric strong monoidal structure

$$(\text{Core}, \text{Core}^{\amalg}, \text{Core}_{\mathbb{1}}^{\amalg}): (\text{Cats}, \amalg, \emptyset_{\text{cat}}) \rightarrow (\text{Grpd}, \amalg, \emptyset_{\text{cat}})$$

being equipped with isomorphisms

$$\text{Core}_{\mathcal{C}, \mathcal{D}}^{\amalg}: \text{Core}(\mathcal{C}) \amalg \text{Core}(\mathcal{D}) \xrightarrow{\cong} \text{Core}(\mathcal{C} \amalg \mathcal{D}),$$

$$\text{Core}_{\mathbb{1}}^{\amalg}: \emptyset_{\text{cat}} \xrightarrow{\cong} \text{Core}(\emptyset_{\text{cat}}),$$

natural in $\mathcal{C}, \mathcal{D} \in \text{Obj}(\text{Cats})$.

PROOF 8.3.3.6 ► PROOF OF PROPOSITION 8.3.3.5

Item 1: Functoriality

Omitted.

Item 2: 2-Functoriality

Omitted.

Item 3: Adjointness

Omitted.

Item 4: 2-Adjointness

Omitted.

Item 5: Symmetric Strong Monoidality With Respect to Products

Omitted.

Item 6: Symmetric Strong Monoidality With Respect to Coproducts

Omitted. 00ZU **8.4 Functors**00ZV **8.4.1 Foundations**Let \mathcal{C} and \mathcal{D} be categories.00ZW **DEFINITION 8.4.1.1 ► FUNCTORS**A **functor** $F: \mathcal{C} \rightarrow \mathcal{D}$ **from \mathcal{C} to \mathcal{D}** ¹ consists of:

1. *Action on Objects.* A map of sets

$$F: \text{Obj}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{D}),$$

called the **action on objects of F** .

2. *Action on Morphisms.* For each $A, B \in \text{Obj}(\mathcal{C})$, a map

$$F_{A,B}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F(A), F(B)),$$

called the **action on morphisms of F at (A, B)** ².

satisfying the following conditions:

1. *Preservation of Identities.* For each $A \in \text{Obj}(C)$, the diagram

$$\begin{array}{ccc} \text{pt} & & \\ \mathbb{1}_A^C \downarrow & \searrow \mathbb{1}_{F(A)}^{\mathcal{D}} & \\ \text{Hom}_C(A, A) & \xrightarrow{F_{A,A}} & \text{Hom}_{\mathcal{D}}(F(A), F(A)) \end{array}$$

commutes, i.e. we have

$$F(\text{id}_A) = \text{id}_{F(A)}.$$

2. *Preservation of Composition.* For each $A, B, C \in \text{Obj}(C)$, the diagram

$$\begin{array}{ccc} \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,C}^C} & \text{Hom}_C(A, C) \\ \downarrow F_{B,C} \times F_{A,B} & & \downarrow F_{A,C} \\ \text{Hom}_{\mathcal{D}}(F(B), F(C)) \times \text{Hom}_{\mathcal{D}}(F(A), F(B)) & \xrightarrow{\circ_{F(A),F(B),F(C)}^{\mathcal{D}}} & \text{Hom}_{\mathcal{D}}(F(A), F(C)) \end{array}$$

commutes, i.e. for each composable pair (g, f) of morphisms of C , we have

$$F(g \circ f) = F(g) \circ F(f).$$

¹Further Terminology: Also called a **covariant functor**.

²Further Terminology: Also called **action on Hom-sets of F at (A, B)** .

00ZX NOTATION 8.4.1.2 ► SUBSCRIPT AND SUPERScript NOTATION FOR FUNCTORS

Let C and \mathcal{D} be categories, and write C^{op} for the opposite category of C of ??.

00ZY 1. Given a functor

$$F: C \rightarrow \mathcal{D},$$

we also write F_A for $F(A)$.

00ZZ 2. Given a functor

$$F: C^{\text{op}} \rightarrow \mathcal{D},$$

we also write F^A for $F(A)$.

0100 3. Given a functor

$$F: C \times C \rightarrow \mathcal{D},$$

we also write $F_{A,B}$ for $F(A, B)$.

0101 4. Given a functor

$$F: \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{D},$$

we also write F_B^A for $F(A, B)$.

We employ a similar notation for morphisms, writing e.g. F_f for $F(f)$ given a functor $F: \mathcal{C} \rightarrow \mathcal{D}$.

0102 NOTATION 8.4.1.3 ► ADDITIONAL NOTATION FOR FUNCTORS

Following the notation $\llbracket x \mapsto f(x) \rrbracket$ for a function $f: X \rightarrow Y$ introduced in Notation 1.1.1.2, we will sometimes denote a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ by

$$F \stackrel{\text{def}}{=} \llbracket A \mapsto F(A) \rrbracket,$$

specially when the action on morphisms of F is clear from its action on objects.

0103 EXAMPLE 8.4.1.4 ► IDENTITY FUNCTORS

The **identity functor** of a category \mathcal{C} is the functor $\text{id}_{\mathcal{C}}: \mathcal{C} \rightarrow \mathcal{C}$ where

1. *Action on Objects.* For each $A \in \text{Obj}(\mathcal{C})$, we have

$$\text{id}_{\mathcal{C}}(A) \stackrel{\text{def}}{=} A.$$

2. *Action on Morphisms.* For each $A, B \in \text{Obj}(\mathcal{C})$, the action on morphisms

$$(\text{id}_{\mathcal{C}})_{A,B}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \underbrace{\text{Hom}_{\mathcal{C}}(\text{id}_{\mathcal{C}}(A), \text{id}_{\mathcal{C}}(B))}_{\stackrel{\text{def}}{=} \text{Hom}_{\mathcal{C}}(A, B)}$$

of $\text{id}_{\mathcal{C}}$ at (A, B) is defined by

$$(\text{id}_{\mathcal{C}})_{A,B} \stackrel{\text{def}}{=} \text{id}_{\text{Hom}_{\mathcal{C}}(A, B)}.$$

PROOF 8.4.1.5 ► PROOF OF EXAMPLE 8.4.1.4


Preservation of Identities

We have $\text{id}_{\mathcal{C}}(\text{id}_A) \stackrel{\text{def}}{=} \text{id}_A$ for each $A \in \text{Obj}(\mathcal{C})$ by definition.

Preservation of Compositions

For each composable pair $A \xrightarrow{f} B \xrightarrow{g} B$ of morphisms of C , we have

$$\begin{aligned} \text{id}_C(g \circ f) &\stackrel{\text{def}}{=} g \circ f \\ &\stackrel{\text{def}}{=} \text{id}_C(g) \circ \text{id}_C(f). \end{aligned}$$

This finishes the proof. 

0104

DEFINITION 8.4.1.6 ► COMPOSITION OF FUNCTORS

The **composition** of two functors $F: C \rightarrow D$ and $G: D \rightarrow E$ is the functor $G \circ F$ where

- *Action on Objects.* For each $A \in \text{Obj}(C)$, we have

$$[G \circ F](A) \stackrel{\text{def}}{=} G(F(A)).$$

- *Action on Morphisms.* For each $A, B \in \text{Obj}(C)$, the action on morphisms

$$(G \circ F)_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_E(G_{F_A}, G_{F_B})$$

of $G \circ F$ at (A, B) is defined by

$$[G \circ F](f) \stackrel{\text{def}}{=} G(F(f)).$$

PROOF 8.4.1.7 ► PROOF OF DEFINITION 8.4.1.6

Preservation of Identities

For each $A \in \text{Obj}(C)$, we have

$$\begin{aligned} G_{F_{\text{id}_A}} &= G_{\text{id}_{F_A}} && \text{(functoriality of } F) \\ &= \text{id}_{G_{F_A}}. && \text{(functoriality of } G) \end{aligned}$$

Preservation of Composition

For each composable pair (g, f) of morphisms of C , we have

$$\begin{aligned} G_{F_{g \circ f}} &= G_{F_g \circ F_f} && \text{(functoriality of } F) \\ &= G_{F_g} \circ G_{F_f}. && \text{(functoriality of } G) \end{aligned}$$

This finishes the proof. 

0105 PROPOSITION 8.4.1.8 ► ELEMENTARY PROPERTIES OF FUNCTORS

Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

- 0106 1. *Preservation of Isomorphisms.* If f is an isomorphism in \mathcal{C} , then $F(f)$ is an isomorphism in \mathcal{D} .¹

¹When the converse holds, we call F *conservative*, see Definition 8.5.4.1.

PROOF 8.4.1.9 ► PROOF OF PROPOSITION 8.4.1.8


Item 1: Preservation of Isomorphisms

Indeed, we have

$$\begin{aligned} F(f)^{-1} \circ F(f) &= F(f^{-1} \circ f) \\ &= F(\text{id}_A) \\ &= \text{id}_{F(A)} \end{aligned}$$

and

$$\begin{aligned} F(f) \circ F(f)^{-1} &= F(f \circ f^{-1}) \\ &= F(\text{id}_B) \\ &= \text{id}_{F(B)}, \end{aligned}$$

showing $F(f)$ to be an isomorphism. 

0107 8.4.2 Contravariant Functors

Let \mathcal{C} and \mathcal{D} be categories, and let \mathcal{C}^{op} denote the opposite category of \mathcal{C} of ??.

0108 DEFINITION 8.4.2.1 ► CONTRAVARIANT FUNCTORS

A **contravariant functor** from \mathcal{C} to \mathcal{D} is a functor from \mathcal{C}^{op} to \mathcal{D} .

0109 REMARK 8.4.2.2 ► UNWINDING DEFINITION 8.4.2.1

In detail, a **contravariant functor** from \mathcal{C} to \mathcal{D} consists of:

1. *Action on Objects.* A map of sets

$$F: \text{Obj}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{D}),$$

called the **action on objects of F** .

2. *Action on Morphisms.* For each $A, B \in \text{Obj}(C)$, a map

$$F_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F(B), F(A)),$$

called the **action on morphisms of F at (A, B)** .

satisfying the following conditions:

1. *Preservation of Identities.* For each $A \in \text{Obj}(C)$, the diagram

$$\begin{array}{ccc} \text{pt} & & \\ \downarrow \mathbb{1}_A^C & \searrow \mathbb{1}_{F(A)}^{\mathcal{D}} & \\ \text{Hom}_C(A, A) & \xrightarrow{F_{A,A}} & \text{Hom}_{\mathcal{D}}(F(A), F(A)) \end{array}$$

commutes, i.e. we have

$$F(\text{id}_A) = \text{id}_{F(A)}.$$

2. *Preservation of Composition.* For each $A, B, C \in \text{Obj}(C)$, the diagram

$$\begin{array}{ccc} & \text{Hom}_{\mathcal{D}}(F(C), F(B)) \times \text{Hom}_{\mathcal{D}}(F(B), F(A)) & \\ & \nearrow F_{B,C} \times F_{A,B} & \searrow \sigma_{\text{Hom}_{\mathcal{D}}(F(C), F(B)), \text{Hom}_{\mathcal{D}}(F(B), F(A))}^{\text{Sets}} \\ \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & & \text{Hom}_{\mathcal{D}}(F(B), F(A)) \times \text{Hom}_{\mathcal{D}}(F(C), F(B)) \\ \downarrow \circ_{A,B,C}^C & & \downarrow \circ_{F(C), F(B), F(A)}^{\mathcal{D}} \\ \text{Hom}_C(A, C) & \xrightarrow{F_{A,C}} & \text{Hom}_{\mathcal{D}}(F(C), F(A)) \end{array}$$

commutes, i.e. for each composable pair (g, f) of morphisms of C , we have

$$F(g \circ f) = F(f) \circ F(g).$$

010A **REMARK 8.4.2.3 ► ON THE TERM CONTRAVARIANT FUNCTOR**

Throughout this work we will not use the term “contravariant” functor, speaking instead simply of functors $F: \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}$. We will usually, however, write

$$F_{A,B}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F(B), F(A))$$

for the action on morphisms

$$F_{A,B}: \text{Hom}_{\mathcal{C}^{\text{op}}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F(A), F(B))$$

of F , as well as write $F(g \circ f) = F(f) \circ F(g)$.

010B **8.4.3 Forgetful Functors**

010C **DEFINITION 8.4.3.1 ► FORGETFUL FUNCTORS**

There isn't a precise definition of a **forgetful functor**.

010D **REMARK 8.4.3.2 ► UNWINDING DEFINITION 8.4.3.1**

Despite there not being a formal or precise definition of a forgetful functor, the term is often very useful in practice, similarly to the word “canonical”. The idea is that a “forgetful functor” is a functor that forgets structure or properties, and is best explained through examples, such as the ones below (see [Examples 8.4.3.3](#) and [8.4.3.4](#)).

010E **EXAMPLE 8.4.3.3 ► FORGETFUL FUNCTORS THAT FORGET STRUCTURE**

Examples of forgetful functors that forget structure include:

- 010F
1. *Forgetting Group Structures.* The functor $\text{Grp} \rightarrow \text{Sets}$ sending a group (G, μ_G, η_G) to its underlying set G , forgetting the multiplication and unit maps μ_G and η_G of G .
- 010G
2. *Forgetting Topologies.* The functor $\text{Top} \rightarrow \text{Sets}$ sending a topological space (X, \mathcal{T}_X) to its underlying set X , forgetting the topology \mathcal{T}_X .
- 010H
3. *Forgetting Fibrations.* The functor $\text{FibSets}(K) \rightarrow \text{Sets}$ sending a K -fibred set $\phi_X: X \rightarrow K$ to the set X , forgetting the map ϕ_X and the base set K .

010J EXAMPLE 8.4.3.4 ► FORGETFUL FUNCTORS THAT FORGET PROPERTIES

Examples of forgetful functors that forget properties include:

- 010K 1. *Forgetting Commutativity.* The inclusion functor $\iota: \text{CMon} \hookrightarrow \text{Mon}$ which forgets the property of being commutative.
- 010L 2. *Forgetting Inverses.* The inclusion functor $\iota: \text{Grp} \hookrightarrow \text{Mon}$ which forgets the property of having inverses.

010M NOTATION 8.4.3.5 ► NOTATION FOR FORGETFUL FUNCTORS THAT FORGET STRUCTURE

Throughout this work, we will denote forgetful functors that forget structure by 忘, e.g. as in

$$\text{忘}: \text{Grp} \rightarrow \text{Sets}.$$

The symbol 忘, pronounced *wasureru* (see [Item 1](#) of [Remark 8.4.3.6](#) below), means *to forget*, and is a kanji found in the following words in Japanese and Chinese:

- 010N 1. 忘れる, transcribed as *wasureru*, meaning *to forget*.
- 010P 2. 忘却関手, transcribed as *boukyaku kanshu*, meaning *forgetful functor*.
- 010Q 3. 忘记 or 忘記, transcribed as *wàngjì*, meaning *to forget*.
- 010R 4. 遗忘函子 or 遺忘函子, transcribed as *yíwàng hánzǐ*, meaning *forgetful functor*.

010S REMARK 8.4.3.6 ► PRONUNCIATION OF THE WORDS IN NOTATION 8.4.3.5

Here we collect the pronunciation of the words in [Notation 8.4.3.5](#) for accuracy and completeness.

- 010T 1. Pronunciation of 忘れる:
- Audio: see <https://topological-modular-forms.github.io/the-clowder-project/static/sounds/wasureru-01.mp3>
 - IPA broad transcription: [wäsũr̥ɕru].
 - IPA narrow transcription: [ɥ̥ʰäsi̯r̥ɕɾ̥ɥ̥ʰ].
- 010U 2. Pronunciation of 忘却関手: Pronunciation:

010V

3. Pronunciation of 忘记:

- Audio: see <https://topological-modular-forms.github.io/the-clowder-project/static/sounds/wasureru-02.mp3>
- IPA broad transcription: [bɔ:kʰäku kǎũɕu].
- IPA narrow transcription: [bɔ:kʰäku^β kǎũɕu^β].

010W

4. Pronunciation of 遗忘函子:

- Audio: see <https://topological-modular-forms.github.io/the-clowder-project/static/sounds/wasureru-03.ogg>
- Broad IPA transcription: [waŋtɕi].
- Sinological IPA transcription: [waŋ⁵¹⁻⁵³tɕi⁵¹].

- Audio: see <https://topological-modular-forms.github.io/the-clowder-project/static/sounds/wasureru-04.mp3>
- Broad IPA transcription: [iwaŋ xǎntɕzi].
- Sinological IPA transcription: [i³⁵waŋ⁵¹ xǎn³⁵tɕz²¹⁴⁻²¹⁽⁴⁾].

010X 8.4.4 The Natural Transformation Associated to a Functor

010Y

DEFINITION 8.4.4.1 ► THE NATURAL TRANSFORMATION ASSOCIATED TO A FUNCTOR

Every functor $F: C \rightarrow D$ defines a natural transformation¹

$$F^\dagger: \text{Hom}_C \Longrightarrow \text{Hom}_D \circ (F^{\text{op}} \times F),$$

called the **natural transformation associated to F** , consisting of the collection

$$\left\{ F_{A,B}^\dagger: \text{Hom}_C(A, B) \rightarrow \text{Hom}_D(F_A, F_B) \right\}_{(A,B) \in \text{Obj}(C^{\text{op}} \times C)}$$

with

$$F_{A,B}^\dagger \stackrel{\text{def}}{=} F_{A,B}.$$

¹This is the 1-categorical version of [Item 1](#) of [Proposition 2.4.1.3](#).

PROOF 8.4.4.2 ► PROOF OF DEFINITION 8.4.4.1

The naturality condition for F^\dagger is the requirement that for each morphism

$$(\phi, \psi): (X, Y) \rightarrow (A, B)$$

of $C^{\text{op}} \times C$, the diagram

$$\begin{array}{ccc} \text{Hom}_C(X, Y) & \xrightarrow{\phi^* \circ \psi_* = \psi_* \circ \phi^*} & \text{Hom}_C(A, B) \\ F_{X,Y} \downarrow & & \downarrow F_{A,B} \\ \text{Hom}_{\mathcal{D}}(F_X, F_Y) & \xrightarrow{F(\phi)^* \circ F(\psi)_* = F(\psi)_* \circ F(\phi)^*} & \text{Hom}_{\mathcal{D}}(F_A, F_B), \end{array}$$

acting on elements as

$$\begin{array}{ccc} f & \longmapsto & \psi \circ f \circ \phi \\ \downarrow & & \downarrow \\ F(f) & \longmapsto & F(\psi) \circ F(f) \circ F(\psi) = F(\psi \circ f \circ \phi) \end{array}$$

commutes, which follows from the functoriality of F . ▢

PROPOSITION 8.4.4.3 ► PROPERTIES OF NATURAL TRANSFORMATIONS ASSOCIATED TO FUNCTORS

010Z

Let $F: C \rightarrow D$ and $G: D \rightarrow E$ be functors.

0110

1. *Interaction With Natural Isomorphisms.* The following conditions are equivalent:

0111

(a) The natural transformation $F^\dagger: \text{Hom}_C \Rightarrow \text{Hom}_D \circ (F^{\text{op}} \times F)$ associated to F is a natural isomorphism.

0112

(b) The functor F is fully faithful.

0113

2. *Interaction With Composition.* We have an equality of pasting diagrams

$$\begin{array}{ccc} C^{\text{op}} \times C & \xrightarrow{F^{\text{op}} \times F} & D^{\text{op}} \times D & \xrightarrow{G^{\text{op}} \times G} & E^{\text{op}} \times E & & C^{\text{op}} \times C & \xrightarrow{(GoF)^{\text{op}} \times (GoF)} & E^{\text{op}} \times E, \\ \downarrow \text{Hom}_C & \nearrow F^\dagger & \downarrow \text{Hom}_D & \nearrow G^\dagger & \downarrow \text{Hom}_E & & \downarrow \text{Hom}_C & \nearrow (GoF)^\dagger & \downarrow \text{Hom}_E \\ & & \text{Sets} & & & = & & & \text{Sets} \end{array}$$

in \mathbf{Cats}_2 , i.e. we have

$$(G \circ F)^\dagger = (G^\dagger \star \text{id}_{F^{\text{op}} \times F}) \circ F^\dagger.$$

0114

3. *Interaction With Identities.* We have

$$\text{id}_C^\dagger = \text{id}_{\text{Hom}_C(-1, -2)},$$

i.e. the natural transformation associated to id_C is the identity natural transformation of the functor $\text{Hom}_C(-1, -2)$.

PROOF 8.4.4.4 ► PROOF OF PROPOSITION 8.4.4.3

Item 1: Interaction With Natural Isomorphisms

Clear.

Item 2: Interaction With Composition

Clear.

Item 3: Interaction With Identities

Clear. 

0115 8.5 Conditions on Functors

0116 8.5.1 Faithful Functors

Let \mathcal{C} and \mathcal{D} be categories.

0117 DEFINITION 8.5.1.1 ► FAITHFUL FUNCTORS

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **faithful** if, for each $A, B \in \text{Obj}(\mathcal{C})$, the action on morphisms

$$F_{A,B}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of F at (A, B) is injective.

0118 PROPOSITION 8.5.1.2 ► PROPERTIES OF FAITHFUL FUNCTORS

Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

0119 1. *Interaction With Postcomposition.* The following conditions are equivalent:

- 011A (a) The functor $F: C \rightarrow \mathcal{D}$ is faithful.
- 011B (b) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor
- $$F_*: \text{Fun}(\mathcal{X}, C) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$
- is faithful.
- 011C (c) The functor $F: C \rightarrow \mathcal{D}$ is a representably faithful morphism in Cats_2 in the sense of [Definition 9.1.1.1](#).
- 011D 2. *Interaction With Precomposition I.* Let $F: C \rightarrow \mathcal{D}$ be a functor.
- 011E (a) If F is faithful, then the precomposition functor
- $$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$
- can fail* to be faithful.
- 011F (b) Conversely, if the precomposition functor
- $$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$
- is faithful, then F *can fail* to be faithful.
- 011G 3. *Interaction With Precomposition II.* If F is essentially surjective, then the precomposition functor
- $$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$
- is faithful.
- 011H 4. *Interaction With Precomposition III.* The following conditions are equivalent:
- 011J (a) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor
- $$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$
- is faithful.
- 011K (b) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor
- $$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$
- is conservative.

011L

(c) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^* : \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is monadic.

011M

(d) The functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is a corepresentably faithful morphism in Cats_2 in the sense of [Definition 9.2.1.1](#).

011N

(e) The components

$$\eta_G : G \Longrightarrow \text{Ran}_F(G \circ F)$$

of the unit

$$\eta : \text{id}_{\text{Fun}(\mathcal{D}, \mathcal{X})} \Longrightarrow \text{Ran}_F \circ F^*$$

of the adjunction $F^* \dashv \text{Ran}_F$ are all monomorphisms.

011P

(f) The components

$$\epsilon_G : \text{Lan}_F(G \circ F) \Longrightarrow G$$

of the counit

$$\epsilon : \text{Lan}_F \circ F^* \Longrightarrow \text{id}_{\text{Fun}(\mathcal{D}, \mathcal{X})}$$

of the adjunction $\text{Lan}_F \dashv F^*$ are all epimorphisms.

011Q

(g) The functor F is dominant ([Definition 8.6.1.1](#)), i.e. every object of \mathcal{D} is a retract of some object in $\text{Im}(F)$:(★) For each $B \in \text{Obj}(\mathcal{D})$, there exist:

- An object A of \mathcal{C} ;
- A morphism $s : B \rightarrow F(A)$ of \mathcal{D} ;
- A morphism $r : F(A) \rightarrow B$ of \mathcal{D} ;

such that $r \circ s = \text{id}_B$.**PROOF 8.5.1.3 ► PROOF OF PROPOSITION 8.5.1.2**

Item 1: Interaction With Postcomposition

Omitted.

Item 2: Interaction With Precomposition I

See [[MSE 733163](#)] for [Item 2a](#). [Item 2b](#) follows from [Item 3](#) and the fact that

there are essentially surjective functors that are not faithful.

Item 3: Interaction With Precomposition II

Omitted, but see https://unimath.github.io/doc/UniMath/d4de26f//UniMath.CategoryTheory.precomp_fully_faithful.html for a formalised proof.

Item 4: Interaction With Precomposition III

We claim [Items 4a](#) to [4g](#) are equivalent:

- [Items 4a](#) and [4d](#) Are Equivalent: This is true by the definition of corepresentably faithful morphism; see [Definition 9.2.1.1](#).
- [Items 4a](#) to [4c](#) and [4g](#) Are Equivalent: See [[Adá+01](#), Proposition 4.1] or alternatively [[Fre09](#), Lemmas 3.1 and 3.2] for the equivalence between [Items 4a](#) and [4g](#).
- [Items 4a](#), [4e](#) and [4f](#) Are Equivalent: See ?? of ??.

This finishes the proof. 

011R 8.5.2 Full Functors

Let \mathcal{C} and \mathcal{D} be categories.

011S DEFINITION 8.5.2.1 ► FULL FUNCTORS

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **full** if, for each $A, B \in \text{Obj}(\mathcal{C})$, the action on morphisms

$$F_{A,B}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of F at (A, B) is surjective.

011T PROPOSITION 8.5.2.2 ► PROPERTIES OF FULL FUNCTORS

Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

011U 1. *Interaction With Postcomposition.* The following conditions are equivalent:

011V (a) The functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is full.

011W (b) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is full.

011X (c) The functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is a representably full morphism in \mathbf{Cats}_2 in the sense of [Definition 9.1.2.1](#).

011Y 2. *Interaction With Precomposition I.* If F is full, then the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

can fail to be full.

011Z 3. *Interaction With Precomposition II.* If the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is full, then F can fail to be full.

0120 4. *Interaction With Precomposition III.* If F is essentially surjective and full, then the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is full (and also faithful by [Item 3](#) of [Proposition 8.5.1.2](#)).

0121 5. *Interaction With Precomposition IV.* The following conditions are equivalent:

0122 (a) For each $\mathcal{X} \in \text{Obj}(\mathbf{Cats})$, the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is full.

0123 (b) The functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is a corepresentably full morphism in \mathbf{Cats}_2 in the sense of [Definition 9.2.1.1](#).

0124 (c) The components

$$\eta_G: G \Longrightarrow \text{Ran}_F(G \circ F)$$

of the unit

$$\eta: \text{id}_{\text{Fun}(\mathcal{D}, \mathcal{X})} \Longrightarrow \text{Ran}_F \circ F^*$$

of the adjunction $F^* \dashv \text{Ran}_F$ are all retractions/split epimorphisms.

0125

(d) The components

$$\epsilon_G : \text{Lan}_F(G \circ F) \Longrightarrow G$$

of the counit

$$\epsilon : \text{Lan}_F \circ F^* \Longrightarrow \text{id}_{\text{Fun}(\mathcal{D}, \mathcal{X})}$$

of the adjunction $\text{Lan}_F \dashv F^*$ are all sections/split monomorphisms.

0126

(e) For each $B \in \text{Obj}(\mathcal{D})$, there exist:

- An object A_B of \mathcal{C} ;
- A morphism $s_B : B \rightarrow F(A_B)$ of \mathcal{D} ;
- A morphism $r_B : F(A_B) \rightarrow B$ of \mathcal{D} ;

satisfying the following condition:

(★) For each $A \in \text{Obj}(\mathcal{C})$ and each pair of morphisms

$$\begin{aligned} r : F(A) &\rightarrow B, \\ s : B &\rightarrow F(A) \end{aligned}$$

of \mathcal{D} , we have

$$[(A_B, s_B, r_B)] = [(A, s, r \circ s_B \circ r_B)]$$

$$\text{in } \int^{A \in \mathcal{C}} h_{F_A}^{B'} \times h_B^{F_A}.$$

PROOF 8.5.2.3 ► PROOF OF PROPOSITION 8.5.2.2

Item 1: Interaction With Postcomposition

Omitted.

Item 2: Interaction With Precomposition I

Omitted.

Item 3: Interaction With Precomposition II

See [BS10, p. 47].

Item 4: Interaction With Precomposition III


Omitted, but see https://unimath.github.io/doc/UniMath/d4de26f//UniMath.CategoryTheory.precomp_fully_faithful.html for a

formalised proof.

Item 5: Interaction With Precomposition IV

We claim **Items 5a to 5e** are equivalent:

- **Items 5a and 5b Are Equivalent:** This is true by the definition of corepresentably full morphism; see **Definition 9.2.2.1**.
- **Items 5a, 5c and 5d Are Equivalent:** See ?? of ??.
- **Items 5a and 5e Are Equivalent:** See [Adá+01, Item (b) of Remark 4.3].

This finishes the proof. 

0127

QUESTION 8.5.2.4 ► BETTER CHARACTERISATIONS OF FUNCTORS WITH FULL PRECOMPOSITION

Item 5 of **Proposition 8.5.2.2** gives a characterisation of the functors F for which F^* is full, but the characterisations given there are really messy. Are there better ones?

This question also appears as [MO 468121b].

0128 8.5.3 Fully Faithful Functors

Let \mathcal{C} and \mathcal{D} be categories.

0129

DEFINITION 8.5.3.1 ► FULLY FAITHFUL FUNCTORS

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **fully faithful** if F is full and faithful, i.e. if, for each $A, B \in \text{Obj}(\mathcal{C})$, the action on morphisms

$$F_{A,B}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of F at (A, B) is bijective.

012A

PROPOSITION 8.5.3.2 ► PROPERTIES OF FULLY FAITHFUL FUNCTORS

Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

012B

1. *Characterisations.* The following conditions are equivalent:

012C

(a) The functor F is fully faithful.

012D

(b) We have a pullback square

$$\begin{array}{ccc}
 \text{Arr}(\mathcal{C}) & \xrightarrow{\text{Arr}(F)} & \text{Arr}(\mathcal{D}) \\
 \text{src} \times \text{tgt} \downarrow & \lrcorner & \downarrow \text{src} \times \text{tgt} \\
 \mathcal{C} \times \mathcal{C} & \xrightarrow{F \times F} & \mathcal{D} \times \mathcal{D}
 \end{array}$$

$\text{Arr}(\mathcal{C}) \cong (\mathcal{C} \times \mathcal{C}) \times_{\mathcal{D} \times \mathcal{D}} \text{Arr}(\mathcal{D}),$

in Cats .

012E

2. *Conservativity*. If F is fully faithful, then F is conservative.

012F

3. *Essential Injectivity*. If F is fully faithful, then F is essentially injective.

012G

4. *Interaction With Co/Limits*. If F is fully faithful, then F reflects co/limits.

012H

5. *Interaction With Postcomposition*. The following conditions are equivalent:

012J

(a) The functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is fully faithful.

012K

(b) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is fully faithful.

012L

(c) The functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is a representably fully faithful morphism in Cats_2 in the sense of [Definition 9.1.3.1](#).

012M

6. *Interaction With Precomposition I*. If F is fully faithful, then the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

can fail to be fully faithful.

012N

7. *Interaction With Precomposition II*. If the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is fully faithful, then F *can fail* to be fully faithful (and in fact it can also fail to be either full or faithful).

- 012P 8. *Interaction With Precomposition III.* If F is essentially surjective and full, then the precomposition functor

$$F^* : \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is fully faithful.

- 012Q 9. *Interaction With Precomposition IV.* The following conditions are equivalent:

- 012R (a) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^* : \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is fully faithful.

- 012S (b) The precomposition functor

$$F^* : \text{Fun}(\mathcal{D}, \text{Sets}) \rightarrow \text{Fun}(\mathcal{C}, \text{Sets})$$

is fully faithful.

- 012T (c) The functor

$$\text{Lan}_F : \text{Fun}(\mathcal{C}, \text{Sets}) \rightarrow \text{Fun}(\mathcal{D}, \text{Sets})$$

is fully faithful.

- 012U (d) The functor F is a corepresentably fully faithful morphism in Cats_2 in the sense of [Definition 9.2.3.1](#).

- 012V (e) The functor F is absolutely dense.

- 012W (f) The components

$$\eta_G : G \Longrightarrow \text{Ran}_F(G \circ F)$$

of the unit

$$\eta : \text{id}_{\text{Fun}(\mathcal{D}, \mathcal{X})} \Longrightarrow \text{Ran}_F \circ F^*$$

of the adjunction $F^* \dashv \text{Ran}_F$ are all isomorphisms.

- 012X (g) The components

$$\epsilon_G : \text{Lan}_F(G \circ F) \Longrightarrow G$$

of the counit

$$\epsilon : \text{Lan}_F \circ F^* \Longrightarrow \text{id}_{\text{Fun}(\mathcal{D}, \mathcal{X})}$$

of the adjunction $\text{Lan}_F \dashv F^*$ are all isomorphisms.

012Y

(h) The natural transformation

$$\alpha: \text{Lan}_{h_F}(h^F) \Longrightarrow h$$

with components

$$\alpha_{B',B}: \int^{A \in C} h_{F_A}^{B'} \times h_B^{F_A} \rightarrow h_B^{B'}$$

given by

$$\alpha_{B',B}([\phi, \psi]) = \psi \circ \phi$$

is a natural isomorphism.

012Z

(i) For each $B \in \text{Obj}(\mathcal{D})$, there exist:

- An object A_B of C ;
- A morphism $s_B: B \rightarrow F(A_B)$ of \mathcal{D} ;
- A morphism $r_B: F(A_B) \rightarrow B$ of \mathcal{D} ;

satisfying the following conditions:

0130

i. The triple $(F(A_B), r_B, s_B)$ is a retract of B , i.e. we have $r_B \circ s_B = \text{id}_B$.

0131

ii. For each morphism $f: B' \rightarrow B$ of \mathcal{D} , we have

$$[(A_B, s_{B'}, f \circ r_{B'})] = [(A_B, s_B \circ f, r_B)]$$

$$\text{in } \int^{A \in C} h_{F_A}^{B'} \times h_B^{F_A}.$$

PROOF 8.5.3.3 ► PROOF OF PROPOSITION 8.5.3.2

Item 1: Characterisations

Omitted.

Item 2: Conservativity

This is a repetition of **Item 2** of **Proposition 8.5.4.2**, and is proved there.

Item 3: Essential Injectivity

Omitted.

Item 4: Interaction With Co/Limits

Omitted.

Item 5: Interaction With Postcomposition

This follows from [Item 1 of Proposition 8.5.1.2](#) and [Item 1 of Proposition 8.5.2.2](#).

Item 6: Interaction With Precomposition I

See [\[MSE 733161\]](#) for an example of a fully faithful functor whose precomposition with which fails to be full.

Item 7: Interaction With Precomposition II

See [\[MSE 749304, Item 3\]](#).


Item 8: Interaction With Precomposition III

Omitted, but see https://unimath.github.io/doc/UniMath/d4de26f//UniMath.CategoryTheory.precomp_fully_faithful.html for a formalised proof.

Item 9: Interaction With Precomposition IV

We claim [Items 9a to 9i](#) are equivalent:

- [Items 9a and 9d](#) Are Equivalent: This is true by the definition of corepresentably fully faithful morphism; see [Definition 9.2.3.1](#).
- [Items 9a, 9f and 9g](#) Are Equivalent: See ?? of ??.
- [Items 9a to 9c](#) Are Equivalent: This follows from [\[Low15, Proposition A.1.5\]](#).
- [Items 9a, 9e, 9h and 9i](#) Are Equivalent: See [\[Fre09, Theorem 4.1\]](#) and [\[Adá+01, Theorem 1.1\]](#).

This finishes the proof. 

0132 8.5.4 Conservative Functors

Let \mathcal{C} and \mathcal{D} be categories.

0133 DEFINITION 8.5.4.1 ► CONSERVATIVE FUNCTORS

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **conservative** if it satisfies the following condition:¹

- (★) For each $f \in \text{Mor}(\mathcal{C})$, if $F(f)$ is an isomorphism in \mathcal{D} , then f is an isomorphism in \mathcal{C} .

¹*Slogan:* A functor F is **conservative** if it reflects isomorphisms.

0134 **PROPOSITION 8.5.4.2 ► PROPERTIES OF CONSERVATIVE FUNCTORS**

Let $F: C \rightarrow \mathcal{D}$ be a functor.

0135 1. *Characterisations.* The following conditions are equivalent:

0136 (a) The functor F is conservative.

0137 (b) For each $f \in \text{Mor}(C)$, the morphism $F(f)$ is an isomorphism in \mathcal{D} iff f is an isomorphism in C .

0138 2. *Interaction With Fully Faithfulness.* Every fully faithful functor is conservative.

0139 3. *Interaction With Precomposition.* The following conditions are equivalent:

013A (a) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$

is conservative.

013B (b) The equivalent conditions of **Item 4** of **Proposition 8.5.1.2** are satisfied.

PROOF 8.5.4.3 ► PROOF OF PROPOSITION 8.5.4.2

Item 1: Characterisations

This follows from **Item 1** of **Proposition 8.4.1.8**.


Item 2: Interaction With Fully Faithfulness

Let $F: C \rightarrow \mathcal{D}$ be a fully faithful functor, let $f: A \rightarrow B$ be a morphism of C , and suppose that F_f is an isomorphism. We have

$$\begin{aligned} F(\text{id}_B) &= \text{id}_{F(B)} \\ &= F(f) \circ F(f)^{-1} \\ &= F(f \circ f^{-1}). \end{aligned}$$

Similarly, $F(\text{id}_A) = F(f^{-1} \circ f)$. But since F is fully faithful, we must have

$$\begin{aligned} f \circ f^{-1} &= \text{id}_B, \\ f^{-1} \circ f &= \text{id}_A, \end{aligned}$$

showing f to be an isomorphism. Thus F is conservative. 

013C QUESTION 8.5.4.4 ► CHARACTERISATIONS OF FUNCTORS WITH CONSERVATIVE PRE/POSTCOMPOSITION

Is there a characterisation of functors $F: \mathcal{C} \rightarrow \mathcal{D}$ satisfying the following condition:

(★) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is conservative?

This question also appears as [MO 468121a].

013D 8.5.5 Essentially Injective Functors

Let \mathcal{C} and \mathcal{D} be categories.

013E DEFINITION 8.5.5.1 ► ESSENTIALLY INJECTIVE FUNCTORS

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **essentially injective** if it satisfies the following condition:

(★) For each $A, B \in \text{Obj}(\mathcal{C})$, if $F(A) \cong F(B)$, then $A \cong B$.

013F QUESTION 8.5.5.2 ► CHARACTERISATIONS OF FUNCTORS WITH ESSENTIALLY INJECTIVE PRE/POSTCOMPOSITION

Is there a characterisation of functors $F: \mathcal{C} \rightarrow \mathcal{D}$ such that:

013G 1. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is essentially injective, i.e. if $\phi \circ F \cong \psi \circ F$, then $\phi \cong \psi$ for all functors ϕ and ψ ?

013H 2. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is essentially injective, i.e. if $F \circ \phi \cong F \circ \psi$, then $\phi \cong \psi$?

This question also appears as [MO 468121a].

013J 8.5.6 Essentially Surjective Functors

Let \mathcal{C} and \mathcal{D} be categories.

013K DEFINITION 8.5.6.1 ► ESSENTIALLY SURJECTIVE FUNCTORS

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **essentially surjective**¹ if it satisfies the following condition:

- (★) For each $D \in \text{Obj}(\mathcal{D})$, there exists some object A of \mathcal{C} such that $F(A) \cong D$.

¹*Further Terminology:* Also called an **eso** functor, where the name “eso” comes from *essentially surjective on objects*.

013L QUESTION 8.5.6.2 ► CHARACTERISATIONS OF FUNCTORS WITH ESSENTIALLY SURJECTIVE PRE/POSTCOMPOSITION

Is there a characterisation of functors $F: \mathcal{C} \rightarrow \mathcal{D}$ such that:

- 013M 1. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is essentially surjective?

- 013N 2. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is essentially surjective?

This question also appears as [MO 468121a].

013P 8.5.7 Equivalences of Categories

013Q DEFINITION 8.5.7.1 ► EQUIVALENCES OF CATEGORIES

Let \mathcal{C} and \mathcal{D} be categories.

- 013R 1. An **equivalence of categories** between \mathcal{C} and \mathcal{D} consists of a pair of functors

$$F: \mathcal{C} \rightarrow \mathcal{D},$$

$$G: \mathcal{D} \rightarrow \mathcal{C}$$

together with natural isomorphisms

$$\eta: \text{id}_C \xrightarrow{\sim} G \circ F,$$

$$\epsilon: F \circ G \xrightarrow{\sim} \text{id}_D.$$

- 013S 2. An **adjoint equivalence of categories** between C and D is an equivalence (F, G, η, ϵ) between C and D which is also an adjunction.

013T **PROPOSITION 8.5.7.2 ► PROPERTIES OF EQUIVALENCES OF CATEGORIES**

Let $F: C \rightarrow D$ be a functor.

- 013U 1. *Characterisations.* If C and D are small¹, then the following conditions are equivalent:²

- 013V (a) The functor F is an equivalence of categories.
 013W (b) The functor F is fully faithful and essentially surjective.
 013X (c) The induced functor

$$\uparrow F\text{Sk}(C): \text{Sk}(C) \rightarrow \text{Sk}(D)$$

is an *isomorphism* of categories.

- 013Y (d) For each $X \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(D, X) \rightarrow \text{Fun}(C, X)$$

is an equivalence of categories.

- 013Z (e) For each $X \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(X, C) \rightarrow \text{Fun}(X, D)$$

is an equivalence of categories.

- 0140 2. *Two-Out-of-Three.* Let

$$\begin{array}{ccc} C & \xrightarrow{G \circ F} & E \\ & \searrow F & \nearrow G \\ & & D \end{array}$$

be a diagram in Cats . If two out of the three functors among F , G , and $G \circ F$ are equivalences of categories, then so is the third.

0141 3. *Stability Under Composition.* Let

$$C \begin{array}{c} \xrightarrow{F} \\ \xleftarrow{G} \end{array} \mathcal{D} \begin{array}{c} \xrightarrow{F'} \\ \xleftarrow{G'} \end{array} \mathcal{E}$$

be a diagram in Cats. If (F, G) and (F', G') are equivalences of categories, then so is their composite $(F' \circ F, G' \circ G)$.

0142 4. *Equivalences vs. Adjoint Equivalences.* Every equivalence of categories can be promoted to an adjoint equivalence.³

0143 5. *Interaction With Groupoids.* If C and \mathcal{D} are groupoids, then the following conditions are equivalent:

0144 (a) The functor F is an equivalence of groupoids.

0145 (b) The following conditions are satisfied:

0146 i. The functor F induces a bijection

$$\pi_0(F) : \pi_0(C) \rightarrow \pi_0(\mathcal{D})$$

of sets.

0147 ii. For each $A \in \text{Obj}(C)$, the induced map

$$F_{x,x} : \text{Aut}_C(A) \rightarrow \text{Aut}_{\mathcal{D}}(F_A)$$

is an isomorphism of groups.

¹Otherwise there will be size issues. One can also work with large categories and universes, or require F to be *constructively* essentially surjective; see [MSE 1465107].

²In ZFC, the equivalence between [Item 1a](#) and [Item 1b](#) is equivalent to the axiom of choice; see [MO 119454].

In Univalent Foundations, this is true without requiring neither the axiom of choice nor the law of excluded middle.

³More precisely, we can promote an equivalence of categories (F, G, η, ϵ) to adjoint equivalences (F, G, η', ϵ) and (F, G, η, ϵ') .

PROOF 8.5.7.3 ► PROOF OF PROPOSITION 8.5.7.2

Item 1: Characterisations

We claim that [Items 1a](#) to [1e](#) are indeed equivalent:

1. [Item 1a](#) \implies [Item 1b](#): Clear.

2. [Item 1b](#) \implies [Item 1a](#): Since F is essentially surjective and C and \mathcal{D} are

small, we can choose, using the axiom of choice, for each $B \in \text{Obj}(\mathcal{D})$, an object j_B of \mathcal{C} and an isomorphism $i_B: B \rightarrow F_{j_B}$ of \mathcal{D} .

Since F is fully faithful, we can extend the assignment $B \mapsto j_B$ to a *unique* functor $j: \mathcal{D} \rightarrow \mathcal{C}$ such that the isomorphisms $i_B: B \rightarrow F_{j_B}$ assemble into a natural isomorphism $\eta: \text{id}_{\mathcal{D}} \xrightarrow{\sim} F \circ j$, with a similar natural isomorphism $\epsilon: \text{id}_{\mathcal{C}} \xrightarrow{\sim} j \circ F$. Hence F is an equivalence.

3. *Item 1a* \implies *Item 1c*: This follows from *Item 4* of [Proposition 8.1.5.3](#).

4. *Item 1c* \implies *Item 1a*: Omitted.

5. *Items 1a, 1d and 1e Are Equivalent*: This follows from ??.

This finishes the proof of [Item 1](#).

Item 2: Two-Out-of-Three

Omitted.

Item 3: Stability Under Composition

Clear.

Item 4: Equivalences vs. Adjoint Equivalences

See [\[Rie17, Proposition 4.4.5\]](#).

Item 5: Interaction With Groupoids

See [\[nLa24, Proposition 4.4\]](#). 

0148 8.5.8 Isomorphisms of Categories

0149 DEFINITION 8.5.8.1 ► ISOMORPHISMS OF CATEGORIES

An **isomorphism of categories** is a pair of functors

$$F: \mathcal{C} \rightarrow \mathcal{D},$$

$$G: \mathcal{D} \rightarrow \mathcal{C}$$

such that we have

$$G \circ F = \text{id}_{\mathcal{C}},$$

$$F \circ G = \text{id}_{\mathcal{D}}.$$

014A **EXAMPLE 8.5.8.2 ▶ EQUIVALENT BUT NON-ISOMORPHIC CATEGORIES**

Categories can be equivalent but non-isomorphic. For example, the category consisting of two isomorphic objects is equivalent to pt , but not isomorphic to it.

014B **PROPOSITION 8.5.8.3 ▶ PROPERTIES OF ISOMORPHISMS OF CATEGORIES**

Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

014C 1. *Characterisations.* If \mathcal{C} and \mathcal{D} are small, then the following conditions are equivalent:

014D (a) The functor F is an isomorphism of categories.

014E (b) The functor F is fully faithful and bijective on objects.

014F (c) For each $X \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, X) \rightarrow \text{Fun}(\mathcal{C}, X)$$

is an isomorphism of categories.

014G (d) For each $X \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(X, \mathcal{C}) \rightarrow \text{Fun}(X, \mathcal{D})$$

is an isomorphism of categories.


PROOF 8.5.8.4 ▶ PROOF OF PROPOSITION 8.5.8.3

Item 1: Characterisations

We claim that **Items 1a to 1d** are indeed equivalent:

1. **Items 1a and 1b Are Equivalent:** Omitted, but similar to **Item 1 of Proposition 8.5.7.2**.

2. **Items 1a, 1c and 1d Are Equivalent:** This follows from ??.

This finishes the proof. 

014H **8.6 More Conditions on Functors**

014J **8.6.1 Dominant Functors**

Let \mathcal{C} and \mathcal{D} be categories.

014K DEFINITION 8.6.1.1 ► DOMINANT FUNCTORS

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **dominant** if every object of \mathcal{D} is a retract of some object in $\text{Im}(F)$, i.e.:

(★) For each $B \in \text{Obj}(\mathcal{D})$, there exist:

- An object A of \mathcal{C} ;
- A morphism $r: F(A) \rightarrow B$ of \mathcal{D} ;
- A morphism $s: B \rightarrow F(A)$ of \mathcal{D} ;

such that we have

$$r \circ s = \text{id}_B,$$

$$\begin{array}{ccc} B & \xrightarrow{s} & F(A) \\ & \searrow \text{id}_B & \downarrow r \\ & & B. \end{array}$$

014L PROPOSITION 8.6.1.2 ► PROPERTIES OF DOMINANT FUNCTORS

Let $F, G: \mathcal{C} \rightarrow \mathcal{D}$ be functors and let $I: \mathcal{X} \rightarrow \mathcal{C}$ be a functor.

014M 1. *Interaction With Right Whiskering.* If I is full and dominant, then the map

$$-\star \text{id}_I: \text{Nat}(F, G) \rightarrow \text{Nat}(F \circ I, G \circ I)$$

is a bijection.

014N 2. *Interaction With Adjunctions.* Let $(F, G): \mathcal{C} \rightleftarrows \mathcal{D}$ be an adjunction.

014P (a) If F is dominant, then G is faithful.

014Q (b) The following conditions are equivalent:

014R i. The functor G is full.

014S ii. The restriction

$$\upharpoonright \text{GIm}_F: \text{Im}(F) \rightarrow \mathcal{C}$$

of G to $\text{Im}(F)$ is full.

PROOF 8.6.1.3 ► PROOF OF PROPOSITION 8.6.1.2

Item 1: Interaction With Right Whiskering

See [DFH75, Proposition 1.4].

Item 2: Interaction With Adjunctions

See [DFH75, Proposition 1.7]. **QUESTION 8.6.1.4 ► CHARACTERISATIONS OF FUNCTORS WITH DOMINANT PRE/POST-COMPOSITION**

014T

Is there a characterisation of functors $F: \mathcal{C} \rightarrow \mathcal{D}$ such that:

014U

1. For each
- $\mathcal{X} \in \text{Obj}(\text{Cats})$
- , the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is dominant?

014V

2. For each
- $\mathcal{X} \in \text{Obj}(\text{Cats})$
- , the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is dominant?

This question also appears as [MO 468121a].

014W 8.6.2 Monomorphisms of CategoriesLet \mathcal{C} and \mathcal{D} be categories.

014X

DEFINITION 8.6.2.1 ► MONOMORPHISMS OF CATEGORIESA functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is a **monomorphism of categories** if it is a monomorphism in Cats (see ??).

014Y

PROPOSITION 8.6.2.2 ► PROPERTIES OF MONOMORPHISMS OF CATEGORIESLet $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

014Z

- 1.
- Characterisations.*
- The following conditions are equivalent:

0150

- (a) The functor
- F
- is a monomorphism of categories.

0151

(b) The functor F is injective on objects and morphisms, i.e. F is injective on objects and the map

$$F: \text{Mor}(C) \rightarrow \text{Mor}(D)$$

is injective.

PROOF 8.6.2.3 ► PROOF OF PROPOSITION 8.6.2.2

Item 1: Characterisations

Omitted. 

0152

QUESTION 8.6.2.4 ► CHARACTERISATIONS OF FUNCTORS WITH MONIC PRE/POSTCOMPOSITION

Is there a characterisation of functors $F: C \rightarrow D$ such that:

0153

1. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(D, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$

is a monomorphism of categories?

0154

2. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, C) \rightarrow \text{Fun}(\mathcal{X}, D)$$

is a monomorphism of categories?

This question also appears as [\[MO 468121a\]](#).

0155 8.6.3 Epimorphisms of Categories

Let C and D be categories.

0156

DEFINITION 8.6.3.1 ► EPIMORPHISMS OF CATEGORIES

A functor $F: C \rightarrow D$ is a **epimorphism of categories** if it is a epimorphism in Cats (see ??).


0157 **PROPOSITION 8.6.3.2 ► PROPERTIES OF EPIMORPHISMS OF CATEGORIES**

Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

0158 1. *Characterisations.* The following conditions are equivalent:¹

0159 (a) The functor F is an epimorphism of categories.

015A (b) For each morphism $f: A \rightarrow B$ of \mathcal{D} , we have a diagram



in \mathcal{D} satisfying the following conditions:

015B i. We have $f = \alpha_0 \circ \phi_1$.

015C ii. We have $f = \psi_m \circ \alpha_{2m}$.

015D iii. For each $0 \leq i \leq 2m$, we have $\alpha_i \in \text{Mor}(\text{Im}(F))$.

015E 2. *Surjectivity on Objects.* If F is an epimorphism of categories, then F is surjective on objects.

¹Further Terminology: This statement is known as **Isbell's zigzag theorem**.

PROOF 8.6.3.3 ► PROOF OF PROPOSITION 8.6.3.2

Item 1: Characterisations

See [Isb68].

Item 2: Surjectivity on Objects

Omitted.



QUESTION 8.6.3.4 ► CHARACTERISATIONS OF FUNCTORS WITH EPIC PRE/POSTCOMPOSITION

015F

Is there a characterisation of functors $F: \mathcal{C} \rightarrow \mathcal{D}$ such that:

015G

1. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is an epimorphism of categories?

015H

2. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is an epimorphism of categories?

This question also appears as [MO 468121a].

015J 8.6.4 Pseudomonadic Functors

Let \mathcal{C} and \mathcal{D} be categories.

015K

DEFINITION 8.6.4.1 ► PSEUDOMONADIC FUNCTORS

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **pseudomonadic** if it satisfies the following conditions:

015L

1. For all diagrams of the form

$$\mathcal{X} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \downarrow \Downarrow \beta \\ \xrightarrow{\psi} \end{array} \mathcal{C} \xrightarrow{F} \mathcal{D},$$

if we have

$$\text{id}_F \star \alpha = \text{id}_F \star \beta,$$

then $\alpha = \beta$.

015M

2. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$ and each natural isomorphism

$$\beta: F \circ \phi \xrightarrow{\sim} F \circ \psi, \quad \mathcal{X} \begin{array}{c} \xrightarrow{F \circ \phi} \\ \beta \downarrow \Downarrow \\ \xrightarrow{F \circ \psi} \end{array} \mathcal{D},$$

there exists a natural isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad \mathcal{X} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{C}$$

such that we have an equality

$$\mathcal{X} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{C} \xrightarrow{F} \mathcal{D} = \mathcal{X} \begin{array}{c} \xrightarrow{F \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{F \circ \psi} \end{array} \mathcal{D}$$

of pasting diagrams, i.e. such that we have

$$\beta = \text{id}_F \star \alpha.$$

015N **PROPOSITION 8.6.4.2 ► PROPERTIES OF PSEUDOMONIC FUNCTORS**

Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

015P 1. *Characterisations.* The following conditions are equivalent:

015Q (a) The functor F is pseudomonadic.

015R (b) The functor F satisfies the following conditions:

015S i. The functor F is faithful, i.e. for each $A, B \in \text{Obj}(\mathcal{C})$, the action on morphisms

$$F_{A,B}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of F at (A, B) is injective.

015T ii. For each $A, B \in \text{Obj}(\mathcal{C})$, the restriction

$$F_{A,B}^{\text{iso}}: \text{Iso}_{\mathcal{C}}(A, B) \rightarrow \text{Iso}_{\mathcal{D}}(F_A, F_B)$$

of the action on morphisms of F at (A, B) to isomorphisms is surjective.

015U (c) We have an isocomma square of the form

$$\mathcal{C} \stackrel{\text{eq.}}{\cong} \mathcal{C} \overset{\leftrightarrow}{\times}_{\mathcal{D}} \mathcal{C}, \quad \begin{array}{ccc} \mathcal{C} & \xrightarrow{\text{id}_{\mathcal{C}}} & \mathcal{C} \\ \text{id}_{\mathcal{C}} \downarrow & \nearrow & \downarrow F \\ \mathcal{C} & \xrightarrow{F} & \mathcal{D} \end{array}$$

015V

in \mathbf{Cats}_2 up to equivalence.

(d) We have an isocomma square of the form

$$C \cong^{\text{eq.}} C \times_{\text{Arr}(\mathcal{D})} \mathcal{D}, \quad \begin{array}{ccc} C & \hookrightarrow & \text{Arr}(C) \\ F \downarrow & \swarrow \dashrightarrow & \downarrow \text{Arr}(F) \\ \mathcal{D} & \hookrightarrow & \text{Arr}(\mathcal{D}) \end{array}$$

in \mathbf{Cats}_2 up to equivalence.

015W

(e) For each $\mathcal{X} \in \text{Obj}(\mathbf{Cats})$, the postcomposition¹ functor

$$F_* : \text{Fun}(\mathcal{X}, C) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is pseudomonic.

015X

2. *Conservativity.* If F is pseudomonic, then F is conservative.

015Y

3. *Essential Injectivity.* If F is pseudomonic, then F is essentially injective.

¹Asking the precomposition functors

$$F^* : \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$

to be pseudomonic leads to pseudoepic functors; see [Item 1b](#) of [Item 1](#) of [Proposition 8.6.5.2](#).

PROOF 8.6.4.3 ► PROOF OF PROPOSITION 8.6.4.2

Item 1: Characterisations

Omitted.

Item 2: Conservativity

Omitted.

Item 3: Essential Injectivity

Omitted. 

015Z 8.6.5 Pseudoepic Functors

Let C and \mathcal{D} be categories.

0160 DEFINITION 8.6.5.1 ► PSEUDOEPIC FUNCTORS

A functor $F: C \rightarrow \mathcal{D}$ is **pseudoepic** if it satisfies the following conditions:

- 0161 1. For all diagrams of the form

$$C \xrightarrow{F} \mathcal{D} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \beta \\ \xrightarrow{\psi} \end{array} \mathcal{X},$$

if we have

$$\alpha \star \text{id}_F = \beta \star \text{id}_F,$$

then $\alpha = \beta$.

- 0162 2. For each $X \in \text{Obj}(C)$ and each 2-isomorphism

$$\beta: \phi \circ F \xrightarrow{\sim} \psi \circ F, \quad C \begin{array}{c} \xrightarrow{\phi \circ F} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ F} \end{array} \mathcal{X}$$

of C , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad \mathcal{D} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{X}$$

of C such that we have an equality

$$C \xrightarrow{F} \mathcal{D} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{X} = C \begin{array}{c} \xrightarrow{\phi \circ F} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ F} \end{array} \mathcal{X}$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \alpha \star \text{id}_F.$$

0163 PROPOSITION 8.6.5.2 ► PROPERTIES OF PSEUDOEPIC FUNCTORS

Let $F: C \rightarrow \mathcal{D}$ be a functor.

- 0164 1. *Characterisations.* The following conditions are equivalent:

0165

(a) The functor F is pseudoepic.

0166

(b) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the functor

$$F^* : \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

given by precomposition by F is pseudomonic.

0167

(c) We have an isococcomma square of the form

$$\mathcal{D} \stackrel{\text{eq.}}{\cong} \mathcal{D} \amalg_C \mathcal{D}, \quad \begin{array}{ccc} \mathcal{D} & \xleftarrow{\text{id}_{\mathcal{D}}} & \mathcal{D} \\ \uparrow \text{id}_{\mathcal{D}} & \nearrow & \uparrow F \\ \mathcal{D} & \xleftarrow{F} & \mathcal{C} \end{array}$$

in Cats_2 up to equivalence.

0168

2. *Dominance*. If F is pseudoepic, then F is dominant ([Definition 8.6.1.1](#)).**PROOF 8.6.5.3 ► PROOF OF PROPOSITION 8.6.5.2**


Item 1: Characterisations

Omitted.

Item 2: Dominance

If F is pseudoepic, then

$$F^* : \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is pseudomonic for all $\mathcal{X} \in \text{Obj}(\text{Cats})$, and thus in particular faithful. By [Item 4g](#) of [Item 4](#) of [Proposition 8.5.1.2](#), this is equivalent to requiring F to be dominant. 

0169

QUESTION 8.6.5.4 ► CHARACTERISATIONS OF PSEUDOEPIC FUNCTORS

Is there a nice characterisation of the pseudoepic functors, similarly to the characterisation of pseudomonic functors given in [Item 1b](#) of [Item 1](#) of [Proposition 8.6.4.2](#)?

This question also appears as [\[MO 321971\]](#).

016A **QUESTION 8.6.5.5 ▶ MUST A PSEUDOMONIC AND PSEUDOEPIC FUNCTOR BE AN EQUIVALENCE OF CATEGORIES**

A pseudomonic and pseudoepic functor is dominant, faithful, essentially injective, and full on isomorphisms. Is it necessarily an equivalence of categories? If not, how bad can this fail, i.e. how far can a pseudomonic and pseudoepic functor be from an equivalence of categories? This question also appears as [MO 468334].

016B **QUESTION 8.6.5.6 ▶ CHARACTERISATIONS OF FUNCTORS WITH PSEUDOEPIC PRE-/POSTCOMPOSITION**

Is there a characterisation of functors $F: \mathcal{C} \rightarrow \mathcal{D}$ such that:

016C 1. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(\mathcal{C}, \mathcal{X})$$

is pseudoepic?

016D 2. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, \mathcal{C}) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is pseudoepic?

This question also appears as [MO 468121a].

016E **8.7 Even More Conditions on Functors**

016F **8.7.1 Injective on Objects Functors**

Let \mathcal{C} and \mathcal{D} be categories.

016G **DEFINITION 8.7.1.1 ▶ INJECTIVE ON OBJECTS FUNCTORS**

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **injective on objects** if the action on objects

$$F: \text{Obj}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{D})$$

of F is injective.

016H **PROPOSITION 8.7.1.2 ► PROPERTIES OF INJECTIVE ON OBJECTS FUNCTORS**

Let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a functor.

- 016J 1. *Characterisations.* The following conditions are equivalent:
- 016K (a) The functor F is injective on objects.
- 016L (b) The functor F is an isocofibration in \mathbf{Cats}_2 .

PROOF 8.7.1.3 ► PROOF OF PROPOSITION 8.7.1.2

Item 1: Characterisations

Omitted. 

016M **8.7.2 Surjective on Objects Functors**

Let \mathcal{C} and \mathcal{D} be categories.

016N **DEFINITION 8.7.2.1 ► SURJECTIVE ON OBJECTS FUNCTORS**

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **surjective on objects** if the action on objects

$$F: \text{Obj}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{D})$$

of F is surjective.

016P **8.7.3 Bijective on Objects Functors**

Let \mathcal{C} and \mathcal{D} be categories.

016Q **DEFINITION 8.7.3.1 ► BIJECTIVE ON OBJECTS FUNCTORS**

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **bijective on objects**¹ if the action on objects

$$F: \text{Obj}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{D})$$

of F is a bijection.

¹*Further Terminology:* Also called a **bo** functor.

016R **8.7.4 Functors Representably Faithful on Cores**

Let \mathcal{C} and \mathcal{D} be categories.

016S DEFINITION 8.7.4.1 ► FUNCTORS REPRESENTABLY FAITHFUL ON CORES

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **representably faithful on cores** if, for each $X \in \text{Obj}(\text{Cats})$, the postcomposition by F functor

$$F_*: \text{Core}(\text{Fun}(X, \mathcal{C})) \rightarrow \text{Core}(\text{Fun}(X, \mathcal{D}))$$

is faithful.

016T REMARK 8.7.4.2 ► UNWINDING DEFINITION 8.7.4.1

In detail, a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **representably faithful on cores** if, given a diagram of the form

$$\mathcal{X} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \downarrow \Downarrow \beta \\ \xrightarrow{\psi} \end{array} \mathcal{C} \xrightarrow{F} \mathcal{D},$$

if α and β are natural isomorphisms and we have

$$\text{id}_F \star \alpha = \text{id}_F \star \beta,$$

then $\alpha = \beta$.

016U QUESTION 8.7.4.3 ► CHARACTERISATION OF FUNCTORS REPRESENTABLY FAITHFUL ON CORES

Is there a characterisation of functors representably faithful on cores?

016V 8.7.5 Functors Representably Full on Cores

Let \mathcal{C} and \mathcal{D} be categories.

016W DEFINITION 8.7.5.1 ► FUNCTORS REPRESENTABLY FULL ON CORES

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **representably full on cores** if, for each $X \in \text{Obj}(\text{Cats})$, the postcomposition by F functor

$$F_*: \text{Core}(\text{Fun}(X, \mathcal{C})) \rightarrow \text{Core}(\text{Fun}(X, \mathcal{D}))$$

is full.

016X **REMARK 8.7.5.2** ► UNWINDING DEFINITION 8.7.5.1

In detail, a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **representably full on cores** if, for each $X \in \text{Obj}(\text{Cats})$ and each natural isomorphism

$$\beta: F \circ \phi \xrightarrow{\sim} F \circ \psi, \quad X \begin{array}{c} \xrightarrow{F \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{F \circ \psi} \end{array} \mathcal{D},$$

there exists a natural isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{C}$$

such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{C} \xrightarrow{F} \mathcal{D} = X \begin{array}{c} \xrightarrow{F \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{F \circ \psi} \end{array} \mathcal{D}$$

of pasting diagrams in Cats_2 , i.e. such that we have

$$\beta = \text{id}_F \star \alpha.$$

016Y

QUESTION 8.7.5.3 ► CHARACTERISATION OF FUNCTORS REPRESENTABLY FULL ON CORES

Is there a characterisation of functors representably full on cores?
This question also appears as [MO 468121a].

016Z **8.7.6 Functors Representably Fully Faithful on Cores**

Let \mathcal{C} and \mathcal{D} be categories.

0170 **DEFINITION 8.7.6.1** ► FUNCTORS REPRESENTABLY FULLY FAITHFUL ON CORES

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **representably fully faithful on cores** if, for each $X \in \text{Obj}(\text{Cats})$, the postcomposition by F functor

$$F_*: \text{Core}(\text{Fun}(X, \mathcal{C})) \rightarrow \text{Core}(\text{Fun}(X, \mathcal{D}))$$

is fully faithful.

0171 **REMARK 8.7.6.2** ▶ UNWINDING DEFINITION 8.7.6.1

In detail, a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **representably fully faithful on cores** if it satisfies the conditions in **Remarks 8.7.4.2** and **8.7.5.2**, i.e.:

- 0172 1. For all diagrams of the form

$$\mathcal{X} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \beta \\ \xrightarrow{\psi} \end{array} \mathcal{C} \xrightarrow{F} \mathcal{D},$$

with α and β natural isomorphisms, if we have $\text{id}_F \star \alpha = \text{id}_F \star \beta$, then $\alpha = \beta$.

- 0173 2. For each
- $\mathcal{X} \in \text{Obj}(\text{Cats})$
- and each natural isomorphism

$$\beta: F \circ \phi \xrightarrow{\sim} F \circ \psi, \quad \mathcal{X} \begin{array}{c} \xrightarrow{F \circ \phi} \\ \beta \Downarrow \Downarrow \\ \xrightarrow{F \circ \psi} \end{array} \mathcal{D}$$

of \mathcal{C} , there exists a natural isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad \mathcal{X} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{C}$$

of \mathcal{C} such that we have an equality

$$\mathcal{X} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{C} \xrightarrow{F} \mathcal{D} = \mathcal{X} \begin{array}{c} \xrightarrow{F \circ \phi} \\ \beta \Downarrow \Downarrow \\ \xrightarrow{F \circ \psi} \end{array} \mathcal{D}$$

of pasting diagrams in Cats_2 , i.e. such that we have

$$\beta = \text{id}_F \star \alpha.$$

0174 **QUESTION 8.7.6.3** ▶ CHARACTERISATION OF FUNCTORS REPRESENTABLY FULLY FAITHFUL ON CORES

Is there a characterisation of functors representably fully faithful on cores?

0175 **8.7.7 Functors Corepresentably Faithful on Cores**

Let \mathcal{C} and \mathcal{D} be categories.

0176

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **corepresentably faithful on cores** if, for each $X \in \text{Obj}(\text{Cats})$, the postcomposition by F functor

$$F_*: \text{Core}(\text{Fun}(X, \mathcal{C})) \rightarrow \text{Core}(\text{Fun}(X, \mathcal{D}))$$

is faithful.

0177 **REMARK 8.7.7.2 ► UNWINDING DEFINITION 8.7.7.1**

In detail, a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **corepresentably faithful on cores** if, given a diagram of the form

$$\mathcal{C} \xrightarrow{F} \mathcal{D} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \Downarrow \beta \\ \xrightarrow{\psi} \end{array} \mathcal{X},$$

if α and β are natural isomorphisms and we have

$$\alpha \star \text{id}_F = \beta \star \text{id}_F,$$

then $\alpha = \beta$.

0178 **QUESTION 8.7.7.3 ► CHARACTERISATION OF FUNCTORS COREPRESENTABLY FAITHFUL ON CORES**

Is there a characterisation of functors corepresentably faithful on cores?

0179 **8.7.8 Functors Corepresentably Full on Cores**

Let \mathcal{C} and \mathcal{D} be categories.

017A **DEFINITION 8.7.8.1 ► FUNCTORS COREPRESENTABLY FULL ON CORES**

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **corepresentably full on cores** if, for each $X \in \text{Obj}(\text{Cats})$, the postcomposition by F functor

$$F_*: \text{Core}(\text{Fun}(X, \mathcal{C})) \rightarrow \text{Core}(\text{Fun}(X, \mathcal{D}))$$

is full.

017B

REMARK 8.7.8.2 ► UNWINDING DEFINITION 8.7.8.1

In detail, a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **corepresentably full on cores** if, for each $\mathcal{X} \in \text{Obj}(\text{Cats})$ and each natural isomorphism

$$\beta: \phi \circ F \xrightarrow{\sim} \psi \circ F, \quad \mathcal{C} \begin{array}{c} \xrightarrow{\phi \circ F} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ F} \end{array} \mathcal{X},$$

there exists a natural isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad \mathcal{D} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{X}$$

such that we have an equality

$$\mathcal{X} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{C} \xrightarrow{F} \mathcal{D} = \mathcal{X} \begin{array}{c} \xrightarrow{F \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{F \circ \psi} \end{array} \mathcal{D}$$

of pasting diagrams in Cats_2 , i.e. such that we have

$$\beta = \alpha \star \text{id}_F.$$

017C

QUESTION 8.7.8.3 ► CHARACTERISATION OF FUNCTORS COREPRESENTABLY FULL ON CORES

Is there a characterisation of functors corepresentably full on cores?
This question also appears as [MO 468121a].

017D 8.7.9 Functors Corepresentably Fully Faithful on Cores

Let \mathcal{C} and \mathcal{D} be categories.

017E

DEFINITION 8.7.9.1 ► FUNCTORS COREPRESENTABLY FULLY FAITHFUL ON CORES

A functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **corepresentably fully faithful on cores** if, for each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition by F functor

$$F_*: \text{Core}(\text{Fun}(\mathcal{X}, \mathcal{C})) \rightarrow \text{Core}(\text{Fun}(\mathcal{X}, \mathcal{D}))$$

is fully faithful.

017F **REMARK 8.7.9.2 ▶ UNWINDING DEFINITION 8.7.9.1**

In detail, a functor $F: \mathcal{C} \rightarrow \mathcal{D}$ is **corepresentably fully faithful on cores** if it satisfies the conditions in **Remarks 8.7.7.2** and **8.7.8.2**, i.e.:

017G 1. For all diagrams of the form

$$\mathcal{C} \xrightarrow{F} \mathcal{D} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \beta \\ \xrightarrow{\psi} \end{array} \mathcal{X},$$

if α and β are natural isomorphisms and we have

$$\alpha \star \text{id}_F = \beta \star \text{id}_F,$$

then $\alpha = \beta$.

017H 2. For each $\mathcal{X} \in \text{Obj}(\text{Cats})$ and each natural isomorphism

$$\beta: \phi \circ F \xrightarrow{\sim} \psi \circ F, \quad \mathcal{C} \begin{array}{c} \xrightarrow{\phi \circ F} \\ \beta \Downarrow \Downarrow \\ \xrightarrow{\psi \circ F} \end{array} \mathcal{X},$$

there exists a natural isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad \mathcal{D} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{X}$$

such that we have an equality

$$\mathcal{X} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{C} \xrightarrow{F} \mathcal{D} = \mathcal{X} \begin{array}{c} \xrightarrow{F \circ \phi} \\ \beta \Downarrow \Downarrow \\ \xrightarrow{F \circ \psi} \end{array} \mathcal{D}$$

of pasting diagrams in Cats_2 , i.e. such that we have

$$\beta = \alpha \star \text{id}_F.$$

QUESTION 8.7.9.3 ► CHARACTERISATION OF FUNCTORS COREPRESENTABLY FULLY FAITHFUL ON CORES

017J

Is there a characterisation of functors corepresentably fully faithful on cores?

017K 8.8 Natural Transformations

017L 8.8.1 Transformations

Let \mathcal{C} and \mathcal{D} be categories and $F, G: \mathcal{C} \Rightarrow \mathcal{D}$ be functors.

017M

DEFINITION 8.8.1.1 ► TRANSFORMATIONS

A **transformation**¹ $\alpha: F \Rightarrow G$ **from F to G** is a collection

$$\{\alpha_A: F(A) \rightarrow G(A)\}_{A \in \text{Obj}(\mathcal{C})}$$

of morphisms of \mathcal{D} .

¹*Further Terminology:* Also called an **unnatural transformation** for emphasis.

017N

NOTATION 8.8.1.2 ► THE SET OF TRANSFORMATIONS BETWEEN TWO FUNCTORS

We write $\text{Trans}(F, G)$ for the set of transformations from F to G .

017P 8.8.2 Natural Transformations

Let \mathcal{C} and \mathcal{D} be categories and $F, G: \mathcal{C} \Rightarrow \mathcal{D}$ be functors.

017Q

DEFINITION 8.8.2.1 ► NATURAL TRANSFORMATIONS

A **natural transformation** $\alpha: F \Longrightarrow G$ **from F to G** is a transformation

$$\{\alpha_A: F(A) \rightarrow G(A)\}_{A \in \text{Obj}(\mathcal{C})}$$

from F to G such that, for each morphism $f: A \rightarrow B$ of \mathcal{C} , the diagram

$$\begin{array}{ccc} F(A) & \xrightarrow{F(f)} & F(B) \\ \alpha_A \downarrow & & \downarrow \alpha_B \\ G(A) & \xrightarrow{G(f)} & G(B) \end{array}$$

commutes.¹

¹*Further Terminology:* The morphism $\alpha_A: F_A \rightarrow G_A$ is called the **component of α at A** .

017R **REMARK 8.8.2.2** ▶ PICTURING NATURAL TRANSFORMATIONS IN DIAGRAMS

We denote natural transformations in diagrams as

$$C \begin{array}{c} \xrightarrow{F} \\ \alpha \Downarrow \\ \xrightarrow{G} \end{array} \mathcal{D}.$$

017S

NOTATION 8.8.2.3 ▶ THE SET OF NATURAL TRANSFORMATIONS BETWEEN TWO FUNCTORS

We write $\text{Nat}(F, G)$ for the set of natural transformations from F to G .

017T

EXAMPLE 8.8.2.4 ▶ IDENTITY NATURAL TRANSFORMATIONS


The **identity natural transformation** $\text{id}_F: F \Rightarrow F$ of F is the natural transformation consisting of the collection

$$\{\text{id}_{F(A)}: F(A) \rightarrow F(A)\}_{A \in \text{Obj}(C)}.$$

PROOF 8.8.2.5 ▶ PROOF OF EXAMPLE 8.8.2.4

The naturality condition for id_F is the requirement that, for each morphism $f: A \rightarrow B$ of C , the diagram

$$\begin{array}{ccc} F(A) & \xrightarrow{F(f)} & F(B) \\ \text{id}_{F(A)} \downarrow & & \downarrow \text{id}_{F(B)} \\ F(A) & \xrightarrow{F(f)} & F(B) \end{array}$$

commutes, which follows from unitality of the composition of C . 

017U

DEFINITION 8.8.2.6 ▶ EQUALITY OF NATURAL TRANSFORMATIONS

Two natural transformations $\alpha, \beta: F \Rightarrow G$ are **equal** if we have

$$\alpha_A = \beta_A$$

for each $A \in \text{Obj}(C)$.

017V **8.8.3 Vertical Composition of Natural Transformations**

017W

DEFINITION 8.8.3.1 ► VERTICAL COMPOSITION OF NATURAL TRANSFORMATIONS

The **vertical composition** of two natural transformations $\alpha: F \Rightarrow G$ and $\beta: G \Rightarrow H$ as in the diagram

$$\begin{array}{ccc}
 & F & \\
 & \curvearrowright & \\
 C & \xrightarrow{G} & \mathcal{D} \\
 & \curvearrowleft & \\
 & H & \\
 & \alpha \Downarrow & \\
 & \beta \Downarrow &
 \end{array}$$

is the natural transformation $\beta \circ \alpha: F \Rightarrow H$ consisting of the collection

$$\{(\beta \circ \alpha)_A: F(A) \rightarrow H(A)\}_{A \in \text{Obj}(\mathcal{C})}$$

with

$$(\beta \circ \alpha)_A \stackrel{\text{def}}{=} \beta_A \circ \alpha_A$$

for each $A \in \text{Obj}(\mathcal{C})$.

PROOF 8.8.3.2 ► PROOF OF DEFINITION 8.8.3.1

The naturality condition for $\beta \circ \alpha$ is the requirement that the boundary of the diagram

$$\begin{array}{ccc}
 F(A) & \xrightarrow{F(f)} & F(B) \\
 \alpha_A \downarrow & (1) & \downarrow \alpha_B \\
 G(A) & \xrightarrow{G(f)} & G(B) \\
 \beta_A \downarrow & (2) & \downarrow \beta_B \\
 H(A) & \xrightarrow{H(f)} & H(B)
 \end{array}$$

commutes. Since

1. Subdiagram (1) commutes by the naturality of α .
2. Subdiagram (2) commutes by the naturality of β .

so does the boundary diagram. Hence $\beta \circ \alpha$ is a natural transformation. \square

PROPOSITION 8.8.3.3 ► PROPERTIES OF VERTICAL COMPOSITION OF NATURAL TRANSFORMATIONS

017X

Let \mathcal{C} , \mathcal{D} , and \mathcal{E} be categories.


017Y

1. *Functionality.* The assignment $(\beta, \alpha) \mapsto \beta \circ \alpha$ defines a function

$$\circ_{F,G,H}: \text{Nat}(G, H) \times \text{Nat}(F, G) \rightarrow \text{Nat}(F, H).$$

017Z

2. *Associativity.* Let $F, G, H, K: \mathcal{C} \rightrightarrows \mathcal{D}$ be functors. The diagram



commutes, i.e. given natural transformations

$$F \xrightarrow{\alpha} G \xrightarrow{\beta} H \xrightarrow{\gamma} K,$$


we have

$$(\gamma \circ \beta) \circ \alpha = \gamma \circ (\beta \circ \alpha).$$

0180

3. *Unitality.* Let $F, G: \mathcal{C} \rightrightarrows \mathcal{D}$ be functors.


(a) *Left Unitality.* The diagram



commutes, i.e. given a natural transformation $\alpha: F \rightrightarrows G$, we have

$$\text{id}_G \circ \alpha = \alpha.$$

(b) *Right Unitality*. The diagram




commutes, i.e. given a natural transformation $\alpha: F \Rightarrow G$, we have


$$\alpha \circ \text{id}_F = \alpha.$$

0181

4. *Middle Four Exchange*. Let $F_1, F_2, F_3: C \rightarrow D$ and $G_1, G_2, G_3: D \rightarrow E$ be functors. The diagram



commutes, i.e. given a diagram



in Cats_2 , we have

$$(\beta' \star \alpha') \circ (\beta \star \alpha) = (\beta' \circ \beta) \star (\alpha' \circ \alpha).$$

PROOF 8.8.3.4 ► PROOF OF PROPOSITION 8.8.3.3

Item 1: Functionality

Clear.

Item 2: Associativity

Indeed, we have

$$\begin{aligned}
 ((\gamma \circ \beta) \circ \alpha)_A &\stackrel{\text{def}}{=} (\gamma \circ \beta)_A \circ \alpha_A \\
 &\stackrel{\text{def}}{=} (\gamma_A \circ \beta_A) \circ \alpha_A \\
 &= \gamma_A \circ (\beta_A \circ \alpha_A) \\
 &\stackrel{\text{def}}{=} \gamma_A \circ (\beta \circ \alpha)_A \\
 &\stackrel{\text{def}}{=} (\gamma \circ (\beta \circ \alpha))_A
 \end{aligned}$$

for each $A \in \text{Obj}(C)$, showing the desired equality.

Item 3: Unitality

We have

$$\begin{aligned}
 (\text{id}_G \circ \alpha)_A &= \text{id}_G \circ \alpha_A \\
 &= \alpha_A, \\
 (\alpha \circ \text{id}_F)_A &= \alpha_A \circ \text{id}_F \\
 &= \alpha_A
 \end{aligned}$$

for each $A \in \text{Obj}(C)$, showing the desired equality.

Item 4: Middle Four Exchange

This is proved in [Item 4 of Proposition 8.8.4.4](#). 

0182 8.8.4 Horizontal Composition of Natural Transformations

0183 DEFINITION 8.8.4.1 ► HORIZONTAL COMPOSITION OF NATURAL TRANSFORMATIONS

The **horizontal composition**^{1,2} of two natural transformations $\alpha: F \Rightarrow G$ and $\beta: H \Rightarrow K$ as in the diagram

$$\begin{array}{ccccc}
 & & F & & H \\
 & \curvearrowright & & \curvearrowright & \\
 C & & & \mathcal{D} & & \mathcal{E} \\
 & \alpha \Downarrow & & \beta \Downarrow & \\
 & \curvearrowleft & & \curvearrowleft & \\
 & & G & & K
 \end{array}$$

of α and β is the natural transformation

$$\beta \star \alpha: (H \circ F) \Rightarrow (K \circ G),$$

as in the diagram

$$\begin{array}{ccc} & H \circ F & \\ \curvearrowright & \parallel & \curvearrowleft \\ C & \beta \star \alpha & \mathcal{E}, \\ \curvearrowleft & \Downarrow & \curvearrowright \\ & K \circ G & \end{array}$$

consisting of the collection

$$\{(\beta \star \alpha)_A : H(F(A)) \rightarrow K(G(A))\}_{A \in \text{Obj}(C)},$$

of morphisms of \mathcal{E} with

$$\begin{array}{ccc} H(F(A)) & \xrightarrow{H(\alpha_A)} & H(G(A)) \\ \beta_{F(A)} \downarrow & & \downarrow \beta_{G(A)} \\ K(F(A)) & \xrightarrow{K(\alpha_A)} & K(G(A)). \end{array}$$

$$\begin{aligned} (\beta \star \alpha)_A &\stackrel{\text{def}}{=} \beta_{G(A)} \circ H(\alpha_A) \\ &= K(\alpha_A) \circ \beta_{F(A)}, \end{aligned}$$

¹Further Terminology: Also called the **Godement product** of α and β .

²Horizontal composition forms a map

$$\star_{(F,H),(G,K)} : \text{Nat}(H, K) \times \text{Nat}(F, G) \rightarrow \text{Nat}(H \circ F, K \circ G).$$

PROOF 8.8.4.2 ► PROOF OF DEFINITION 8.8.4.1

First, we claim that we indeed have

$$\beta_{G(A)} \circ H(\alpha_A) = K(\alpha_A) \circ \beta_{F(A)},$$

$$\begin{array}{ccc} H(F(A)) & \xrightarrow{H(\alpha_A)} & H(G(A)) \\ \beta_{F(A)} \downarrow & & \downarrow \beta_{G(A)} \\ K(F(A)) & \xrightarrow{K(\alpha_A)} & K(G(A)). \end{array}$$


This is, however, simply the naturality square for β applied to the morphism $\alpha_A : F(A) \rightarrow G(A)$. Next, we check the naturality condition for $\beta \star \alpha$, which

is the requirement that the boundary of the diagram

$$\begin{array}{ccc}
 H(F(A)) & \xrightarrow{H(F(f))} & H(F(B)) \\
 \downarrow H(\alpha_A) & (1) & \downarrow H(\alpha_B) \\
 H(G(A)) & \xrightarrow{H(G(f))} & H(G(B)) \\
 \downarrow \beta_{G(A)} & (2) & \downarrow \beta_{G(B)} \\
 K(G(A)) & \xrightarrow{K(G(f))} & K(G(B))
 \end{array}$$

commutes. Since

1. Subdiagram (1) commutes by the naturality of α .
2. Subdiagram (2) commutes by the naturality of β .

so does the boundary diagram. Hence $\beta \circ \alpha$ is a natural transformation.¹ 

¹Reference: [Bor94, Proposition 1.3.4].

DEFINITION 8.8.4.3 ► WHISKERING OF FUNCTORS WITH NATURAL TRANSFORMATIONS

0184

Let

$$\mathcal{X} \xrightarrow{F} \mathcal{C} \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} \mathcal{D} \xrightarrow{G} \mathcal{Y}$$

be a diagram in \mathbf{Cats}_2 .

0185

1. The **left whiskering of α with G** is the natural transformation¹

$$\mathrm{id}_G \star \alpha: G \circ \phi \Longrightarrow G \circ \psi.$$

0186

2. The **right whiskering of α with F** is the natural transformation²

$$\alpha \star \mathrm{id}_F: \phi \circ F \Longrightarrow \psi \circ F.$$

¹Further Notation: Also written $G\alpha$ or $G \star \alpha$, although we won't use either of these notations in this work.

²Further Notation: Also written αF or $\alpha \star F$, although we won't use either of these notations in this work.

PROPOSITION 8.8.4.4 ► PROPERTIES OF HORIZONTAL COMPOSITION OF NATURAL TRANSFORMATIONS

0187

Let \mathcal{C} , \mathcal{D} , and \mathcal{E} be categories.

0188

1. *Functionality.* The assignment $(\beta, \alpha) \mapsto \beta \star \alpha$ defines a function

$$\star_{(F,G),(H,K)} : \text{Nat}(H, K) \times \text{Nat}(F, G) \rightarrow \text{Nat}(H \circ F, K \circ G).$$

0189

2. *Associativity.* Let

$$\mathcal{C} \begin{array}{c} \xrightarrow{F_1} \\ \xrightarrow{G_1} \end{array} \mathcal{D} \begin{array}{c} \xrightarrow{F_2} \\ \xrightarrow{G_2} \end{array} \mathcal{E} \begin{array}{c} \xrightarrow{F_3} \\ \xrightarrow{G_3} \end{array} \mathcal{F}$$

be a diagram in Cats_2 . The diagram

$$\begin{array}{ccc} \text{Nat}(F_3, G_3) \times \text{Nat}(F_2, G_2) \times \text{Nat}(F_1, G_1) & \xrightarrow{\star_{(F_2, G_2), (F_3, G_3)} \times \text{id}} & \text{Nat}(F_3 \circ F_2, G_3 \circ G_2) \times \text{Nat}(F_1, G_1) \\ \downarrow \text{id} \times \star_{(F_1, G_1), (F_2, G_2)} & & \downarrow \star_{(F_3 \circ F_2), (G_3 \circ G_2, F_1, G_1)} \\ \text{Nat}(F_3, G_3) \times \text{Nat}(F_2 \circ F_1, G_2 \circ G_1) & \xrightarrow{\star_{(F_2 \circ F_1), (G_2 \circ G_1, F_3, G_3)}} & \text{Nat}(F_3 \circ F_2 \circ F_1, G_3 \circ G_2 \circ G_1) \end{array}$$

commutes, i.e. given natural transformations

$$\mathcal{C} \begin{array}{c} \xrightarrow{F_1} \\ \alpha \Downarrow \\ \xrightarrow{G_1} \end{array} \mathcal{D} \begin{array}{c} \xrightarrow{F_2} \\ \beta \Downarrow \\ \xrightarrow{G_2} \end{array} \mathcal{E} \begin{array}{c} \xrightarrow{F_3} \\ \gamma \Downarrow \\ \xrightarrow{G_3} \end{array} \mathcal{F},$$

we have

$$(\gamma \star \beta) \star \alpha = \gamma \star (\beta \star \alpha).$$

018A

3. *Interaction With Identities.* Let $F: \mathcal{C} \rightarrow \mathcal{D}$ and $G: \mathcal{D} \rightarrow \mathcal{E}$ be functors. The diagram


$$\begin{array}{ccc} \text{pt} \times \text{pt} & \xrightarrow{[\text{id}_G] \times [\text{id}_F]} & \text{Nat}(G, G) \times \text{Nat}(F, F) \\ \uparrow \text{?} & & \downarrow \star_{(F,F),(G,G)} \\ \text{pt} & \xrightarrow{[\text{id}_{G \circ F}]} & \text{Nat}(G \circ F, G \circ F) \end{array}$$

commutes, i.e. we have


$$\text{id}_G \star \text{id}_F = \text{id}_{G \circ F}.$$

018B

4. *Middle Four Exchange.* Let $F_1, F_2, F_3 : C \rightarrow D$ and $G_1, G_2, G_3 : D \rightarrow E$ be functors. The diagram



commutes, i.e. given a diagram



in Cats_2 , we have

$$(\beta' \star \alpha') \circ (\beta \star \alpha) = (\beta' \circ \beta) \star (\alpha' \circ \alpha).$$

PROOF 8.8.4.5 ► PROOF OF PROPOSITION 8.8.4.4

Item 1: Functionality

Clear.

Item 2: Associativity

Omitted.

Item 3: Interaction With Identities


We have

$$\begin{aligned}
 (\text{id}_G \star \text{id}_F)_A &\stackrel{\text{def}}{=} (\text{id}_G)_{F_A} \circ G_{(\text{id}_F)_A} \\
 &\stackrel{\text{def}}{=} \text{id}_{G_{F_A}} \circ G_{\text{id}_{F_A}} \\
 &= \text{id}_{G_{F_A}} \circ \text{id}_{G_{F_A}} \\
 &= \text{id}_{G_{F_A}} \\
 &\stackrel{\text{def}}{=} (\text{id}_{G \circ F})_A
 \end{aligned}$$


for each $A \in \text{Obj}(\mathcal{C})$, showing the desired equality.

Item 4: Middle Four Exchange


Let $A \in \text{Obj}(\mathcal{C})$ and consider the diagram



The top composition



is given by $((\beta' \circ \beta) \star (\alpha' \circ \alpha))_A$, while the bottom composition



is given by $((\beta' \star \alpha') \circ (\beta \star \alpha))_A$. Now, Subdiagram (1) corresponds to the naturality condition

$$G_2(\alpha'_A) \circ \beta_{F_2(A)} = \beta_{F_3(A)} \circ G_1(\alpha'_A), \quad \begin{array}{ccc} G_1(F_2(A)) & \xrightarrow{G_1(\alpha'_A)} & G_1(F_3(A)) \\ \beta_{F_2(A)} \downarrow & & \downarrow \beta_{F_3(A)} \\ G_2(F_2(A)) & \xrightarrow{G_2(\alpha'_A)} & G_2(F_3(A)) \end{array}$$

for $\beta: G_1 \Rightarrow G_2$ at $\alpha'_A: F_2(A) \rightarrow F_3(A)$, and thus commutes. Thus we have

$$((\beta' \circ \beta) \star (\alpha' \circ \alpha))_A = ((\beta' \star \alpha') \circ (\beta \star \alpha))_A$$

for each $A \in \text{Obj}(C)$ and therefore

$$(\beta' \star \alpha') \circ (\beta \star \alpha) = (\beta' \circ \beta) \star (\alpha' \circ \alpha).$$


This finishes the proof. ▢

018C **8.8.5 Properties of Natural Transformations**

018D **PROPOSITION 8.8.5.1 ► NATURAL TRANSFORMATIONS AS CATEGORICAL HOMOTOPIES**


Let $F, G: C \Rightarrow D$ be functors. The following data are equivalent:¹

- 018E 1. A natural transformation $\alpha: F \Rightarrow G$.
- 018F 2. A functor $[\alpha]: C \rightarrow D^{\mathbb{1}}$ filling the diagram



018G

3. A functor $[\alpha]: C \times \mathbb{1} \rightarrow \mathcal{D}$ filling the diagram



¹Taken from [MO 64365].

PROOF 8.8.5.2 ► PROOF OF PROPOSITION 8.8.5.1

From Item 1 to Item 2 and Back

We may identify $\mathcal{D}^{\mathbb{1}}$ with $\text{Arr}(\mathcal{D})$. Given a natural transformation $\alpha: F \Rightarrow G$, we have a functor

$$\begin{array}{ccc}
 [\alpha]: C & \longrightarrow & \mathcal{D}^{\mathbb{1}} \\
 A & \longmapsto & \alpha_A \\
 \\
 (f: A \rightarrow B) & \longmapsto & \left(\begin{array}{ccc} F_A & \xrightarrow{F_f} & F_B \\ \alpha_A \downarrow & & \downarrow \alpha_B \\ G_A & \xrightarrow{G_f} & G_B \end{array} \right)
 \end{array}$$

making the diagram in **Item 2** commute. Conversely, every such functor gives rise to a natural transformation from F to G , and these constructions are inverse to each other.

From Item 2 to Item 3 and Back

This follows from **Item 3** of **Proposition 8.9.1.2**. 

018H 8.8.6 Natural Isomorphisms

Let C and \mathcal{D} be categories and let $F, G: C \rightrightarrows \mathcal{D}$ be functors.

018J DEFINITION 8.8.6.1 ► NATURAL ISOMORPHISMS

A natural transformation $\alpha: F \Rightarrow G$ is a **natural isomorphism** if there exists a natural transformation $\alpha^{-1}: G \Rightarrow F$ such that

$$\begin{aligned}\alpha^{-1} \circ \alpha &= \text{id}_F, \\ \alpha \circ \alpha^{-1} &= \text{id}_G.\end{aligned}$$

018K PROPOSITION 8.8.6.2 ► PROPERTIES OF NATURAL ISOMORPHISMS

Let $\alpha: F \Rightarrow G$ be a natural transformation.

- 018L 1. *Characterisations.* The following conditions are equivalent:
- 018M (a) The natural transformation α is a natural isomorphism.
- 018N (b) For each $A \in \text{Obj}(C)$, the morphism $\alpha_A: F_A \rightarrow G_A$ is an isomorphism.
- 018P 2. *Componentwise Inverses of Natural Transformations Assemble Into Natural Transformations.* Let $\alpha^{-1}: G \Rightarrow F$ be a transformation such that, for each $A \in \text{Obj}(C)$, we have

$$\begin{aligned}\alpha_A^{-1} \circ \alpha_A &= \text{id}_{F(A)}, \\ \alpha_A \circ \alpha_A^{-1} &= \text{id}_{G(A)}.\end{aligned}$$

Then α^{-1} is a natural transformation.

PROOF 8.8.6.3 ► PROOF OF PROPOSITION 8.8.6.2

Item 1: Characterisations

The implication **Item 1a** \Rightarrow **Item 1b** is clear, whereas the implication **Item 1b** \Rightarrow **Item 1a** follows from **Item 2**.

Item 2: Componentwise Inverses of Natural Transformations Assemble Into

The naturality condition for α^{-1} corresponds to the commutativity of the diagram

$$\begin{array}{ccc} G(A) & \xrightarrow{G(f)} & G(B) \\ \alpha_A^{-1} \downarrow & & \downarrow \alpha_B^{-1} \\ F(A) & \xrightarrow{F(f)} & F(B) \end{array}$$

for each $A, B \in \text{Obj}(C)$ and each $f \in \text{Hom}_C(A, B)$. Considering the diagram


$$\begin{array}{ccc}
 G(A) & \xrightarrow{G(f)} & G(B) \\
 \alpha_A^{-1} \downarrow & (1) & \downarrow \alpha_B^{-1} \\
 F(A) & \xrightarrow{F(f)} & F(B) \\
 \alpha_A \downarrow & (2) & \downarrow \alpha_B \\
 G(A) & \xrightarrow{G(f)} & G(B),
 \end{array}$$

where the boundary diagram as well as Subdiagram (2) commute, we have

$$\begin{aligned}
 G(f) &= G(f) \circ \text{id}_{G(A)} \\
 &= G(f) \circ \alpha_A \circ \alpha_A^{-1} \\
 &= \alpha_B \circ F(f) \circ \alpha_A^{-1}.
 \end{aligned}$$

Postcomposing both sides with α_B^{-1} , we get

$$\begin{aligned}
 \alpha_B^{-1} \circ G(f) &= \alpha_B^{-1} \circ \alpha_B \circ F(f) \circ \alpha_A^{-1} \\
 &= \text{id}_{F(B)} \circ F(f) \circ \alpha_A^{-1} \\
 &= F(f) \circ \alpha_A^{-1},
 \end{aligned}$$

which is the naturality condition we wanted to show. Thus α^{-1} is a natural transformation. 

018Q 8.9 Categories of Categories

018R 8.9.1 Functor Categories

Let C be a category and \mathcal{D} be a small category.

018S DEFINITION 8.9.1.1 ► FUNCTOR CATEGORIES

The **category of functors from C to \mathcal{D}** ¹ is the category $\text{Fun}(C, \mathcal{D})$ ² where

- *Objects.* The objects of $\text{Fun}(C, \mathcal{D})$ are functors from C to \mathcal{D} .
- *Morphisms.* For each $F, G \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$, we have

$$\text{Hom}_{\text{Fun}(C, \mathcal{D})}(F, G) \stackrel{\text{def}}{=} \text{Nat}(F, G).$$

- *Identities.* For each $F \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$, the unit map

$$\mathbb{1}_F^{\text{Fun}(C, \mathcal{D})} : \text{pt} \rightarrow \text{Nat}(F, F)$$

of $\text{Fun}(C, \mathcal{D})$ at F is given by

$$\text{id}_F^{\text{Fun}(C, \mathcal{D})} \stackrel{\text{def}}{=} \text{id}_F,$$

where $\text{id}_F : F \implies F$ is the identity natural transformation of F of [Example 8.8.2.4](#).

- *Composition.* For each $F, G, H \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$, the composition map

$$\circ_{F, G, H}^{\text{Fun}(C, \mathcal{D})} : \text{Nat}(G, H) \times \text{Nat}(F, G) \rightarrow \text{Nat}(F, H)$$

of $\text{Fun}(C, \mathcal{D})$ at (F, G, H) is given by

$$\beta \circ_{F, G, H}^{\text{Fun}(C, \mathcal{D})} \alpha \stackrel{\text{def}}{=} \beta \circ \alpha,$$

where $\beta \circ \alpha$ is the vertical composition of α and β of [Item 1 of Proposition 8.8.3.3](#).

¹*Further Terminology:* Also called the **functor category** $\text{Fun}(C, \mathcal{D})$.

²*Further Notation:* Also written \mathcal{D}^C and $[C, \mathcal{D}]$.

018T PROPOSITION 8.9.1.2 ► PROPERTIES OF FUNCTOR CATEGORIES

Let C and \mathcal{D} be categories and let $F : C \rightarrow \mathcal{D}$ be a functor.

- 018U 1. *Functoriality.* The assignments $C, \mathcal{D}, (C, \mathcal{D}) \mapsto \text{Fun}(C, \mathcal{D})$ define functors

$$\begin{aligned} \text{Fun}(C, -_2) &: \text{Cats} \rightarrow \text{Cats}, \\ \text{Fun}(-_1, \mathcal{D}) &: \text{Cats}^{\text{op}} \rightarrow \text{Cats}, \\ \text{Fun}(-_1, -_2) &: \text{Cats}^{\text{op}} \times \text{Cats} \rightarrow \text{Cats}. \end{aligned}$$

- 018V 2. *2-Functoriality.* The assignments $C, \mathcal{D}, (C, \mathcal{D}) \mapsto \text{Fun}(C, \mathcal{D})$ define 2-functors

$$\begin{aligned} \text{Fun}(C, -_2) &: \text{Cats}_2 \rightarrow \text{Cats}_2, \\ \text{Fun}(-_1, \mathcal{D}) &: \text{Cats}_2^{\text{op}} \rightarrow \text{Cats}_2, \\ \text{Fun}(-_1, -_2) &: \text{Cats}_2^{\text{op}} \times \text{Cats}_2 \rightarrow \text{Cats}_2. \end{aligned}$$

018W

3. *Adjointness.* We have adjunctions

$$(C \times - \dashv \text{Fun}(C, -)): \text{Cats} \begin{array}{c} \xrightarrow{C \times -} \\ \perp \\ \xleftarrow{\text{Fun}(C, -)} \end{array} \text{Cats},$$

$$(- \times \mathcal{D} \dashv \text{Fun}(\mathcal{D}, -)): \text{Cats} \begin{array}{c} \xrightarrow{- \times \mathcal{D}} \\ \perp \\ \xleftarrow{\text{Fun}(\mathcal{D}, -)} \end{array} \text{Cats},$$

witnessed by bijections of sets

$$\text{Hom}_{\text{Cats}}(C \times \mathcal{D}, \mathcal{E}) \cong \text{Hom}_{\text{Cats}}(\mathcal{D}, \text{Fun}(C, \mathcal{E})),$$

$$\text{Hom}_{\text{Cats}}(C \times \mathcal{D}, \mathcal{E}) \cong \text{Hom}_{\text{Cats}}(C, \text{Fun}(\mathcal{D}, \mathcal{E})),$$

natural in $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats})$.

018X

4. *2-Adjointness.* We have 2-adjunctions

$$(C \times - \dashv \text{Fun}(C, -)): \text{Cats}_2 \begin{array}{c} \xrightarrow{C \times -} \\ \perp_2 \\ \xleftarrow{\text{Fun}(C, -)} \end{array} \text{Cats}_2,$$

$$(- \times \mathcal{D} \dashv \text{Fun}(\mathcal{D}, -)): \text{Cats}_2 \begin{array}{c} \xrightarrow{- \times \mathcal{D}} \\ \perp_2 \\ \xleftarrow{\text{Fun}(\mathcal{D}, -)} \end{array} \text{Cats}_2,$$

witnessed by isomorphisms of categories

$$\text{Fun}(C \times \mathcal{D}, \mathcal{E}) \cong \text{Fun}(\mathcal{D}, \text{Fun}(C, \mathcal{E})),$$

$$\text{Fun}(C \times \mathcal{D}, \mathcal{E}) \cong \text{Fun}(C, \text{Fun}(\mathcal{D}, \mathcal{E})),$$

natural in $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats}_2)$.

018Y

5. *Interaction With Punctual Categories.* We have a canonical isomorphism of categories

$$\text{Fun}(\text{pt}, C) \cong C,$$

natural in $C \in \text{Obj}(\text{Cats})$.

018Z

6. *Objectwise Computation of Co/Limits.* Let

$$D: \mathcal{I} \rightarrow \text{Fun}(C, \mathcal{D})$$

be a diagram in $\text{Fun}(C, \mathcal{D})$. We have isomorphisms

$$\begin{aligned}\lim(D)_A &\cong \lim_{i \in I}(D_i(A)), \\ \text{colim}(D)_A &\cong \text{colim}_{i \in I}(D_i(A)),\end{aligned}$$

naturally in $A \in \text{Obj}(C)$.

0190 7. *Interaction With Co/Completeness.* If \mathcal{E} is co/complete, then so is $\text{Fun}(C, \mathcal{E})$.

0191 8. *Monomorphisms and Epimorphisms.* Let $\alpha: F \Rightarrow G$ be a morphism of $\text{Fun}(C, \mathcal{D})$. The following conditions are equivalent:

0192 (a) The natural transformation

$$\alpha: F \Rightarrow G$$

is a monomorphism (resp. epimorphism) in $\text{Fun}(C, \mathcal{D})$.

0193 (b) For each $A \in \text{Obj}(C)$, the morphism

$$\alpha_A: F_A \rightarrow G_A$$

is a monomorphism (resp. epimorphism) in \mathcal{D} .

PROOF 8.9.1.3 ► PROOF OF PROPOSITION 8.9.1.2

Item 1: Functoriality

Omitted.

Item 2: 2-Functoriality

Omitted.

Item 3: Adjointness

Omitted.

Item 4: 2-Adjointness

Omitted.

Item 5: Interaction With Punctual Categories

Omitted.

Item 6: Objectwise Computation of Co/Limits

Omitted.

Item 7: Interaction With Co/Completeness

This follows from ??.

Item 8: Monomorphisms and Epimorphisms

Omitted. 

0194 8.9.2 The Category of Categories and Functors

0195 DEFINITION 8.9.2.1 ► THE CATEGORY OF CATEGORIES AND FUNCTORS

The **category of (small) categories and functors** is the category \mathbf{Cats} where

- *Objects.* The objects of \mathbf{Cats} are small categories.
- *Morphisms.* For each $C, D \in \text{Obj}(\mathbf{Cats})$, we have

$$\text{Hom}_{\mathbf{Cats}}(C, D) \stackrel{\text{def}}{=} \text{Obj}(\text{Fun}(C, D)).$$

- *Identities.* For each $C \in \text{Obj}(\mathbf{Cats})$, the unit map

$$\mathbb{1}_C^{\mathbf{Cats}} : \text{pt} \rightarrow \text{Hom}_{\mathbf{Cats}}(C, C)$$

of \mathbf{Cats} at C is defined by

$$\text{id}_C^{\mathbf{Cats}} \stackrel{\text{def}}{=} \text{id}_C,$$

where $\text{id}_C : C \rightarrow C$ is the identity functor of C of [Example 8.4.1.4](#).

- *Composition.* For each $C, D, E \in \text{Obj}(\mathbf{Cats})$, the composition map

$$\circ_{C,D,E}^{\mathbf{Cats}} : \text{Hom}_{\mathbf{Cats}}(D, E) \times \text{Hom}_{\mathbf{Cats}}(C, D) \rightarrow \text{Hom}_{\mathbf{Cats}}(C, E)$$

of \mathbf{Cats} at (C, D, E) is given by

$$G \circ_{C,D,E}^{\mathbf{Cats}} F \stackrel{\text{def}}{=} G \circ F,$$

where $G \circ F : C \rightarrow E$ is the composition of F and G of [Definition 8.4.1.6](#).

0196 PROPOSITION 8.9.2.2 ► PROPERTIES OF THE CATEGORY \mathbf{Cats}

Let C be a category.

- 0197 1. *Co/Completeness.* The category \mathbf{Cats} is complete and cocomplete.

0198

2. *Cartesian Monoidal Structure.* The quadruple $(\text{Cats}, \times, \text{pt}, \text{Fun})$ is a Cartesian closed monoidal category.

PROOF 8.9.2.3 ► PROOF OF PROPOSITION 8.9.2.2

Item 1: Co/Completeness

Omitted.

Item 2: Cartesian Monoidal Structure

Omitted. 

8.9.3 The 2-Category of Categories, Functors, and Natural Transformations

0199

019A

DEFINITION 8.9.3.1 ► THE 2-CATEGORY OF CATEGORIES

The **2-category of (small) categories, functors, and natural transformations** is the 2-category Cats_2 where

- *Objects.* The objects of Cats_2 are small categories.
- *Hom-Categories.* For each $C, \mathcal{D} \in \text{Obj}(\text{Cats}_2)$, we have

$$\text{Hom}_{\text{Cats}_2}(C, \mathcal{D}) \stackrel{\text{def}}{=} \text{Fun}(C, \mathcal{D}).$$

- *Identities.* For each $C \in \text{Obj}(\text{Cats}_2)$, the unit functor

$$\mathbb{1}_C^{\text{Cats}_2} : \text{pt} \rightarrow \text{Fun}(C, C)$$

of Cats_2 at C is the functor picking the identity functor $\text{id}_C : C \rightarrow C$ of C .

- *Composition.* For each $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats}_2)$, the composition bifunctor

$$\circ_{C, \mathcal{D}, \mathcal{E}}^{\text{Cats}_2} : \text{Hom}_{\text{Cats}_2}(\mathcal{D}, \mathcal{E}) \times \text{Hom}_{\text{Cats}_2}(C, \mathcal{D}) \rightarrow \text{Hom}_{\text{Cats}_2}(C, \mathcal{E})$$

of Cats_2 at $(C, \mathcal{D}, \mathcal{E})$ is the functor where

- *Action on Objects.* For each object $(G, F) \in \text{Obj}(\text{Hom}_{\text{Cats}_2}(\mathcal{D}, \mathcal{E}) \times \text{Hom}_{\text{Cats}_2}(C, \mathcal{D}))$, we have

$$\circ_{C, \mathcal{D}, \mathcal{E}}^{\text{Cats}_2}(G, F) \stackrel{\text{def}}{=} G \circ F.$$

– *Action on Morphisms.* For each morphism $(\beta, \alpha): (K, H) \Rightarrow (G, F)$ of $\text{Hom}_{\text{Cats}_2}(\mathcal{D}, \mathcal{E}) \times \text{Hom}_{\text{Cats}_2}(C, \mathcal{D})$, we have

$$\circ_{C, \mathcal{D}, \mathcal{E}}^{\text{Cats}_2}(\beta, \alpha) \stackrel{\text{def}}{=} \beta \star \alpha,$$

where $\beta \star \alpha$ is the horizontal composition of α and β of [Definition 8.8.4.1](#).

019B **PROPOSITION 8.9.3.2 ▶ PROPERTIES OF THE 2-CATEGORY Cats_2**

Let C be a category.

- 019C 1. *2-Categorical Co/Completeness.* The 2-category Cats_2 is complete and cocomplete as a 2-category, having all 2-categorical and bicategorical co/limits.

PROOF 8.9.3.3 ▶ PROOF OF PROPOSITION 8.9.3.2

Item 1: Co/Completeness

Omitted. 

019D **8.9.4 The Category of Groupoids**

019E **DEFINITION 8.9.4.1 ▶ THE CATEGORY OF SMALL GROUPOIDS**

The **category of (small) groupoids** is the full subcategory Grpd of Cats spanned by the groupoids.

019F **8.9.5 The 2-Category of Groupoids**

019G **DEFINITION 8.9.5.1 ▶ THE 2-CATEGORY OF SMALL GROUPOIDS**

The **2-category of (small) groupoids** is the full sub-2-category Grpd_2 of Cats_2 spanned by the groupoids.

Appendices

8.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets

Relations

5. Relations

6. Constructions With Relations

7. Equivalence Relations and Apartness Relations

Category Theory

8. Categories

Bicategories

9. Types of Morphisms in Bicategories

Part IV

Bicategories

Chapter 9

Types of Morphisms in Bicategories

Ø19H In this chapter, we study special kinds of morphisms in bicategories:

1. *Monomorphisms and Epimorphisms in Bicategories* (Sections 9.1 and 9.2). There is a large number of different notions capturing the idea of a “monomorphism” or of an “epimorphism” in a bicategory.

Arguably, the notion that best captures these concepts is that of a *pseudomonoid morphism* (Definition 9.1.10.1) and of a *pseudoepic morphism* (Definition 9.2.10.1), although the other notions introduced in Sections 9.1 and 9.2 are also interesting on their own.

Contents

9.1	Monomorphisms in Bicategories	516
9.1.1	Representably Faithful Morphisms	516
9.1.2	Representably Full Morphisms	517
9.1.3	Representably Fully Faithful Morphisms	518
9.1.4	Morphisms Representably Faithful on Cores	519
9.1.5	Morphisms Representably Full on Cores	520
9.1.6	Morphisms Representably Fully Faithful on Cores ...	521
9.1.7	Representably Essentially Injective Morphisms	522
9.1.8	Representably Conservative Morphisms	523
9.1.9	Strict Monomorphisms	524
9.1.10	Pseudomonoid Morphisms	525
9.2	Epimorphisms in Bicategories	527
9.2.1	Corepresentably Faithful Morphisms	527
9.2.2	Corepresentably Full Morphisms	528
9.2.3	Corepresentably Fully Faithful Morphisms	529
9.2.4	Morphisms Corepresentably Faithful on Cores	530
9.2.5	Morphisms Corepresentably Full on Cores	531

9.2.6	Morphisms Corepresentably Fully Faithful on Cores.	532
9.2.7	Corepresentably Essentially Injective Morphisms	533
9.2.8	Corepresentably Conservative Morphisms	534
9.2.9	Strict Epimorphisms	535
9.2.10	Pseudoepic Morphisms	536
9.A	Other Chapters	538

019J **9.1 Monomorphisms in Bicategories**

019K **9.1.1 Representably Faithful Morphisms**

Let C be a bicategory.

019L **DEFINITION 9.1.1.1 ► REPRESENTABLY FAITHFUL MORPHISMS**

A 1-morphism $f: A \rightarrow B$ of C is **representably faithful**¹ if, for each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is faithful.

¹*Further Terminology:* Also called simply a **faithful morphism**, based on [Item 1 of Example 9.1.1.3](#).

019M **REMARK 9.1.1.2 ► UNWINDING DEFINITION 9.1.1.1**

In detail, f is representably faithful if, for all diagrams in C of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \Downarrow \beta \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then $\alpha = \beta$.

019N **EXAMPLE 9.1.1.3 ► EXAMPLES OF REPRESENTABLY FAITHFUL MORPHISMS**

Here are some examples of representably faithful morphisms.

- 019P 1. *Representably Faithful Morphisms in Cats_2* . The representably faithful

morphisms in \mathbf{Cats}_2 are precisely the faithful functors; see [Item 1](#) of [Proposition 8.5.1.2](#).

019Q

2. *Representably Faithful Morphisms in Rel.* Every morphism of **Rel** is representably faithful; see [Item 1](#) of [Proposition 5.3.8.1](#).

019R **9.1.2 Representably Full Morphisms**

Let C be a bicategory.

019S

DEFINITION 9.1.2.1 ▶ REPRESENTABLY FULL MORPHISMS

A 1-morphism $f: A \rightarrow B$ of C is **representably full**¹ if, for each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is full.

¹*Further Terminology:* Also called simply a **full morphism**, based on [Item 1](#) of [Example 9.1.2.3](#).

019T

REMARK 9.1.2.2 ▶ UNWINDING DEFINITION 9.1.2.1

In detail, f is representably full if, for each $X \in \text{Obj}(C)$ and each 2-morphism

$$\beta: f \circ \phi \Rightarrow f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of C , there exists a 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of C such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

019U **EXAMPLE 9.1.2.3 ▶ EXAMPLES OF REPRESENTABLY FULL MORPHISMS**

Here are some examples of representably full morphisms.

- 019V 1. *Representably Full Morphisms in \mathbf{Cats}_2* . The representably full morphisms in \mathbf{Cats}_2 are precisely the full functors; see [Item 1](#) of [Proposition 8.5.2.2](#).
- 019W 2. *Representably Full Morphisms in \mathbf{Rel}* . The representably full morphisms in \mathbf{Rel} are characterised in [Item 2](#) of [Proposition 5.3.8.1](#).

019X **9.1.3 Representably Fully Faithful Morphisms**

Let C be a bicategory.

019Y **DEFINITION 9.1.3.1 ▶ REPRESENTABLY FULLY FAITHFUL MORPHISMS**

A 1-morphism $f : A \rightarrow B$ of C is **representably fully faithful**¹ if the following equivalent conditions are satisfied:

- 019Z 1. The 1-morphism f is representably faithful ([Definition 9.1.1.1](#)) and representably full ([Definition 9.1.2.1](#)).
- 01A0 2. For each $X \in \text{Obj}(C)$, the functor

$$f_* : \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is fully faithful.

¹*Further Terminology:* Also called simply a **fully faithful morphism**, based on [Item 1](#) of [Example 9.1.3.3](#).

01A1 **REMARK 9.1.3.2 ▶ UNWINDING REPRESENTABLY FULLY FAITHFUL MORPHISMS**

In detail, f is representably fully faithful if the conditions in [Remark 9.1.1.2](#) and [Remark 9.1.2.2](#) hold:

- 1. For all diagrams in C of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \downarrow \downarrow \downarrow \beta \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then $\alpha = \beta$.

2. For each $X \in \text{Obj}(C)$ and each 2-morphism

$$\beta: f \circ \phi \Longrightarrow f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of C , there exists a 2-morphism

$$\alpha: \phi \Longrightarrow \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of C such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

01A2 **EXAMPLE 9.1.3 ► EXAMPLES OF REPRESENTABLY FULLY FAITHFUL MORPHISMS**

Here are some examples of representably fully faithful morphisms.

- 01A3 1. *Representably Fully Faithful Morphisms in \mathbf{Cats}_2* . The representably fully faithful morphisms in \mathbf{Cats}_2 are precisely the fully faithful functors; see [Item 5 of Proposition 8.5.3.2](#).
- 01A4 2. *Representably Fully Faithful Morphisms in \mathbf{Rel}* . The representably fully faithful morphisms of \mathbf{Rel} coincide ([Item 3 of Proposition 5.3.8.1](#)) with the representably full morphisms in \mathbf{Rel} , which are characterised in [Item 2 of Proposition 5.3.8.1](#).

01A5 **9.1.4 Morphisms Representably Faithful on Cores**

Let C be a bicategory.

01A6 DEFINITION 9.1.4.1 ► MORPHISMS REPRESENTABLY FAITHFUL ON CORES

A 1-morphism $f: A \rightarrow B$ of C is **representably faithful on cores** if, for each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Core}(\text{Hom}_C(X, A)) \rightarrow \text{Core}(\text{Hom}_C(X, B))$$

given by postcomposition by f is faithful.

01A7 REMARK 9.1.4.2 ► UNWINDING DEFINITION 9.1.4.1

In detail, f is representably faithful on cores if, for all diagrams in C of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \beta \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if α and β are 2-isomorphisms and we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then $\alpha = \beta$.

01A8 9.1.5 Morphisms Representably Full on Cores

Let C be a bicategory.

01A9 DEFINITION 9.1.5.1 ► MORPHISMS REPRESENTABLY FULL ON CORES

A 1-morphism $f: A \rightarrow B$ of C is **representably full on cores** if, for each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Core}(\text{Hom}_C(X, A)) \rightarrow \text{Core}(\text{Hom}_C(X, B))$$

given by postcomposition by f is full.

01AA REMARK 9.1.5.2 ► UNWINDING DEFINITION 9.1.5.1

In detail, f is representably full on cores if, for each $X \in \text{Obj}(C)$ and each 2-isomorphism

$$\beta: f \circ \phi \xrightarrow{\sim} f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of C , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of C such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

01AB 9.1.6 Morphisms Representably Fully Faithful on Cores

Let C be a bicategory.

01AC DEFINITION 9.1.6.1 ► MORPHISMS REPRESENTABLY FULLY FAITHFUL ON CORES

A 1-morphism $f: A \rightarrow B$ of C is **representably fully faithful on cores** if the following equivalent conditions are satisfied:

- 01AD 1. The 1-morphism f is representably faithful on cores (Definition 9.1.5.1) and representably full on cores (Definition 9.1.4.1).
- 01AE 2. For each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Core}(\text{Hom}_C(X, A)) \rightarrow \text{Core}(\text{Hom}_C(X, B))$$

given by postcomposition by f is fully faithful.

01AF

REMARK 9.1.6.2 ► UNWINDING DEFINITION 9.1.6.1

In detail, f is representably fully faithful on cores if the conditions in Remark 9.1.4.2 and Remark 9.1.5.2 hold:

1. For all diagrams in C of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \beta \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if α and β are 2-isomorphisms and we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then $\alpha = \beta$.

2. For each $X \in \text{Obj}(C)$ and each 2-isomorphism

$$\beta: f \circ \phi \xrightarrow{\sim} f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of C , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of C such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

01AG 9.1.7 Representably Essentially Injective Morphisms

Let C be a bicategory.

01AH DEFINITION 9.1.7.1 ► REPRESENTABLY ESSENTIALLY INJECTIVE MORPHISMS

A 1-morphism $f: A \rightarrow B$ of C is **representably essentially injective** if, for each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is essentially injective.

01AJ REMARK 9.1.7.2 ► UNWINDING DEFINITION 9.1.7.1

In detail, f is representably essentially injective if, for each pair of morphisms $\phi, \psi: X \rightrightarrows A$ of C , the following condition is satisfied:

$$(\star) \text{ If } f \circ \phi \cong f \circ \psi, \text{ then } \phi \cong \psi.$$

01AK 9.1.8 Representably Conservative Morphisms

Let C be a bicategory.

01AL DEFINITION 9.1.8.1 ► REPRESENTABLY CONSERVATIVE MORPHISMS

A 1-morphism $f: A \rightarrow B$ of C is **representably conservative** if, for each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is conservative.

01AM REMARK 9.1.8.2 ► UNWINDING DEFINITION 9.1.8.1

In detail, f is representably conservative if, for each pair of morphisms $\phi, \psi: X \rightrightarrows A$ and each 2-morphism

$$\alpha: \phi \rightrightarrows \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of C , if the 2-morphism

$$\text{id}_f \star \alpha: f \circ \phi \implies f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \parallel \\ \text{id}_f \star \alpha \\ \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

is a 2-isomorphism, then so is α .

01AN 9.1.9 Strict Monomorphisms

Let C be a bicategory.

01AP DEFINITION 9.1.9.1 ▶ STRICT MONOMORPHISMS

A 1-morphism $f: A \rightarrow B$ of C is a **strict monomorphism** if, for each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is injective on objects, i.e. its action on objects

$$f_*: \text{Obj}(\text{Hom}_C(X, A)) \rightarrow \text{Obj}(\text{Hom}_C(X, B))$$

is injective.

01AQ REMARK 9.1.9.2 ▶ UNWINDING DEFINITION 9.1.9.1

In detail, f is a strict monomorphism in C if, for each diagram in C of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if $f \circ \phi = f \circ \psi$, then $\phi = \psi$.

01AR EXAMPLE 9.1.9.3 ▶ EXAMPLES OF STRICT MONOMORPHISMS

Here are some examples of strict monomorphisms.

- 01AS**
1. *Strict Monomorphisms in Cats_2* . The strict monomorphisms in Cats_2 are precisely the functors which are injective on objects and injective on morphisms; see [Item 1](#) of [Proposition 8.6.2.2](#).

01AT 2. *Strict Monomorphisms in Rel.* The strict monomorphisms in **Rel** are characterised in [Proposition 5.3.7.1](#).

01AU **9.1.10 Pseudomonic Morphisms**

Let C be a bicategory.

01AV **DEFINITION 9.1.10.1 ► PSEUDOMONIC MORPHISMS**

A 1-morphism $f: A \rightarrow B$ of C is **pseudomonic** if, for each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is pseudomonic.

01AW **REMARK 9.1.10.2 ► UNWINDING DEFINITION 9.1.10.1**

In detail, a 1-morphism $f: A \rightarrow B$ of C is pseudomonic if it satisfies the following conditions:

01AX 1. For all diagrams in C of the form

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \beta \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B,$$

if we have

$$\text{id}_f \star \alpha = \text{id}_f \star \beta,$$

then $\alpha = \beta$.

01AY 2. For each $X \in \text{Obj}(C)$ and each 2-isomorphism

$$\beta: f \circ \phi \xrightarrow{\sim} f \circ \psi, \quad X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of C , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A$$

of C such that we have an equality

$$X \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} A \xrightarrow{f} B = X \begin{array}{c} \xrightarrow{f \circ \phi} \\ \beta \Downarrow \\ \xrightarrow{f \circ \psi} \end{array} B$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \text{id}_f \star \alpha.$$

01AZ

PROPOSITION 9.1.10.3 ► PROPERTIES OF PSEUDOMONIC MORPHISMS

Let $f: A \rightarrow B$ be a 1-morphism of C .

01B0

1. *Characterisations.* The following conditions are equivalent:

01B1

(a) The morphism f is pseudomonic.

01B2

(b) The morphism f is representably full on cores and representably faithful.

01B3

(c) We have an isocomma square of the form

$$A \overset{\text{eq.}}{\cong} A \times_B A, \quad \begin{array}{ccc} A & \xrightarrow{\text{id}_A} & A \\ \text{id}_A \downarrow & \swarrow \text{dashed} & \downarrow F \\ A & \xrightarrow{F} & B \end{array}$$

in C up to equivalence.

01B4

2. *Interaction With Cotensors.* If C has cotensors with $\mathbb{1}$, then the following conditions are equivalent:

(a) The morphism f is pseudomonic.

(b) We have an isocomma square of the form

$$A \overset{\text{eq.}}{\cong} A \times_{\mathbb{1} \pitchfork F} B, \quad \begin{array}{ccc} A & \xrightarrow{\quad} & \mathbb{1} \pitchfork A \\ F \downarrow & \swarrow \text{dashed} & \downarrow \mathbb{1} \pitchfork F \\ B & \xrightarrow{\quad} & \mathbb{1} \pitchfork B \end{array}$$

in C up to equivalence.

PROOF 9.1.10.4 ► PROOF OF PROPOSITION 9.1.10.3

Item 1: Characterisations

Omitted.

Item 2: Interaction With Cotensors

Omitted. 

01B5 9.2 Epimorphisms in Bicategories

01B6 9.2.1 Corepresentably Faithful Morphisms

Let C be a bicategory.

01B7 DEFINITION 9.2.1.1 ► COREPRESENTABLY FAITHFUL MORPHISMS

A 1-morphism $f: A \rightarrow B$ of C is **corepresentably faithful** if, for each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is faithful.

01B8 REMARK 9.2.1.2 ► UNWINDING DEFINITION 9.2.1.1

In detail, f is corepresentably faithful if, for all diagrams in C of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \downarrow \downarrow \downarrow \beta \\ \xrightarrow{\psi} \end{array} X,$$

if we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then $\alpha = \beta$.

01B9 EXAMPLE 9.2.1.3 ► EXAMPLES OF COREPRESENTABLY FAITHFUL MORPHISMS

Here are some examples of corepresentably faithful morphisms.

- 01BA 1. *Corepresentably Faithful Morphisms in Cats_2 .* The corepresentably faithful morphisms in Cats_2 are characterised in [Item 4 of Proposition 8.5.1.2](#).

01BB 2. *Corepresentably Faithful Morphisms in Rel.* Every morphism of **Rel** is corepresentably faithful; see [Item 1](#) of [Proposition 5.3.10.1](#).

01BC **9.2.2 Corepresentably Full Morphisms**

Let C be a bicategory.

01BD **DEFINITION 9.2.2.1 ► COREPRESENTABLY FULL MORPHISMS**

A 1-morphism $f: A \rightarrow B$ of C is **corepresentably full** if, for each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is full.

01BE **REMARK 9.2.2.2 ► UNWINDING DEFINITION 9.2.2.1**

In detail, f is corepresentably full if, for each $X \in \text{Obj}(C)$ and each 2-morphism

$$\beta: \phi \circ f \Longrightarrow \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of C , there exists a 2-morphism

$$\alpha: \phi \Longrightarrow \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of C such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

01BF **EXAMPLE 9.2.2.3 ► EXAMPLES OF COREPRESENTABLY FULL MORPHISMS**

Here are some examples of corepresentably full morphisms.

- 01BG 1. *Corepresentably Full Morphisms in \mathbf{Cats}_2* . The corepresentably full morphisms in \mathbf{Cats}_2 are characterised in **Item 5** of **Proposition 8.5.2.2**.
- 01BH 2. *Corepresentably Full Morphisms in \mathbf{Rel}* . The corepresentably full morphisms in \mathbf{Rel} are characterised in ?? of **Proposition 5.3.8.1**.

01BJ **9.2.3 Corepresentably Fully Faithful Morphisms**

Let C be a bicategory.

01BK **DEFINITION 9.2.3.1 ► COREPRESENTABLY FULLY FAITHFUL MORPHISMS**

A 1-morphism $f: A \rightarrow B$ of C is **corepresentably fully faithful**¹ if the following equivalent conditions are satisfied:

- 01BL 1. The 1-morphism f is corepresentably full (**Definition 9.2.2.1**) and corepresentably faithful (**Definition 9.2.1.1**).
- 01BM 2. For each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is fully faithful.

¹*Further Terminology:* Corepresentably fully faithful morphisms have also been called **lax epimorphisms** in the literature (e.g. in [Adá+01]), though we will always use the name “corepresentably fully faithful morphism” instead in this work.

01BN **REMARK 9.2.3.2 ► UNWINDING DEFINITION 9.2.3.1**

In detail, f is corepresentably fully faithful if the conditions in **Remark 9.2.1.2** and **Remark 9.2.2.2** hold:

- 1. For all diagrams in C of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \downarrow \downarrow \downarrow \beta \\ \xrightarrow{\psi} \end{array} X,$$

if we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then $\alpha = \beta$.

2. For each $X \in \text{Obj}(C)$ and each 2-morphism

$$\beta: \phi \circ f \Rightarrow \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of C , there exists a 2-morphism

$$\alpha: \phi \Rightarrow \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of C such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

01BP EXAMPLE 9.2.3.3 ► EXAMPLES OF COREPRESENTABLY FULLY FAITHFUL MORPHISMS

Here are some examples of corepresentably fully faithful morphisms.

- 01BQ** 1. *Corepresentably Fully Faithful Morphisms in \mathbf{Cats}_2* . The fully faithful epimorphisms in \mathbf{Cats}_2 are characterised in **Item 9** of **Proposition 8.5.3.2**.
- 01BR** 2. *Corepresentably Fully Faithful Morphisms in \mathbf{Rel}* . The corepresentably fully faithful morphisms of \mathbf{Rel} coincide (**Item 3** of **Proposition 5.3.10.1**) with the corepresentably full morphisms in \mathbf{Rel} , which are characterised in **Item 2** of **Proposition 5.3.10.1**.

01BS 9.2.4 Morphisms Corepresentably Faithful on Cores

Let C be a bicategory.

01BT

DEFINITION 9.2.4.1 ► MORPHISMS COREPRESENTABLY FAITHFUL ON CORES

A 1-morphism $f: A \rightarrow B$ of C is **corepresentably faithful on cores** if, for each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Core}(\text{Hom}_C(B, X)) \rightarrow \text{Core}(\text{Hom}_C(A, X))$$

given by precomposition by f is faithful.

01BU

REMARK 9.2.4.2 ► UNWINDING DEFINITION 9.2.4.1

In detail, f is corepresentably faithful on cores if, for all diagrams in C of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \downarrow \Downarrow \beta \\ \xrightarrow{\psi} \end{array} X,$$

if α and β are 2-isomorphisms and we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then $\alpha = \beta$.

01BV 9.2.5 Morphisms Corepresentably Full on Cores

Let C be a bicategory.

01BW

DEFINITION 9.2.5.1 ► MORPHISMS COREPRESENTABLY FULL ON CORES

A 1-morphism $f: A \rightarrow B$ of C is **corepresentably full on cores** if, for each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Core}(\text{Hom}_C(B, X)) \rightarrow \text{Core}(\text{Hom}_C(A, X))$$

given by precomposition by f is full.

01BX

REMARK 9.2.5.2 ► UNWINDING DEFINITION 9.2.5.1

In detail, f is corepresentably full on cores if, for each $X \in \text{Obj}(C)$ and each 2-isomorphism

$$\beta: \phi \circ f \xrightarrow{\sim} \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of C , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of C such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

01BY 9.2.6 Morphisms Corepresentably Fully Faithful on Cores

Let C be a bicategory.

01BZ DEFINITION 9.2.6.1 ► MORPHISMS COREPRESENTABLY FULLY FAITHFUL ON CORES

A 1-morphism $f: A \rightarrow B$ of C is **corepresentably fully faithful on cores** if the following equivalent conditions are satisfied:

- 01C0 1. The 1-morphism f is corepresentably full on cores (Definition 9.2.5.1) and corepresentably faithful on cores (Definition 9.2.1.1).
- 01C1 2. For each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Core}(\text{Hom}_C(B, X)) \rightarrow \text{Core}(\text{Hom}_C(A, X))$$

given by precomposition by f is fully faithful.

01C2

REMARK 9.2.6.2 ► UNWINDING DEFINITION 9.2.6.1

In detail, f is corepresentably fully faithful on cores if the conditions in Remark 9.2.4.2 and Remark 9.2.5.2 hold:

1. For all diagrams in C of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \Downarrow \beta \\ \xrightarrow{\psi} \end{array} X,$$

if α and β are 2-isomorphisms and we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then $\alpha = \beta$.

2. For each $X \in \text{Obj}(C)$ and each 2-isomorphism

$$\beta: \phi \circ f \xrightarrow{\sim} \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of C , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of C such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

01C3 9.2.7 Corepresentably Essentially Injective Morphisms

Let C be a bicategory.

01C4 DEFINITION 9.2.7.1 ► COREPRESENTABLY ESSENTIALLY INJECTIVE MORPHISMS

A 1-morphism $f: A \rightarrow B$ of C is **corepresentably essentially injective** if, for each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is essentially injective.

01C5 REMARK 9.2.7.2 ► UNWINDING DEFINITION 9.2.7.1

In detail, f is corepresentably essentially injective if, for each pair of morphisms $\phi, \psi: B \rightrightarrows X$ of C , the following condition is satisfied:

$$(\star) \text{ If } \phi \circ f \cong \psi \circ f, \text{ then } \phi \cong \psi.$$

01C6 9.2.8 Corepresentably Conservative Morphisms

Let C be a bicategory.

01C7 DEFINITION 9.2.8.1 ► COREPRESENTABLY CONSERVATIVE MORPHISMS

A 1-morphism $f: A \rightarrow B$ of C is **corepresentably conservative** if, for each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is conservative.

01C8 REMARK 9.2.8.2 ► UNWINDING DEFINITION 9.2.8.1

In detail, f is corepresentably conservative if, for each pair of morphisms $\phi, \psi: B \rightrightarrows X$ and each 2-morphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of C , if the 2-morphism

$$\alpha \star \text{id}_f : \phi \circ f \implies \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \parallel \\ \alpha \star \text{id}_f \\ \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

is a 2-isomorphism, then so is α .

01C9 9.2.9 Strict Epimorphisms

Let C be a bicategory.

01CA DEFINITION 9.2.9.1 ▶ STRICT EPIMORPHISMS

A 1-morphism $f : A \rightarrow B$ is a **strict epimorphism in C** if, for each $X \in \text{Obj}(C)$, the functor

$$f^* : \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is injective on objects, i.e. its action on objects

$$f_* : \text{Obj}(\text{Hom}_C(B, X)) \rightarrow \text{Obj}(\text{Hom}_C(A, X))$$

is injective.

01CB REMARK 9.2.9.2 ▶ UNWINDING DEFINITION 9.2.9.1

In detail, f is a strict epimorphism if, for each diagram in C of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \Downarrow \\ \xrightarrow{\psi} \end{array} X,$$

if $\phi \circ f = \psi \circ f$, then $\phi = \psi$.

01CC EXAMPLE 9.2.9.3 ▶ EXAMPLES OF STRICT EPIMORPHISMS

Here are some examples of strict epimorphisms.

01CD 1. *Strict Epimorphisms in Cats_2 .* The strict epimorphisms in Cats_2 are characterised in **Item 1** of **Proposition 8.6.3.2**.

01CE 2. *Strict Epimorphisms in Rel .* The strict epimorphisms in Rel are charac-

terised in [Proposition 5.3.9.1](#).

01CF 9.2.10 Pseudoepic Morphisms

Let C be a bicategory.

01CG DEFINITION 9.2.10.1 ► PSEUDOEPIC MORPHISMS

A 1-morphism $f: A \rightarrow B$ of C is **pseudoepic** if, for each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is pseudomononic.

01CH REMARK 9.2.10.2 ► UNWINDING DEFINITION 9.2.10.1

In detail, a 1-morphism $f: A \rightarrow B$ of C is pseudoepic if it satisfies the following conditions:

- 01CJ** 1. For all diagrams in C of the form

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \beta \\ \xrightarrow{\psi} \end{array} X,$$

if we have

$$\alpha \star \text{id}_f = \beta \star \text{id}_f,$$

then $\alpha = \beta$.

- 01CK** 2. For each $X \in \text{Obj}(C)$ and each 2-isomorphism

$$\beta: \phi \circ f \xrightarrow{\sim} \psi \circ f, \quad A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of C , there exists a 2-isomorphism

$$\alpha: \phi \xrightarrow{\sim} \psi, \quad B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X$$

of C such that we have an equality

$$A \xrightarrow{f} B \begin{array}{c} \xrightarrow{\phi} \\ \alpha \Downarrow \\ \xrightarrow{\psi} \end{array} X = A \begin{array}{c} \xrightarrow{\phi \circ f} \\ \beta \Downarrow \\ \xrightarrow{\psi \circ f} \end{array} X$$

of pasting diagrams in C , i.e. such that we have

$$\beta = \alpha \star \text{id}_f.$$

01CL

PROPOSITION 9.2.10.3 ► PROPERTIES OF PSEUDOEPIC MORPHISMS

Let $f : A \rightarrow B$ be a 1-morphism of C .

01CM

1. *Characterisations.* The following conditions are equivalent:

01CN

(a) The morphism f is pseudoepic.

01CP

(b) The morphism f is corepresentably full on cores and corepresentably faithful.

01CQ

(c) We have an isococcomma square of the form

$$B \stackrel{\text{eq.}}{\cong} B \amalg_A B, \quad \begin{array}{ccc} B & \xleftarrow{\text{id}_B} & B \\ \text{id}_B \uparrow & \dashrightarrow & \uparrow F \\ B & \xleftarrow{F} & A \end{array}$$

in C up to equivalence.

PROOF 9.2.10.4 ► PROOF OF PROPOSITION 9.2.10.3

Item 1: Characterisations

Omitted.



Appendices

9.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets

Relations

5. Relations

6. Constructions With Relations

7. Equivalence Relations and Apartness Relations

Category Theory

8. Categories

Bicategories

9. Types of Morphisms in Bicategories

Part V

Extra Part

Chapter 10

Miscellaneous Notes

01CR **Contents**

10.1 To Do List.....	540
10.1.1 Omitted Proofs To Add.....	540
10.1.2 Things To Explore/Add.....	541
10.A Other Chapters.....	547

01CS **10.1 To Do List**

01CT **10.1.1 Omitted Proofs To Add**

Не так благотворна истина, как
зловредна ее видимость.

Даниил Данковский

Truth does not do as much good in the
world as the appearance of truth does
evil.

Daniil Dankovsky

There's a very large number of omitted proofs throughout these notes. Here I list them in decreasing order of how nice it would be to add them.

01CU **REMARK 10.1.1.1 ► OMITTED PROOFS TO ADD**

Proofs that *need* to be added at some point:

1. ??.
2. ??.
3. Horizontal composition of natural transformations is associative: ?? of

??.

4. Fully faithful functors are essentially injective: ?? of ??.

Proofs that *would be very nice* to be added at some point:

1. Properties of pseudomonadic functors: ??.
2. Characterisation of fully faithful functors: ?? of ??.

Proofs that *would be nice* to be added at some point:

1. Properties of posetal categories: ??.
2. The quadruple adjunction between categories and sets: ??.
3. Properties of groupoid completions: ??.
4. Properties of cores: ??.
5. F_* faithful iff F faithful: ?? of ??.
6. F_* full iff F full: ?? of ??.
7. Injective on objects functors are precisely the isofibrations in \mathbf{Cats}_2 : ?? of ??.
8. Characterisations of monomorphisms of categories: ?? of ??.
9. Epimorphisms of categories are surjective on objects: ?? of ??.
10. Properties of pseudoepic functors: ??.

01CV 10.1.2 Things To Explore/Add

Here we list things to be explored/added to this work in the future.

01CW REMARK 10.1.2.1 ► THINGS TO EXPLORE/ADD

Set theory through a category theory lens:

1. Isbell duality for sets.
2. Density comonads and codensity monads for sets.

Relations:

1. 2-Categorical monomorphisms and epimorphisms in **Rel**.
2. Co/limits in **Rel**.
3. Apartness composition, categorical properties of **Rel** with apartness, and apartness relations.
4. Apartness defines a composition for relations, but its analogue

$$q \square p \stackrel{\text{def}}{=} \int_{A \in C} p_A^{-1} \amalg q_{-2}^A$$

fails to be unital for profunctors. Is there a less obvious analogue of apartness composition for profunctors?

5. Codensity monad $\text{Ran}_J(J)$ of a relation (What about $\text{Rift}_J(J)$?)
6. Relative comonads in the 2-category of relations
7. Discrete fibrations and Street fibrations in **Rel**.
8. Consider adding the sections
 - The Monoidal Bicategory of Relations
 - The Monoidal Double Category of Relations

to **Relations**.

Spans:

1. Universal property of the bicategory of spans, <https://ncatlab.org/nlab/show/span>
2. Write about cospans.

Un/Straightening:

1. Write proper sections on straightening for lax functors from sets to Rel or Span (displayed sets)

Categories:

1. Expand ?? and add a proof to it.
2. Sections and retractions; retracts, <https://ncatlab.org/nlab/show/retract>.

3. Regular categories: <https://arxiv.org/pdf/2004.08964.pdf>.
4. Are pseudoepic functors those functors whose restricted Yoneda embedding is pseudomonadic and Yoneda preserves absolute colimits?
5. Absolutely dense functors enriched over \mathbb{R}^+ apparently reduce to topological density

Types of Morphisms in Categories:

1. Behaviour in $\text{Fun}(\mathcal{C}, \mathcal{D})$, e.g. pointwise sections vs. sections in $\text{Fun}(\mathcal{C}, \mathcal{D})$.
2. A faithful functor from balanced category is conservative

Yoneda stuff:

1. Properties of restricted Yoneda embedding, e.g. if the restricted Yoneda embedding is full, then what can we conclude? Related: <https://qc.hu.wordpress.com/2015/05/17/generators/>

Adjunctions:

1. Adjunctions, units, counits, and fully faithfulness as in <https://mathoverflow.net/questions/100808/properties-of-functors-and-their-adjoints>.
2. Morphisms between adjunctions and bicategory $\text{Adj}(\mathcal{C})$.
3. <https://ncatlab.org/nlab/show/transformation+of+adjoints>

Constructions With Categories:

1. Comparison between pseudopullbacks and isocomma categories: the “evident” functor $\mathcal{C} \times_{\mathcal{E}}^{\text{ps}} \mathcal{D} \rightarrow \mathcal{C} \overset{\leftrightarrow}{\times}_{\mathcal{E}} \mathcal{D}$ is essentially surjective and full, but not faithful in general.

Co/limits:

1. Add the characterisations of absolutely dense functors given in ?? to ??.
2. Absolutely dense functors, <https://ncatlab.org/nlab/show/absolutely+dense+functor>. Also theorem 1.1 here: <http://www.tac.mta.ca/tac/volumes/8/n20/n20.pdf>.

3. Dense functors, codense functors, and absolutely codense functors.

Co/ends:

1. Examples of co/ends: <https://mathoverflow.net/a/461814>
2. Cofinality for co/ends, <https://mathoverflow.net/questions/353876>

Fibred category theory:

1. Internal **Hom** in categories of co/Cartesian fibrations.
2. *Tensor structures on fibered categories* by Luca Terenzi: <https://arxiv.org/abs/2401.13491>. Check also the other papers by Luca Terenzi.
3. <https://ncatlab.org/nlab/show/cartesian+natural+transformation> (this is a cartesian morphism in $\text{Fun}(C, \mathcal{D})$ apparently)
4. CoCartesian fibration classifying $\text{Fun}(F, G)$, <https://mathoverflow.net/questions/457533/cocartesian-fibration-classifying-mathrmfunf-g>

Monoidal categories:

1. Free braided monoidal category with a braided monoid: <https://ncatlab.org/nlab/show/vine>

Skew monoidal categories:

1. Does the \mathbb{E}_1 tensor product of monoids admit a skew monoidal category structure?
2. Is there a (right?) skew monoidal category structure on $\text{Fun}(C, \mathcal{D})$ using right Kan extensions instead of left Kan extensions?
3. Similarly, are there skew monoidal category structures on the subcategory of $\mathbf{Rel}(A, B)$ spanned by the functions using left Kan extensions and left Kan lifts?

Higher categories:

1. Internal adjunctions in Mod as in [Y21, Section 6.3]; see [Y21, Example 6.2.6].
2. Comonads in the bicategory of profunctors.

Monoids:

1. Isbell's zigzag theorem for semigroups: the following conditions are equivalent:

(a) A morphism $f: A \rightarrow B$ of semigroups is an epimorphism.

(b) For each $b \in B$, one of the following conditions is satisfied:

- We have $f(a) = b$.
- There exist some $m \in \mathbb{N}_{\geq 1}$ and two factorisations

$$b = a_0 y_1,$$

$$b = x_m a_{2m}$$

connected by relations

$$a_0 = x_1 a_1,$$

$$a_1 y_1 = a_2 y_2,$$

$$x_1 a_2 = x_2 a_3,$$

$$a_{2m-1} y_m = a_{2m}$$

such that, for each $1 \leq i \leq m$, we have $a_i \in \text{Im}(f)$.

Wikipedia says in https://en.wikipedia.org/wiki/Isbell%27s_zigzag_theorem:

For monoids, this theorem can be written more concisely:

Types of morphisms in bicategories:

1. Behaviour in 2-categories of pseudofunctors (or lax functors, etc.), e.g. pointwise pseudoepic morphisms in vs. pseudoepic morphisms in 2-categories of pseudofunctors.
2. Statements like “coequifiers are lax epimorphisms”, Item 2 of Examples 2.4 of <https://arxiv.org/abs/2109.09836>, along with most of the other statements/examples there.
3. Dense, absolutely dense, etc. morphisms in bicategories

Other:

1. <https://qchu.wordpress.com/>

2. <https://aroundtoposes.com/>
3. <https://ncatlab.org/nlab/show/essentially+surjective+and+full+functor>
4. <https://mathoverflow.net/questions/415363/objects-who-se-representable-presheaf-is-a-fibration>
5. <https://mathoverflow.net/questions/460146/universal-property-of-isbell-duality>
6. <http://www.tac.mta.ca/tac/volumes/36/12/36-12abs.html> (Isbell conjugacy and the reflexive completion)
7. <https://ncatlab.org/nlab/show/enrichment+versus+internalisation>
8. The works of Philip Saville, <https://philipsaville.co.uk/>
9. https://golem.ph.utexas.edu/category/2024/02/from_cartesian_to_symmetric_mo.html
10. <https://mathoverflow.net/q/463855> (One-object lax transformations)
11. <https://ncatlab.org/nlab/show/analytic+completion+of+a+ring>
12. https://en.wikipedia.org/wiki/Quaternionic_analysis
13. <https://arxiv.org/abs/2401.15051> (The Norm Functor over Schemes)
14. <https://mathoverflow.net/questions/407291/> (Adjunctions with respect to profunctors)
15. <https://mathoverflow.net/a/462726> (Prof is free completion of Cats under right extensions)
16. there's some cool stuff in <https://arxiv.org/abs/2312.00990> (Polynomial Functors: A Mathematical Theory of Interaction), e.g. on cofunctors.
17. <https://ncatlab.org/nlab/show/adjoint+lifting+theorem>
18. <https://ncatlab.org/nlab/show/Gabriel%E2%80%93Ulmer+duality>

Appendices

10.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets

Relations

5. Relations

6. Constructions With Relations

7. Equivalence Relations and Apartness Relations

Category Theory

8. Categories

Bicategories

9. Types of Morphisms in Bicategories

Bibliography

- [MO 119454] user30818. *Category and the axiom of choice*. MathOverflow. URL: <https://mathoverflow.net/q/119454> (cit. on p. 471).
- [MO 321971] Ivan Di Liberti. *Characterization of pseudo monomorphisms and pseudo epimorphisms in Cat*. MathOverflow. URL: <https://mathoverflow.net/q/321971> (cit. on p. 482).
- [MO 350788] Peter LeFanu Lumsdaine. *Epimorphisms of relations*. MathOverflow. URL: <https://mathoverflow.net/q/455260> (cit. on p. 316).
- [MO 382264] Neil Strickland. *Proof that a cartesian category is monoidal*. MathOverflow. URL: <https://mathoverflow.net/q/382264> (cit. on p. 25).
- [MO 460656] Emily de Oliveira Santos. *Existence and characterisations of left Kan extensions and liftings in the bicategory of relations I*. MathOverflow. URL: <https://mathoverflow.net/q/460656> (cit. on pp. 337, 338).
- [MO 461592] Emily de Oliveira Santos. *Existence and characterisations of left Kan extensions and liftings in the bicategory of relations II*. MathOverflow. URL: <https://mathoverflow.net/q/461592> (cit. on pp. 337–339).
- [MO 467527] Emily de Oliveira Santos. *What are the 2-categorical mono/epimorphisms in the 2-category of relations?* MathOverflow. URL: <https://mathoverflow.net/q/467527> (cit. on pp. 313, 320).
- [MO 468121a] Emily de Oliveira Santos. *Characterisations of functors F such that F^* or F_* is [property], e.g. faithful, conservative, etc*. MathOverflow. URL: <https://mathoverflow.net/q/468125> (cit. on pp. 468, 469, 475, 476, 478, 483, 486, 490).
- [MO 468121b] Emily de Oliveira Santos. *Looking for a nice characterisation of functors F whose precomposition functor F^* is full*. MathOverflow. URL: <https://mathoverflow.net/q/468121> (cit. on p. 462).

- [MO 468334] **Emily de Oliveira Santos**. *Is a pseudomonoid and pseudoepimorphism necessarily an equivalence of categories?* MathOverflow. URL: <https://mathoverflow.net/q/468334> (cit. on p. 483).
- [MO 64365] **Giorgio Mossa**. *Natural transformations as categorical homotopies*. MathOverflow. URL: <https://mathoverflow.net/q/64365> (cit. on p. 504).
- [MSE 1465107] **kilian**. *Equivalence of categories and axiom of choice*. Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/1465107> (cit. on p. 471).
- [MSE 2096272] **Akiva Weinberger**. *Is composition of two transitive relations transitive? If not, can you give me a counterexample?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/2096272> (cit. on p. 403).
- [MSE 267469] **Zhen Lin**. *Show that the powerset partial order is a cartesian closed category*. Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/267469> (cit. on p. 63).
- [MSE 2855868] **Qiaochu Yuan**. *Is the category of pointed sets Cartesian closed?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/2855868> (cit. on p. 126).
- [MSE 350788] **Qiaochu Yuan**. *Mono's and epi's in the category Rel?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/350788> (cit. on pp. 308, 315).
- [MSE 733161] **Stefan Hamcke**. *Precomposition with a faithful functor*. Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/733161> (cit. on p. 466).
- [MSE 733163] **Zhen Lin**. *Precomposition with a faithful functor*. Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/733163> (cit. on p. 458).
- [MSE 749304] **Zhen Lin**. *If the functor on presheaf categories given by precomposition by F is ff , is F full? faithful?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/749304> (cit. on p. 466).
- [MSE 884460] **Martin Brandenburg**. *Why are the category of pointed sets and the category of sets and partial functions “essentially the same”?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/884460> (cit. on p. 126).
- [Adá+01] Jiří Adámek, Robert El Bashir, Manuela Sobral, and Jiří Velebil. “On Functors Which Are Lax Epimorphisms”. In: *Theory Appl. Categ.* 8 (2001), pp. 509–521. ISSN: 1201-561X (cit. on pp. 459, 462, 466, 529).

- [Bor94] Francis Borceux. *Handbook of Categorical Algebra I*. Vol. 50. Encyclopedia of Mathematics and its Applications. Basic Category Theory. Cambridge University Press, Cambridge, 1994, pp. xvi+345. ISBN: 0-521-44178-1 (cit. on p. 499).
- [BS10] John C. Baez and Michael Shulman. “Lectures on n -Categories and Cohomology”. In: *Towards higher categories*. Vol. 152. IMA Vol. Math. Appl. Springer, New York, 2010, pp. 1–68. DOI: [10.1007/978-1-4419-1524-5_1](https://doi.org/10.1007/978-1-4419-1524-5_1). URL: https://doi.org/10.1007/978-1-4419-1524-5_1 (cit. on pp. 4, 5, 461).
- [Cie97] Krzysztof Ciesielski. *Set Theory for the Working Mathematician*. Vol. 39. London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1997, pp. xii+236. ISBN: 0-521-59441-3; 0-521-59465-0. DOI: [10.1017/CB09781139173131](https://doi.org/10.1017/CB09781139173131). URL: <https://doi.org/10.1017/CB09781139173131> (cit. on p. 55).
- [DFH75] Aristide Deleanu, Armin Frei, and Peter Hilton. “Idempotent Triples and Completion”. In: *Math. Z.* 143 (1975), pp. 91–104. ISSN: 0025-5874,1432-1823. DOI: [10.1007/BF01173053](https://doi.org/10.1007/BF01173053). URL: <https://doi.org/10.1007/BF01173053> (cit. on p. 475).
- [Fre09] Jonas Frey. *On the 2-Categorical Duals of (Full and) Faithful Functors*. <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=4c289321d622f8fcf947e7a7cfd1bdf75c95ca33>. Archived at <https://web.archive.org/web/20240331195546/https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=4c289321d622f8fcf947e7a7cfd1bdf75c95ca33>. July 2009. URL: <https://citeseerx.ist.psu.edu/document?repid=rep1%5C&type=pdf%5C&doi=4c289321d622f8fcf947e7a7cfd1bdf75c95ca33> (cit. on pp. 459, 466).
- [GGN15] David Gepner, Moritz Groth, and Thomas Nikolaus. “Universality of Multiplicative Infinite Loop Space Machines”. In: *Algebr. Geom. Topol.* 15.6 (2015), pp. 3107–3153. ISSN: 1472-2747. DOI: [10.2140/agt.2015.15.3107](https://doi.org/10.2140/agt.2015.15.3107). URL: <https://doi.org/10.2140/agt.2015.15.3107> (cit. on p. 269).
- [Isb68] John R. Isbell. “Epimorphisms and Dominions. III”. In: *Amer. J. Math.* 90 (1968), pp. 1025–1030. ISSN: 0002-9327,1080-6377. DOI: [10.2307/2373286](https://doi.org/10.2307/2373286). URL: <https://doi.org/10.2307/2373286> (cit. on p. 477).
- [JY21] Niles Johnson and Donald Yau. *2-Dimensional Categories*. Oxford University Press, Oxford, 2021, pp. xix+615. ISBN: 978-0-19-887138-5; 978-0-19-887137-8. DOI: [10.1093/oso/9780198871378.001.0001](https://doi.org/10.1093/oso/9780198871378.001.0001). URL: <https://doi.org/10.1093/oso/9780198871378.001.0001> (cit. on p. 544).

- [Low15] Zhen Lin Low. *Notes on Homotopical Algebra*. Nov. 2015. URL: <https://zll22.user.srcf.net/writing/homotopical-algebra/2015-11-10-Main.pdf> (cit. on p. 466).
- [nLaz24] nLab Authors. *Groupoid*. <https://ncatlab.org/nlab/show/groupoid>. Oct. 2024 (cit. on p. 472).
- [nLab23] nLab Authors. *Skeleton*. 2024. URL: <https://ncatlab.org/nlab/show/skeleton> (cit. on p. 427).
- [Pro24a] ProofWiki Contributors. *Bijection Between $R \times (S \times T)$ and $(R \times S) \times T$ — ProofWiki*. 2024. URL: [https://proofwiki.org/wiki/Bijection_between_R_x_\(S_x_T\)_and_\(R_x_S\)_x_T](https://proofwiki.org/wiki/Bijection_between_R_x_(S_x_T)_and_(R_x_S)_x_T) (cit. on p. 25).
- [Pro24b] Proof Wiki Contributors. *Bijection Between $S \times T$ and $T \times S$ — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Bijection_between_S_x_T_and_T_x_S (cit. on p. 25).
- [Pro24c] Proof Wiki Contributors. *Cartesian Product Distributes Over Set Difference — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Set_Difference (cit. on p. 25).
- [Pro24d] Proof Wiki Contributors. *Cartesian Product Distributes Over Symmetric Difference — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Symmetric_Difference (cit. on p. 25).
- [Pro24e] Proof Wiki Contributors. *Cartesian Product Distributes Over Union — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Union (cit. on p. 25).
- [Pro24f] Proof Wiki Contributors. *Cartesian Product Is Empty Iff Factor Is Empty — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Cartesian_Product_is_Empty_iff_Factor_is_Empty (cit. on p. 25).
- [Pro24g] Proof Wiki Contributors. *Cartesian Product of Intersections — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Cartesian_Product_of_Intersections (cit. on p. 25).
- [Pro24h] Proof Wiki Contributors. *Characteristic Function of Intersection — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Intersection (cit. on p. 63).
- [Pro24i] Proof Wiki Contributors. *Characteristic Function of Set Difference — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Set_Difference (cit. on p. 67).

- [Pro24j] Proof Wiki Contributors. *Characteristic Function of Symmetric Difference*—Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Symmetric_Difference (cit. on p. 75).
- [Pro24k] Proof Wiki Contributors. *Characteristic Function of Union*—Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Union (cit. on p. 59).
- [Pro24l] Proof Wiki Contributors. *Complement of Complement*—Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Complement_of_Complement (cit. on p. 70).
- [Pro24m] Proof Wiki Contributors. *Condition For Mapping from Quotient Set To Be A Surjection*—Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Surjection (cit. on p. 413).
- [Pro24n] Proof Wiki Contributors. *Condition For Mapping From Quotient Set To Be An Injection*—Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Injection (cit. on p. 413).
- [Pro24o] Proof Wiki Contributors. *Condition For Mapping From Quotient Set To Be Well-Defined*—Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Well-Defined (cit. on p. 413).
- [Pro24p] Proof Wiki Contributors. *De Morgan's Laws (Set Theory)*—Proof Wiki. 2024. URL: [https://proofwiki.org/wiki/De_Morgan%27s_Laws_\(Set_Theory\)](https://proofwiki.org/wiki/De_Morgan%27s_Laws_(Set_Theory)) (cit. on pp. 67, 70).
- [Pro24q] Proof Wiki Contributors. *De Morgan's Laws (Set Theory)/Set Difference/Difference with Union*—Proof Wiki. 2024. URL: [https://proofwiki.org/wiki/De_Morgan%27s_Laws_\(Set_Theory\)/Set_Difference/Difference_with_Union](https://proofwiki.org/wiki/De_Morgan%27s_Laws_(Set_Theory)/Set_Difference/Difference_with_Union) (cit. on p. 67).
- [Pro24r] Proof Wiki Contributors. *Equivalence of Definitions of Symmetric Difference*—Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Equivalence_of_Definitions_of_Symmetric_Difference (cit. on p. 74).
- [Pro24s] Proof Wiki Contributors. *Image of Intersection Under Mapping*—Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Image_of_Intersection_under_Mapping (cit. on p. 103).
- [Pro24t] Proof Wiki Contributors. *Image of Union Under Mapping*—Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Image_of_Union_under_Mapping (cit. on p. 104).

- [Pro24u] Proof Wiki Contributors. *Intersection Distributes Over Symmetric Difference*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Intersection_Distributes_over_Symmetric_Difference (cit. on p. 75).
- [Pro24v] Proof Wiki Contributors. *Intersection Is Associative*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Intersection_is_Associative (cit. on p. 63).
- [Pro24w] Proof Wiki Contributors. *Intersection Is Commutative*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Intersection_is_Commutative (cit. on p. 63).
- [Pro24x] Proof Wiki Contributors. *Intersection With Empty Set*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Intersection_with_Empty_Set (cit. on p. 63).
- [Pro24y] Proof Wiki Contributors. *Intersection With Set Difference Is Set Difference With Intersection*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Intersection_with_Set_Difference_is_Set_Difference_with_Intersection (cit. on p. 67).
- [Pro24z] Proof Wiki Contributors. *Intersection With Subset Is Subset*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Intersection_with_Subset_is_Subset (cit. on p. 63).
- [Pro24aa] Proof Wiki Contributors. *Mapping From Quotient Set When Defined Is Unique*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Mapping_from_Quotient_Set_when_Defined_is_Unique (cit. on p. 413).
- [Pro24ab] Proof Wiki Contributors. *Preimage of Intersection Under Mapping*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Preimage_of_Intersection_under_Mapping (cit. on p. 108).
- [Pro24ac] Proof Wiki Contributors. *Preimage of Union Under Mapping*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Preimage_of_Union_under_Mapping (cit. on p. 108).
- [Pro24ad] Proof Wiki Contributors. *Quotient Mapping Is Coequalizer*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Quotient_Mapping_is_Coequalizer (cit. on p. 52).
- [Pro24ae] Proof Wiki Contributors. *Set Difference as Intersection With Complement*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Set_Difference_as_Intersection_with_Complement (cit. on p. 67).
- [Pro24af] Proof Wiki Contributors. *Set Difference as Symmetric Difference With Intersection*—ProofWiki. 2024. URL: https://proofwiki.org/wiki/Set_Difference_as_Symmetric_Difference_With_Intersection

- [org/wiki/Set_Difference_as_Symmetric_Difference_with_Intersection](https://proofwiki.org/wiki/Set_Difference_as_Symmetric_Difference_with_Intersection) (cit. on p. 67).
- [Pro24ag] Proof Wiki Contributors. *Set Difference Is Right Distributive Over Union* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_is_Right_Distributive_over_Union (cit. on p. 67).
- [Pro24ah] Proof Wiki Contributors. *Set Difference Over Subset* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_over_Subset (cit. on p. 67).
- [Pro24ai] Proof Wiki Contributors. *Set Difference With Empty Set Is Self* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_with_Empty_Set_is_Self (cit. on p. 67).
- [Pro24aj] Proof Wiki Contributors. *Set Difference With Self Is Empty Set* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_with_Self_is_Empty_Set (cit. on p. 67).
- [Pro24ak] Proof Wiki Contributors. *Set Difference With Set Difference Is Union of Set Difference With Intersection* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_with_Set_Difference_is_Union_of_Set_Difference_with_Intersection (cit. on p. 67).
- [Pro24al] Proof Wiki Contributors. *Set Difference With Subset Is Superset of Set Difference* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_with_Subset_is_Superset_of_Set_Difference (cit. on p. 67).
- [Pro24am] Proof Wiki Contributors. *Set Difference With Union* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_with_Union (cit. on p. 67).
- [Pro24an] Proof Wiki Contributors. *Set Intersection Distributes Over Union* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Intersection_Distributes_over_Union (cit. on p. 63).
- [Pro24ao] Proof Wiki Contributors. *Set Intersection Is Idempotent* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Set_Intersection_is_Idempotent (cit. on p. 63).
- [Pro24ap] Proof Wiki Contributors. *Set Intersection Preserves Subsets* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Set_Intersection_Preserves_Subsets (cit. on p. 63).
- [Pro24aq] Proof Wiki Contributors. *Set Union Is Idempotent* — *Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Set_Union_is_Idempotent (cit. on p. 59).

- [Pro24ar] Proof Wiki Contributors. *Set Union Preserves Subsets*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Union_Preserves_Subsets (cit. on p. 59).
- [Pro24as] Proof Wiki Contributors. *Symmetric Difference Is Associative*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_is_Associative (cit. on p. 74).
- [Pro24at] Proof Wiki Contributors. *Symmetric Difference Is Commutative*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_is_Commutative (cit. on p. 74).
- [Pro24au] Proof Wiki Contributors. *Symmetric Difference of Complements*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_of_Complements (cit. on p. 74).
- [Pro24av] Proof Wiki Contributors. *Symmetric Difference on Power Set Forms Abelian Group*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_on_Power_Set_forms_Abelian_Group (cit. on p. 75).
- [Pro24aw] Proof Wiki Contributors. *Symmetric Difference With Complement*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Complement (cit. on p. 74).
- [Pro24ax] Proof Wiki Contributors. *Symmetric Difference With Empty Set*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Empty_Set (cit. on p. 74).
- [Pro24ay] Proof Wiki Contributors. *Symmetric Difference With Intersection Forms Ring*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Intersection_forms_Ring (cit. on p. 75).
- [Pro24az] Proof Wiki Contributors. *Symmetric Difference With Self Is Empty Set*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Self_is_Empty_Set (cit. on p. 74).
- [Pro24ba] Proof Wiki Contributors. *Symmetric Difference With Union Does Not Form Ring*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Union_does_not_form_Ring (cit. on p. 74).
- [Pro24bb] Proof Wiki Contributors. *Symmetric Difference With Universe*—*ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Universe (cit. on p. 74).
- [Pro24bc] Proof Wiki Contributors. *Union as Symmetric Difference With Intersection*—*ProofWiki*. 2024. URL: <https://proofwiki.org/>

- wiki/Union_as_Symmetric_Difference_with_Intersection (cit. on p. 59).
- [Pro24bd] Proof Wiki Contributors. *Union Distributes Over Intersection*—*Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_Distributes_over_Intersection (cit. on p. 59).
- [Pro24be] Proof Wiki Contributors. *Union Is Associative*—*Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_is_Associative (cit. on p. 59).
- [Pro24bf] Proof Wiki Contributors. *Union Is Commutative*—*Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_is_Commutative (cit. on p. 59).
- [Pro24bg] Proof Wiki Contributors. *Union of Symmetric Differences*—*Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_of_Symmetric_Differences (cit. on p. 74).
- [Pro24bh] Proof Wiki Contributors. *Union With Empty Set*—*Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_with_Empty_Set (cit. on p. 59).
- [PS19] Maximilien Péroux and Brooke Shipley. “Coalgebras in Symmetric Monoidal Categories of Spectra”. In: *Homology Homotopy Appl.* 21.1 (2019), pp. 1–18. ISSN: 1532-0073. DOI: [10.4310/HHA.2019.v21.n1.a1](https://doi.org/10.4310/HHA.2019.v21.n1.a1). URL: <https://doi.org/10.4310/HHA.2019.v21.n1.a1> (cit. on p. 269).
- [Rie17] Emily Riehl. *Category Theory in Context*. Vol. 10. Aurora: Dover Modern Math Originals. Courier Dover Publications, 2017, pp. xviii+240. ISBN: 978-0486809038. URL: <http://www.math.jhu.edu/~eriehl/context.pdf> (cit. on p. 472).
- [Wik24] Wikipedia Contributors. *Multivalued Function*—*Wikipedia, The Free Encyclopedia*. 2024. URL: https://en.wikipedia.org/wiki/Multivalued_function (cit. on p. 282).

Index of Notation

A

$[a]$, 409
 (A, B) , 55
 $A \amalg B$, 39
 $A \amalg_C B$, 42
 αF , 499
 $\alpha \star F$, 499
 $A \odot X$, 166
 $a \odot x$, 168
 $A \odot (X, x_0)$, 166
 $A \multimap X$, 174
 $A \multimap (X, x_0)$, 174
 $A \triangle B$, 70
 $A \times B$, 17
 $A \times_C B$, 25
 $A \times_{f,C,g} B$, 26
 $A \cup B$, 56

B

$\beta \circ \alpha$, 494
 $\beta \star \alpha$, 497

C

$C(A, B)$, 420
Cats, 510
Cats₂, 511
 $[C, \mathcal{D}]$, 507
 $\chi(-)$, 76
 χ_{-2}^{-1} , 77
 χ_U , 76
 χ^x , 77

χ_x , 76
 $\chi_X(-1, -2)$, 76
 $\chi_X(-, U)$, 77
 $\chi_X(-, x)$, 77
 $\chi_X(U, -)$, 77
 $\chi_X(x, -)$, 77
CoEq(f, g), 50
Coim(f), 412
Coll(R), 364
Coll(R), 364
 $\coprod_{i \in I} A_i$, 37
Core(C), 443
 C^{\approx} , 443

D

\mathcal{D}^C , 507
dom(R), 349

E

\emptyset_{cat} , 422
 \emptyset , 36, 54
Eq(f, g), 32
eq(f, g), 32

F

$F\alpha$, 499
 f^{-1} , 105
 $f^{-1}(V)$, 105
 \mathbb{F}_1 , 121
 $f!$, 110
 $f!_{\text{cp}}(U)$, 112

$f_{!,im}(U)$, 111
 $f_!(U)$, 110
 f^* , 99, 429
 $F \star \alpha$, 499
 $\alpha \star F$, 499
 $f(U)$, 99
 $\text{Fun}(C, \mathcal{D})$, 506

G

$\text{Gr}(A)$, 342
 Grpd , 512
 Grpd_2 , 512

H

$\text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z)$, 163
 $\text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z)$, 164
 $\text{Hom}_{\text{Sets}_*}^{\otimes}(X \times Y, Z)$, 166
 $\text{Hom}_{\text{Sets}}(A, B)$, 55
 $\text{Hom}_{\text{Sets}_*}(X, Y)$, 245

I

id_C , 448
 id_F , 493
 $\text{id}_F \star \alpha$, 499
 $\bigcup_{i \in I} R_i$, 353
 $\bigcap_{X \in \mathcal{F}} X$, 59
 $\text{Iso}_C(A, B)$, 439

K

$K_0(C)$, 439

L

\triangleleft , 182
 $[-, -]_{\text{Sets}_*}^{\triangleleft}$, 187
 $\triangleleft_{\text{Sets}_*}$, 183

M

$\text{Mor}(C)$, 420

N

n , 422
 $\text{Nat}(F, G)$, 493

P

$\pi_0(C)$, 433
 \mathcal{P}^{-1} , 84, 391
 \star_X , 154
 $\prod_{i \in I} A_i$, 14, 127
 $\mathcal{P}!$, 85, 391
 \mathcal{P}_* , 83, 391
 pt , 13, 121, 126, 139
 pt , 421
 $\mathcal{P}(X)$, 82

R

$\text{range}(R)$, 349
 R^\dagger , 357
 Rel , 284
 Rel , 290
 $\text{Rel}(A, B)$, 275
 $\text{Rel}(A, B)$, 275
 Rel^{dbl} , 291
 $\text{Rel}^{\text{eq}}(A, B)$, 406
 $\text{Rel}^{\text{eq}}(A, B)$, 406
 $\text{Rel}^{\text{refl}}(A, A)$, 396
 $\text{Rel}^{\text{refl}}(A, A)$, 396
 $\text{Rel}^{\text{symm}}(A, A)$, 399
 $\text{Rel}^{\text{symm}}(A, A)$, 399
 $\text{Rel}^{\text{trans}}(A)$, 402
 $\text{Rel}^{\text{trans}}(A)$, 402
 R^{eq} , 407
 \triangleright , 208
 $[-, -]_{\text{Sets}_*}^{\triangleright}$, 212
 $\triangleright_{\text{Sets}_*}$, 208
 $R \cap S$, 352
 R^{-1} , 378
 R_{-1} , 372
 $R^{-1}(V)$, 378
 $R_{-1}(V)$, 372
 R^{refl} , 397

$R!$, 385
 $R!(U)$, 385
 R^* , 366
 R^{symm} , 400
 $R \times S$, 354
 R^{trans} , 403
 $R(U)$, 366
 $R \cup S$, 349

S

$S \diamond R$, 359
 $\text{Sets}(A, B)$, 56
 Sets_* , 122, 123
 $\text{Sets}_*((X, x_0), (Y, y_0))$, 245
 \sim_{cotriv} , 280
 \sim_{id} , 77
 \sim_{refl} , 397
 \sim_R^{symm} , 400
 \sim_R^{eq} , 407
 \sim_R^{trans} , 403
 \sim_{triv} , 279
 S^0 , 121

T

$\{t, f\}$, 5
 $\prod_{i \in I} R_i$, 356
 $\text{Trans}(F, G)$, 492
 $\{\text{true}, \text{false}\}$, 5

U

U^c , 69
 $\bigcup_{i \in I} A_i$, 56
 $\bigcup_{i \in I} R_i$, 351

W

忘, 453
 $\bigwedge_{i \in I} X_i$, 270

X

$\{X\}$, 54
 X_{disc} , 435
 X_{indisc} , 437
 $X \cap Y$, 60
 $x \triangleleft_{\text{Sets}_*} y$, 184
 $x \triangleleft y$, 184
 $X \setminus Y$, 64
 X / \sim_R , 409
 $X \otimes_{\mathbb{F}_1} Y$, 233
 X^+ , 154
 X_{pos} , 424
 $x \triangleright y$, 209
 $x \triangleright_{\text{Sets}_*} y$, 209
 $x \wedge y$, 236
 $X \wedge Y$, 233
 $\{X, Y\}$, 55

Index of Set Theory

B

- bilinear morphism
 - of pointed sets, 164
- bilinear morphism of pointed sets
 - left, 163
 - right, 164
- binary intersection, 60

C

- Cartesian product, 15, 17
- category of relations, 284
 - associator of, 285
 - internal Hom of, 288
 - left unitor of, 286
 - monoidal product of, 284
 - monoidal unit of, 285
 - right unitor of, 287
 - symmetry of, 287
- characteristic embedding, 76
- characteristic function
 - of a set, 76
 - of an element, 76
- characteristic relation, 76
- coequaliser of sets, 50
- coimage, 412
- complement of a set, 69
- coproduct
 - of a family of pointed sets, 140

D

- difference of sets, 64

- disjoint union, 39
 - of a family of sets, 37

E

- empty set, 54
- equaliser of sets, 32
- equivalence class, 409
- equivalence relation
 - kernel, 406

F

- fibre coproduct of sets, 43
- fibre product of sets, 26
- field with one element
 - module over, 121
 - module over, morphism of, 122
- function, 2
 - associated direct image
 - function, 99
 - associated direct image with compact support function, 110
 - associated direct image with compact support function, complement part, 111
 - associated direct image with compact support function, image part, 111
 - associated inverse image
 - function, 105
 - coimage of, 412
 - graph of, 342

inverse of, 346
kernel of, 406

I

indicator function, *see* characteristic function
initial pointed set, 139
initial set, 36
intersection of a family of sets, 59

L

left tensor product of pointed sets
 diagonal, 198
 left skew unit of, 189
 skew associator, 189
 skew left unitor, 193
 skew right unitor, 196

M

(−1)-category, 4
(−2)-category, 4
module
 underlying pointed set of, 122

O

ordered pairing, 55

P

pairing of two sets, 55
pointed function, *see* pointed set, morphism of
pointed set, 121
 category of, 122
 coequaliser of, 152
 copower by a set, *see* pointed set, tensor by a set
 coproduct of, 142
 cotensor by a set, 174
 equaliser of, 137
 \mathbb{F}_1 , 121

free, 154
internal Hom of, 245
left internal Hom of, 187
left tensor product, 182
morphism of, 122
of morphisms of pointed sets, 245
power by a set, *see* pointed set, cotensor by a set
product of, 129
pullback of, 132
pushout of, 146
right internal Hom of, 212
right tensor product, 208
tensor by a set, 166
trivial, 121
underlying a module, 122
underlying a semimodule, 122
wedge sum, 142
wedge sum of a family of pointed sets, 140
0-sphere, 121
poset
 of (−1)-categories, 5
 of truth values, 5
powerset, 82
product of a family of pointed sets, 127
product of a family of sets, 14
product of sets, 17
pullback of sets, 25
pushout of sets, 42

Q

quotient
 by an equivalence relation, 409

R

relation, 274
 associated direct image function, 366

- associated direct image with compact support function, 385
- associated strong inverse image function, 372
- associated weak inverse image function, 378
- category of, 275
- category of, 284
- closed symmetric monoidal category of, 289
- collage of, 364
- composition of, 359
- corepresentable, 348
- cotrivial, 280
- domain of, 349
- double category of, 291
- equivalence closure of, 407
- equivalence relation, 406
- equivalence, set of, 406
- functional, 282
- intersection of, 352
- intersection of a family of, 353
- inverse of, 357
- on powersets associated to a relation, 392
- partial equivalence relation, 406
- poset of, 275, 399, 402, 406
- product of, 354
- product of a family of, 356
- range of, 349
- reflexive, 396
- reflexive closure of, 397
- reflexive, poset of, 396
- reflexive, set of, 396
- representable, 348
- set of, 275
- symmetric, 399
- symmetric closure of, 400
- symmetric, set of, 399
- total, 283
- transitive, 402

- transitive closure of, 403
- transitive, set of, 402
- trivial, 279
- 2-category of, 290
- union of, 349
- union of a family of, 351
- right tensor product of pointed sets
 - diagonal, 223
 - right skew unit of, 215
 - skew associator, 215
 - skew left unitor, 218
 - skew right unitor, 220

S

- semimodule
 - underlying pointed set of, 122
- set
 - Hom, 56
 - of maps, 55
- set of bilinear morphisms of pointed sets, 166
 - left, 163
 - right, 164
- singleton set, 54
- smash product
 - of a family of pointed sets, 270
 - of pointed sets, 233
- smash product of pointed sets
 - associator, 247
 - diagonal, 259
 - left unitor, 251
 - monoidal unit of, 247
 - right unitor, 254
 - symmetry, 257
- symmetric difference of sets, 70

T

- terminal pointed set, 126
- terminal set, 13

U

union, 56
of a family of sets, 56

Z

0-category, 8
0-groupoid, 9

Index of Category Theory

A

adjoint equivalence of categories,
470

C

category, 418
adjoint equivalence of, 470
connected, 435
connected component, 432
connected component, set of,
433
core of, 443
disconnected, 435
discrete, 435
empty, 422
equivalence of, 469
groupoid completion of, 439
indiscrete, 437
isomorphism of, 472
 κ -small, 420
locally essentially small, 420
locally small, 420
of small groupoids, 512
skeletal, 427
skeleton of, 427
small, 420
thin, 425
category of categories, 510
contravariant functor, 450

D

discrete category
on a set, 435

E

equivalence of categories, 469

F

forgetful functor, 452
functor, 446
bijective on objects, 484
bo, 484
composition of, 449
conservative, 466
contravariant, 450
corepresentably faithful on
cores, 489
corepresentably full on cores,
489
corepresentably fully faithful on
cores, 490
dominant, 474
epimorphism, 476
eso, 469
essentially injective, 468
essentially surjective, 469
faithful, 456
forgetful, 452
full, 459
fully faithful, 462
identity, 448
injective on objects, 483
monomorphism, 475

pseudoepic, 481
pseudomonadic, 478
representably faithful on cores,
485
representably full on cores, 485
representably fully faithful on
cores, 486
surjective on objects, 484
transformation of, 492
functor category, 506

G

Godement product, *see* natural
transformation, horizontal
composition
Grothendieck groupoid
of a category, 440
groupoid, 439
groupoid completion, 439

I

indiscrete category
on a set, 437
Isbell's zigzag theorem, 477
isomorphism, 439
isomorphism of categories, 472

M

middle four exchange
in Cats, 496, 501

N

natural isomorphism, 505
natural transformation, 492
associated to a functor, 454
equality of, 493
horizontal composition, 497
identity natural transformation,
493

vertical composition, 494

O

ordinal category, 422

P

posetal category, 425
associated to a poset, 424
postcomposition, 429
precomposition, 429
punctual category, 421

S

singleton category, *see* punctual
category
subcategory, 425
full, 426
lluf, 426
strictly full, 426
wide, 426

T

transformation between functors,
492
2-category
of small categories, 511
of small groupoids, 512

W

whiskering
left, 499
right, 499

Z

(0, 1)-category, 425

Index of Higher Category Theory

D

double category
of relations, 291

E

epimorphism
strict, 535

M

monomorphism
strict, 524

O

1-morphism
corepresentably conservative,
534
corepresentably essentially
injective, 534
corepresentably faithful, 527

corepresentably faithful on
cores, 531
corepresentably full, 528
corepresentably full on cores,
531
corepresentably fully faithful,
529
corepresentably fully faithful on
cores, 532
pseudoepic, 536
pseudomononic, 525
representably conservative, 523
representably essentially
injective, 523
representably faithful, 516
representably faithful on cores,
520
representably full, 517
representably full on cores, 520
representably fully faithful, 518
representably fully faithful on
cores, 521